

**RESPONSE OF GRASSPEA TO THE APPLICATION OF PRILLED
UREA AND UREA SUPER GRANULE**

ABDULLAH HEL MAFI



**DEPARTMENT OF AGRONOMY
SHER-E-BANGLA AGRICULTURAL UNIVERSITY
DHAKA-1207**

JUNE, 2014

**RESPONSE OF GRASSPEA TO THE APPLICATION OF PRILLED
UREA AND UREA SUPER GRANULE**

BY

ABDULLAH HEL MAFI

REG. NO. : 07-02428

A Thesis
*Submitted to the Faculty of Agriculture
Sher-e-Bangla Agricultural University, Dhaka
in partial fulfilment of the requirements
for the degree
of*

MASTER OF SCIENCE (MS)

IN

AGRONOMY

SEMESTER: JANUARY-JUNE, 2014

APPROVED BY:

Prof. Dr. Md. Fazlul Karim

Supervisor


Sheikh Muhammad Masum

Co-Supervisor

Prof. Dr. H. M. M. Tariq Hossain

Chairman

Examination Committee



*DEDICATED
TO
MY BELOVED PARENTS*



DEPARTMENT OF AGRONOMY
Sher-e-Bangla Agricultural University
Sher-e-Bangla Nagar, Dhaka-1207

CERTIFICATE

This is to certify that the thesis entitled '**Response of Grasspea to the Application of Prilled Urea and Urea Super Granule**' submitted to the Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka, in partial fulfilment of the requirements for the degree of **Master of Science (MS) in Agronomy**, embodies the result of a piece of bonafide research work carried out by **Abdullah Hel Mafi**, Registration number: **07-02428** under my supervision and guidance. No part of the thesis has been submitted for any other degree or diploma.

I further certify that any help or source of information, received during the course of this investigation, has duly been acknowledged.

Dated:

Dhaka, Bangladesh

Prof. Dr. Md. Fazlul Karim

Supervisor

Department of Agronomy
Sher-e-Bangla Agricultural University
Dhaka-1207

ACKNOWLEDGEMENTS

The author praises to the Kindful Omnipotent “Allah” for His never-ending blessing on to who deems it a great pleasure to express profound gratefulness to “Allah” for successful completion of the research studies followed by this thesis manuscript.

*The author feels proud to express his heartiest scense of gratitude, sincere appreciation and immense indebtedness to his supervisor Professor **Dr. Md. Fazlul Karim**, Department of Agronomy, Shis-e-Bangla Agricultural University (SAU), Dhaka, for his continuous scholastic and intellectual guidance, cooperation, constructive criticism and suggestions in carrying out the research work and preparation of thesis.*

*The author feels proud to express his deepest respect, sincere appreciation and immense indebtedness to his co-supervisor **Sheikh Muhammad Masum**, Assistant Profeser, Department of Agronomy, SAU, Dhaka, for his intellectionalism guidance and valuable suggestions during the entire period of research work and preparation of this thesis.*

*The author expresses his sincere respect to the Professor **Dr. H. M. M. Tariq Hossain**, Chairman, Departement of Agronomy, SAU, Dhaka for valuable suggestions and cooperation during the study period. The author also expresses his heartfelt thanks to all the teachers of the Department of Agronomy, SAU, for their valuable teaching, suggestions and encouragement during the period of the study.*

The author expresses his sincere appreciation to his wife, brothis, sisters, relatives, well wishiss and friends for their inspiration, help and encouragement throughout the study.

The respected parents of author are to be grately reserved for their hadrship look after the author continuously from day one is this stage.

The Author

RESPONSE OF GRASSPEA TO THE APPLICATION OF PRILLED UREA AND UREA SUPER GRANULE

BY

ABDULLAH HEL MAFI

ABSTRACT

The experiment was conducted at the farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka during the period from November 2012 to March, 2013 to study the response of Grasspea to the application of prilled urea and urea super granules. The variety BARI Grasspea 3 was used as test crop. The experiment consists of the following treatments: T₁= Prilled urea (PU) broadcasted; T₂ = PU given between two rows; T₃ = Urea Super Granules (USG) placed at 10 cm distance (avoid one row); T₄ = USG placed at 10 cm distance (avoid two rows); T₅ = USG placed at 10 cm distance (avoid three rows); T₆ = USG placed at 20 cm distance (avoid one row); T₇ = USG placed at 20 cm distance (avoid two rows); T₈ = USG placed at 20 cm distance (avoid three rows); T₉= USG placed at 30 cm distance (avoid one row); T₁₀ = USG placed at 30 cm distance (avoid two rows); T₁₁ = USG placed at 30 cm distance (avoid three rows); T₁₂ = USG placed at 40 cm distance (avoid one row); T₁₃ = USG placed at 40 cm distance (avoid two rows) and T₁₄ = USG placed at 40 cm distance (avoid three rows). USG was placed at 10 cm depth at each case. The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. Results showed that tallest plant 34.64, 40.30, 47.35, 59.32 and 65.49 cm were obtained from T₇ i.e. USG placed at 20 cm distance (avoid two rows) and the shortest plant 19.34, 27.30, 30.55, 38.11 and 57.01 cm from T₁₄ treatment i.e. USG placed at 40 cm distance (avoid three rows) at 30, 50, 70, 90 DAS and harvest, respectively. Treatment T₇ also took minimum days (45.33 days) to flowering and it was statistically identical with T₅ (47 days), T₆ (48 days) and T₁ (49.67 days), while T₁₄ treatment took maximum days (57.67 days). The highest pods plant⁻¹ (54.48), seeds pod⁻¹ (4.87), weight of 1000 seeds (40.57 g), seed yield (1.94 t ha⁻¹) and stover yield (2.92 t ha⁻¹) was recorded from T₇, while the lowest number of pods plant⁻¹ (45.09), lowest number of seeds pod⁻¹ (3.43), lowest weight of 1000 seeds (34.79 g), lowest seed yield (1.19 t ha⁻¹) and lowest stover yield (2.11 t ha⁻¹) was observed from T₁₄. The study revealed that USG application showed better performance than prilled when placed in two rows but not with broadcasted prilled urea.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE NO.
	ACKNOWLEDGEMENTS	i
	ABSTRACT	ii
	LIST OF CONTENTS	iii
	LIST OF TABLES	v
	LIST OF FIGURES	v
	LIST OF APPENDICES	vi
I	INTRODUCTION	01
II	REVIEW OF LITERATURE	04
	2.1 Effects of prilled urea on different pulse crop	04
	2.2 Effect of urea super granules on different crops	22
III	MATERIALS AND METHODS	30
	3.1 Experimental site	30
	3.2 Climatic condition	30
	3.3 Characteristics of soil	30
	3.4 Planting material	31
	3.5 Land preparation	31
	3.6 Fertilizer application	31
	3.7 Treatments of the experiment	32
	3.8 Experimental design and layout	32
	3.9 Sowing of seeds in the field	34
	3.10 Intercultural operations	34
	3.11 Crop sampling and data collection	34
	3.12 Harvest and post harvest operations	34
	3.13 Data collection	35
	3.14 Statistical analysis	37

CHAPTER	TITLE	PAGE NO.
IV	RESULTS AND DISCUSSION	38
	4.1 Crop growth characters of chickpea	38
	4.1.1 Plant height	38
	4.1.2 Branches plant ⁻¹	41
	4.1.3 Dry matter plant ⁻¹	43
	4.1.4 Crop growth rate	45
	4.1.5 Relative growth rate	45
	4.1.6 Days from emergence to flowering	45
	4.2 Yield contributing characters and yield of chickpea	45
	4.2.1 Pods plant ⁻¹	49
	4.2.2 Seeds pod ⁻¹	49
	4.2.3 Weight of 1000 seeds	49
	4.2.4 Seed yield ha ⁻¹	50
	4.2.5 Stover yield ha ⁻¹	52
	4.2.6 Biological yield ha ⁻¹	52
	4.2.7 Harvest index	52
V	SUMMARY AND CONCLUSION	54
	REFERENCES	57
	APPENDICES	69

LIST OF TABLES

	TITLE	PAGE NO.
Table 1.	Effect of application of prilled urea and urea super granule on plant height of grasspea at different growth stages	40
Table 2.	Effect of application of prilled urea and urea super granule on branches plant ⁻¹ of grasspea at different growth stages	42
Table 3.	Effect of application of prilled urea and urea super granule on dry matter plant ⁻¹ of grasspea at different growth stages	44
Table 4.	Effect of application of prilled urea and urea super granule on crop growth rate of grasspea	46
Table 5.	Effect of application of prilled urea and urea super granule on relative growth rate of grasspea	47
Table 6.	Effect of application of prilled urea and urea super granule on yield contributing characters and yield of grasspea	51

LIST OF FIGURES

	TITLE	PAGE NO.
Figure 1.	Field layout of the experimental plot	33
Figure 2.	Effect of application of prilled urea and urea super granules on days from emergence to flowering of grasspea	48
Figure 3.	Effect of application of prilled urea and urea super granules on biological yield of grasspea	53

LIST OF APPENDICES

	TITLE	PAGE NO.
Appendix I.	Monthly average of air temperature, relative humidity and total rainfall of the experimental site during the period from November, 2012 to March, 2013	69
Appendix II.	Characteristics of soil of experimental field	69
Appendix III.	Analysis of variance of the data on plant height of grasspea as influenced by application of prilled urea and urea super granule	70
Appendix IV.	Analysis of variance of the data on number of branches plant ⁻¹ of grasspea as influenced by application of prilled urea and urea super granule	70
Appendix V.	Analysis of variance of the data on dry matter content plant ⁻¹ of grasspea as influenced by application of prilled urea and urea super granule	71
Appendix VI.	Analysis of variance of the data on crop growth rate and relative growth rate of grasspea as influenced by application of prilled urea and urea super granule	71
Appendix VII.	Analysis of variance of the data on yield contributing characters, yields and harvest index of grasspea as influenced by application of prilled urea and urea super granule	72

CHAPTER I

INTRODUCTION

Grasspea (*Lathyrus sativus* L.) is an important pulse crop of Bangladesh and is commonly known as khesari. It belongs to the family Fabaceae. This is also an important pulse crop in India, Nepal, Pakistan, China, Middle East, Myanmar, Srilanka, Southern Europe and part of Africa and South America (Tadesse, 1977). It is the most important pulse crop not in terms of area (65,341 ha) and production (56,672 ton) but also for its high consumption as a common pulse in Bangladesh (BBS, 2013).

As a cheap source of easily digestible dietary protein pulse crop play an important role, it provides better nourishment of human body which also complements the staple rice food. According to the FAO (2006) yield of grasspea in Bangladesh is miserably low (663 kg ha⁻¹) as compared to that of other countries like India (789 kg ha⁻¹), Myanmar (1,054 kg ha⁻¹), Mexico (1,534 kg ha⁻¹), Israel (1,634 kg ha⁻¹), Russian Federation (2,423 kg ha⁻¹), Kazakjasthan (3,127 kg ha⁻¹) and China (5,900 kg ha⁻¹). Such low yield in Bangladesh, however is not an indication of low yielding potentiality of this crop, but may be attributed to a number of reasons, viz. unavailability of quality seeds of high yielding varieties, delayed sowing after the harvest of aman rice, inadequate fertilization, poor managements of pests etc. Among all application of nitrogen fertilizer may get important consideration. Pulse crop stop to nourish *Rhizobia* during its reproductive stage rather translocate energy towards development of flowers and pods. Being pulse crop, grasspea needs continuous uptake of nitrogen from vegetative stage to flowering. Otherwise available low nitrogen from basal application of nitrogen eventually hampers the development of reproductive traits. In this situation nitrogen given as basal to the crop is not sufficiently available to the plant for nourishing its flowers and pods thus seed yield is low (Patel *et al.*, 1984; BARC, 2005). Keeping this in mind attention is given on nitrogen placement or use of

Urea Super Granule (USG) in mungbean. Both these ways of nitrogen placement might have some influencing technique that would be better utilization of nitrogen by the mungbean for its maximum seed yield (Bhardwaj and Singh, 1993).

Grasspea plant can fix nitrogen by symbiotic process with *rhizobium* bacteria. At the time of flowering, nodules generally lose their ability to fix nitrogen, because the plant feeds the developing seed rather than nodule. Legume nodules that are no longer fixing nitrogen usually turn green may actually be discarded by the plant. The basal application of urea is very much limiting during reproductive stage thus abortion of flower to fruit is normal phenomenon. Application of USG may be an alternative approach to release nitrogen slowly from vegetative growth to reproductive stage. Alam (2002) reported that USG has many fold advantages to improve crop growth and yield. Triggering nitrogen at the plant demand would be attempt towards yield improvements of pulse (Deolankar, 2005; Mukesh, 2006). So, nitrogen management is required synchronizing this demand of plant growth stages when plant may uptake nitrogen during its reproductive stage. Recovery of nitrogen from point placement of urea as super granule was 49% higher than prilled urea. The increase in plant nitrogen recovery consequently increases plant height, number of leaf, tillering and a yield or agronomic efficiency of plant (Humphreys *et al.*, 2006).

Lack of attention on fertilizer application in proper way with appropriate amount is identified for lowering yields (Mansoor, 2007). Deep point placement of USG increases N use efficiency in wetland crop and the placement of USG at 8-10 cm depth of soil can save 30% nitrogen than prilled urea, increases nutrient absorption and ultimately increases yields (Singh *et al.*, 1993; Singh and Singh, 1986). Nitrogen available before flowering might have some influencing technique that would be better utilization of nitrogen for reducing flower and pod droppings. Applied basal nitrogen is lost due to leaching, surface run off, NH₃ volatilization, denitrification and other processes consequently decreases nitrogen use efficiency. The scientist all over the world is fighting to increase fertilizer use efficiency through the use of slow release fertilizers through right timing and

placement of USG. Among these deep placements of large granule instead of broadcasting prilled urea is most promising (Hossain, 2008).

Under the above considerations, the present study was undertaken to maximize the seed yield of grasspea answering the questions of application of prilled urea or urea super granules (USG) with the following objectives:

1. To investigate the influence of prilled and USG on the plant characters, yield and yield attributes of grasspea.
2. To find out the best method of USG application for maximum seed yield of grasspea.

CHAPTER II

REVIEW OF LITERATURE

Grasspea is an important pulse crop in Bangladesh and in many other countries of the world. The crop has been given less attention by the researchers in respect of different agronomic practices especially placement of nitrogen through prilled urea and urea super granules (USG) and normally it grows without any management including fertilization. Based on this a very few research work related to growth, yield and development of grasspea with different management practices have been carried out in our country. Research works related to the response of prilled urea and urea super granules are very limited in Bangladesh context. However, research findings related to the effect of prilled urea and USG on other crops so far have been done at home and abroad have been reviewed in this chapter under the following headings-

2.1 Effects of prilled urea on different pulse crop

2.1.1 Plant height

An experiment was conducted by Sardana and Verma (1987) in Delhi, India, they reported that application of nitrogen, phosphorus and potassium fertilizers in combination resulted in significant increase in plant height of mungbean.

Patra and Padhi (1989) noticed increased plant height of chickpea over control with 20 kg N along with 40 kg P ha⁻¹. Arvadia and Patel (1988) observed stimulatory effect of nitrogen or phosphorus alone at the rate of 25 kg ha⁻¹ on the growth of chickpea plants. They also reported appreciable increase in the plant height than those in control plots. Application of phosphorus alone at the rate of 50 kg ha⁻¹ did not show any significant effect on plant height over 25 kg P ha⁻¹.

Rathore and Patel (1991) noticed that application of 18 kg N along with 46 kg P ha⁻¹ increased plant height of chickpea over no N application. Vadavia *et al.*

(1991) noticed that application of 20 kg ha⁻¹ N and 40 kg P ha⁻¹ increased plant height of chickpea significantly over no N and P application.

Dahiya *et al.* (1993) reported increased in plant height of chickpea using N and P at the rate of 18-27 and 46-69 kg ha⁻¹, respectively.

Chaudhari *et al.* (1998) found a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on increased in chickpea plant height.

Oad and Buriro (2005) conducted a field experiment to determine the effect of different NPK levels (0-0-0, 10-20-20, 10-30-30, 10-30-40 and 10-40-40 kg ha⁻¹) on the growth and yield of mungbean cv. AEM 96 in Tandojam, Pakistan. The different NPK levels significantly affected the crop parameters. The 10-30-30 kg NPK ha⁻¹ was the best treatment, recording plant height of 56.25 cm.

A field experiment was conducted by Raman and Venkataramana (2006) in Annamalainagar, Tamil Nadu, India to investigate the effect of foliar nutrition on crop nutrient uptake and yield of greengram (*V. radiata*). Crop nutrient uptake, yield and its attributes (number of pods per plant and number of seeds per pod) of greengram augmented significantly due to foliar nutrition. The foliar application of 2% diammonium phosphate + NAA + Penshibao was significantly superior to other treatments in increasing the values of plant height.

Aliloo *et al.* (2012) studied the effects of foliar spraying of aqueous solutions 2 and 4% urea at two stages (before and after flowering) and 20 kg ha⁻¹ urea application in soil. Results showed that the effect of urea treatment on plant height was notable the highest plant height was obtained by application of 20 kg ha⁻¹ urea in soil.

Rhizobium and inorganic nitrogen fertilization on some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, investigated by Namvar *et al.* (2013) with mineral nitrogen fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with *Rhizobium*

bacteria as sub plots and found that plant height were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

An experiment was conducted by Abbasi *et al.* (2013) with nitrogen rates at four levels (0, 25, 50 and 75 kg urea ha⁻¹). The result reveal that plant height were significantly affected by urea nitrogen rates and seed inoculation.

Farzana (2013) conducted an experiment with different forms of urea (prilled urea, PU and urea super granules, USG) for BARI masur-6 cultivation and the tallest plant was observed with USG placed at 20 cm distance (avoid one row) and lowest was recorded from USG placed at 40 cm distance (avoid three rows). USG placed at 10 cm depth at 20-30 cm distance showed similar performance. She also reported that USG performed better than prilled urea.

Rumana (2013) carried out an experiment with prilled urea and urea super granules in chickpea cultivation and reported that, the tallest plant (14.37 cm, 27.28 cm, 45.07 cm, 52.52 cm, 58.76 cm and 61.23 cm) was observed from USG placed at 20 cm distance (avoid two rows) and the shortest plant (10.35 cm, 23.26 cm, 35.32 cm, 41.52 cm, 47.40 cm and 49.27 cm) in USG placed at 40 cm distance (avoid three rows) at 20, 40, 60, 80 and 100 days after sowing (DAS).

2.1.2 Branches plant⁻¹

Rathore and Patel (1991) found that the doses of 18 kg N and 46 kg P ha⁻¹ were most effective in increasing the number of branches plant⁻¹ of chickpea. Vadavia *et al.* (1991) reported that application of 20 kg N ha⁻¹ and P ha⁻¹ increased number of branches plant⁻¹ of chickpea.

An experiment was conducted by Dahiya *et al.* (1993) and reported that application of 18-27 kg N and 46-69 kg P ha⁻¹ increased number of branches plant⁻¹ in chickpea.

Chaudhari *et al.* (1998) found a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on increased in relation to the chickpea number of primary and secondary branches plant⁻¹.

The effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) on mungbean cultivars MH 85-111 and T44 were determined in a field experiment conducted by Rajender *et al.* (2002) in Hisar, Haryana, India during the summer. The number of branches increased with increasing N rates.

Malik *et al.* (2003) conducted a study to determine the effect of varying levels of nitrogen (0, 25, and 50 kg ha⁻¹) and phosphorus (0, 50, 75 and 100 kg ha⁻¹) on the yield and quality of mungbean cv. NM-98. They observed that number of branches per plant was found to be significantly higher at 25 kg N ha⁻¹.

Namvar *et al.* (2013) conducted an experiment on rhizobium and inorganic nitrogen fertilization on some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, with mineral nitrogen fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with *Rhizobium* bacteria in sub plots and found that number of primary and secondary branches were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

2.1.3 Dry matter content

Chaudhari *et al.* (1998) found a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on dry matter content plant⁻¹ in chickpea.

Yakadri *et al.* (2002) studied the effect of nitrogen (40 and 60 kg ha⁻¹) on crop growth and yield of green gram (cv. ML-267). Application of nitrogen at 20 kg ha⁻¹ resulted in the significant increase in dry matter content in above ground part.

Mukesh (2006) conducted a field experiment to study the impact of starter doses of nitrogen (0, 15 and 30 kg ha⁻¹) on nodulation and yield of different cultivars

(Radhey, Avarodhi and K-850) of chickpea under irrigated condition and reported that dry matter content plant^{-1} was highest in the crop treated with 30 kg N ha^{-1} .

2.1.4 Pods plant^{-1}

In trials, on clay soils during the summer season Patel *et al.* (1984) observed the effect of N levels (0, 10, 20 and 30 kg N ha^{-1}) and that of the P (0, 10, 20, 40, 60 and $80 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$) on the growth and seed yield of mungbean. In that experiment, it was found that application of 30 kg N ha^{-1} along with $40 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ significantly increased the number of pods plant^{-1} .

Patel and Parmer (1986) conducted an experiment on the response of greengram to varying levels of nitrogen and phosphorus. They observed that increasing N application to rainfed mungbean (cv. Gujrat-1) from 0 to 50 kg N ha^{-1} increased the number of pods plant^{-1} .

Patra *et al.* (1989) noticed that number of pods plant^{-1} of chickpea increased over control with 20 kg N along with 40 kg P ha^{-1} .

Vijai *et al.* (1990) carried out an experiment with gardenpea cv. Bonneville on N or P. They found that increasing rates of N or P up to 40 kg ha^{-1} significantly increased pod yield.

Rathore and Patel (1991) observed that maximum number of pods plant^{-1} when chickpea was provided with 18 kg N along with 46 kg P ha^{-1} .

Bhopal and Singh (1990) conducted an experiment with the semi dwarf garden pea cv. Lincoln, which received N at the rate of 0, 20, 40 and 60 kg ha^{-1} , P_2O_5 at 0, 30, 60 and 90 kg ha^{-1} and K_2O at 30 kg ha^{-1} . They reported that increasing N rates up to 40 kg ha^{-1} increased green pod yield. Further addition of nitrogen (60 kg ha^{-1}) tended to decrease the yield.

Vadavia *et al.* (1991) found that number of pods plant^{-1} of chickpea increased following application of 20 kg N ha^{-1} and 40 kg P ha^{-1} .

Tank *et al.* (1992) observed when mungbean was fertilized with 20 kg N along with level of 40 kg P₂O₅ ha⁻¹ increased seed yield significantly over the unfertilized control. They also reported that mungbean fertilized with 20 kg N ha⁻¹ along with 75 kg P₂O₅ ha⁻¹ significantly increased the number of pods plant⁻¹.

Khan *et al.* (1992) reported that the application of 20 kg N + 50 kg P₂O₅ ha⁻¹ in chickpea produced significantly higher number of pods plant⁻¹. Negi (1992) carried out an experiment with 4 levels of N (10, 20, 40, 60 kg ha⁻¹) and 3 of P₂O₅ on vegetable pea. He reported that the application of 20 kg N ha⁻¹ gave the highest green pod yield (1.72 t ha⁻¹).

A field study was conducted by Chaudhari *et al.* (1998) and reported a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on increased in chickpea pods plant⁻¹ and protein content in seed over control.

Karadavut and Ozdemir (2001) conducted a field trial on *Rhizobium sp.* and nitrogen on chickpea cultivars. They found that *Rhizobium* inoculation and 30 kg N ha⁻¹ significantly increased pods plant⁻¹.

Srinivas *et al.* (2002) conducted an experiment on the performance of mungbean at 0, 25 and 60 kg N ha⁻¹) and 0, 25, 50 and 60 kg P ha⁻¹) were tested. They observed that the number of pods plant⁻¹ was increased with the increasing rates of N up to 40 kg ha⁻¹ followed by a decrease with further increase in N.

The effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) on mung bean cultivars MH 85-111 and T44 were determined in a field experiment conducted by Rajender *et al.* (2002) in Hisar, Haryana, India during the summer and reported that the number of pods plant⁻¹ increased with increasing N rates.

A field experiment to study the impact of starter doses of nitrogen (0, 15 and 30 kg ha⁻¹) on nodulation and yield of different cultivars (Radhey, Avarodhi and K-850) of chickpea under irrigated condition was conducted by Mukesh (2006) and

reported that number of pods plant⁻¹ were highest in the crop treated with 30 kg N ha⁻¹.

A study was conducted by Nigamananda and Elamathi (2007) in Uttar Pradesh, India to evaluate the effect of N application time as basal urea spray and use of plant growth regulator (NAA at 40 ppm) on the yield and yield components of greengram cv. K-851. The recommended rate of N:P:K (20:50:20 kg ha⁻¹) as basal was used as a control. Treatments were as included: 1/2 basal N + foliar N as urea at 25 or 35 days after sowing (DAS); 1/2 basal N + 1/4 at 25 DAS + 1/4 at 35 DAS as urea; and 1/2 basal N + 1/2 foliar spraying as urea + 40 ppm NAA. 2% foliar spray of urea and NAA, applied at 35 DAS, resulted in the highest values for number of pods plant⁻¹ (38.3).

A research work was conducted by Namvar *et al.* (2013) with the application of rhizobium and inorganic nitrogen fertilization on some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, with mineral nitrogen fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with *Rhizobium* bacteria as sub plots and found that number of pods plant⁻¹ were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

An experiment was conducted by Abbasi *et al.* (2013) with nitrogen rates at four levels (0, 25, 50 and 75 kg urea ha⁻¹) and the result showed that number of pod plant⁻¹, were significantly affected by nitrogen rates and number of pod plant⁻¹ (32.48) was obtained in the higher nitrogen rates.

Farzana (2013) conducted an experiment with different forms of urea (prilled urea, PU and urea super granules, USG) for BARI masur-6 cultivation and found that pods plant⁻¹ (47.96) was statistically maximum with USG placed at 20 cm distance (avoid one row) and minimum values were recorded from USG placed at 40 cm distance (avoid three rows).

Rumana (2013) carried out an experiment with prilled urea and urea super granules in chickpea cultivation and found the highest pods plant⁻¹ (68.60) from USG placed at 20 cm distance (avoid two rows), while the lowest number of pods plant⁻¹ (53.27) from USG placed at 40 cm distance (avoid three rows).

2.1.5 Seeds pod⁻¹

Patel and Parmer (1986) conducted an experiment on the response of greengram to varying levels of nitrogen and phosphorus. They observed that increasing N application to rainfed mungbean (cv. Gujrat-1) from 0 to 50 kg N ha⁻¹ increased the number of seeds pod⁻¹.

Patra and Padhi (1989) noticed in chickpea increased number of seeds pod⁻¹ over control with 20 kg N along with 40 kg P ha⁻¹.

Rathore and Patel (1991) performed an experiment on chickpea with different levels of nitrogen and phosphorus fertilizers. They reported that application of 18 kg N along with 46 kg P ha⁻¹ resulted with significant increase in the chickpea seeds pod⁻¹.

Malik *et al.* (2003) investigated the effect of varying levels of nitrogen (0, 25 and 50 kg ha⁻¹) and P (0, 50, 75 and 100 kg ha⁻¹) on the yield and quality of mungbean cv. NM-98. They found that number of seeds pod⁻¹ was significantly affected by varying levels of nitrogen and phosphorus.

An experiment was carried out by Mukesh (2006) to study the impact of starter doses of nitrogen (0, 15 and 30 kg ha⁻¹) on nodulation and yield of different cultivars (Radhey, Avarodhi and K-850) of chickpea under irrigated condition and reported that number of seeds plant⁻¹ was highest in the crop treated with 30 kg N ha⁻¹.

Nigamananda and Elamathi (2007) conducted an experiment in Uttar Pradesh, India to evaluate the effect of N application time as basal urea spray and use of plant growth regulator (NAA at 40 ppm) on the yield and yield components of

greengram cv. K-851. The recommended rate of N:P:K (20:50:20 kg ha⁻¹) as basal was used as a control. Treatments were as included: 1/2 basal N + foliar N as urea at 25 or 35 days after sowing (DAS); 1/2 basal N + 1/4 at 25 DAS + 1/4 at 35 DAS as urea; and 1/2 basal N + 1/2 foliar spraying as urea + 40 ppm NAA. 2% foliar spray of urea and NAA, applied at 35 DAS, resulted in the highest seeds pod⁻¹ (7.67).

A field experiment was successfully finished by Namvar *et al.* (2013) to find out the effect of rhizobium and inorganic nitrogen fertilization on some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, with mineral nitrogen fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with *Rhizobium* bacteria as sub plots and found that number of grains plant⁻¹ were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

Rumana (2013) carried out an experiment with prilled urea and urea super granules in chickpea cultivation and found the highest seeds pod⁻¹ (2.87) in USG placed at 20 cm distance (avoid two rows), while the lowest number of seeds pod⁻¹ (1.77) was observed from USG placed at 40 cm distance (avoid three rows).

2.1.6 Weight of 1000 seeds

In trials, on clay soils during the summer season Patel *et al.* (1984) observed the effect of N levels (0, 10, 20 and 30 kg N ha⁻¹) and that of the P (0, 10, 20, 40, 60 and 80 kg P₂O₅ ha⁻¹) on the growth and seed yield of mungbean. They observed that application of 40 kg P₂O₅ ha⁻¹ along with 20 kg N ha⁻¹ significantly increased the 1000 seeds weight of mungbean.

Results of an experiment, conducted by Sardana and Verma (1987) in Delhi, India, revealed that application of nitrogen, phosphorus and potassium fertilizers in combination resulted in significant increases in 1000 seeds weight of mungbean.

Patra and Padhi (1989) reported that when 20 kg N along with 40 kg P ha⁻¹ were applied, it increased 1000 seeds weight of chickpea over control.

Bali *et al.* (1991) conducted a field trial on mungbean in kharif seasons on silty clay loam soil. They revealed that 1000 seeds weight increased with 40 kg N ha⁻¹ and 60 kg P₂O₅ ha⁻¹.

Rathore and Patel (1991) reported that application of 18 kg N ha⁻¹ along with 40 kg P ha⁻¹ increased 1000 seeds weight. Vadavia *et al.* (1991) found that seed weight increased following application of 20 kg N ha⁻¹ and 40 kg P ha⁻¹ of chickpea.

Srinivas *et al.* (2002) conducted an experiment on the performance of mungbean at 0, 25 and 60 kg N ha⁻¹ and 0, 25, 50 and 60 kg P ha⁻¹ were tested. They observed 1000 seeds weight increased with increasing rates of N up to 40 kg ha⁻¹.

Mukesh (2006) conducted a field experiment to study the impact of starter doses of nitrogen (0, 15 and 30 kg ha⁻¹) on nodulation and yield of different cultivars (Radhey, Avarodhi and K-850) of chickpea under irrigated condition and reported that 100 seeds weight were highest in the crop treated with 30 kg N ha⁻¹.

An experiment was conducted by Abbasi *et al.* (2013) with nitrogen rates at four levels (0, 25, 50 and 75 kg urea ha⁻¹) the result reveal that 100 grains weight were significantly affected by nitrogen rates and seed inoculation.

Farzana (2013) conducted an experiment in SAU with different forms of urea (prilled urea, PU and urea super granules, USG) for BARI Masur-6 cultivation found that 1000-seed weight (26.33g) was statistically maximum with USG placed at 20 cm distance (avoid one row) and minimum values were recorded from USG placed at 40 cm distance (avoid three rows). USG placed at 10 cm depth at 20-30 cm distance showed similar performance. She also reported that USG performed better than prilled urea.

Rumana (2013) carried out an experiment in SAU with prilled urea and urea super granules in chickpea cultivation and found the highest weight of 1000 seeds from USG placed at 20 cm distance (avoid two rows), while the lowest weight of 1000 seeds (188.79 g) from USG placed at 40 cm distance (avoid three rows).

2.1.7 Seed yield

Tomar and Sharma (1985) obtained highest seed yield in chickpea of two consecutive years with the application of N, P and K at the rate of 20, 40 and 20 kg ha⁻¹ respectively over control. An experiment was conducted by Trung and Yoshida (1983) using 0-100 ppm N as treatment in the form of ammonium nitrate or 10 or 100 ppm N as urea, sodium nitrate, ammonium nitrate or ammonium sulphate. They found that seed yield of mungbean increased with the increase in N up to 50 ppm.

Shamim and Naimat (1987) reported that application of 10 kg N + 75 kg P₂O₅ ha⁻¹ to *Cicer arietinum* cv. C-727 increases seed yields cover uninoculated seed from 583 to 878 kg ha⁻¹. Application of 20 kg N ha⁻¹ increased grain yield of chickpea reported by Subba Rao *et al.* (1986).

Khokar and Warsi (1987) reported maximum seed yield in chickpea with application of 18 kg N ha⁻¹. On the other hand, Patel *et al.* (1989) observed no significant yield variation in chickpea with the application of 15-30 kg N ha⁻¹.

Pongkao and Inthong (1988) applied N at the rate of 0-60 kg ha⁻¹ on mungbean and reported that application of 15 kg N ha⁻¹ was found to be superior giving 23% higher seed yield over the control. Addition of 60 kg N ha⁻¹ tended to produced seed yield which was at par of 15 kg N ha⁻¹.

Arvadia and Patel (1988) observed stimulatory effect of nitrogen at the rate of 25 kg ha⁻¹ on chickpea plants and reported appreciable increased in seed yield than those in control plots. They also found application of nitrogen alone at the rate of 50 kg ha⁻¹ showed no additional of that parameter over 25 kg N ha⁻¹.

Arya and Kalra (1988) reported that application of N at the rate of 50 kg ha⁻¹ along with 50 kg P ha⁻¹ increased mungbean yield. Another experimental result from field experiments conducted by Mahadkar and Saraf (1988) of mungbean revealed that the application of N with P and K at 20:25 kg ha⁻¹ gave higher seed yield.

Dahiya *et al.* (1989) noted an increase in seed yield in chickpea over control with the application of N, P and K at the rate of 20, 40 and 20 kg ha⁻¹ respectively. Patra *et al.* (1989) conducted an experiment on chickpea with different N and P rates. They stated that application of 20 kg N and 40 kg P ha⁻¹ increased grain yield of chickpea. Application of 25 kg N + 50 kg P ha⁻¹ gave the highest yield in the experiment of Javiya *et al.* (1989).

A field experiments was conducted by Sarkar and Banik (1991) to study the effect of N and P on yield of mungbean. Results showed that application of N along with P significantly increased the seed yield of mungbean. The maximum seed yield was obtained with the combination of 20 kg N and 60 kg P₂O₅ ha⁻¹.

Vadavia *et al.* (1991) found significant higher seed yield of chickpea following application of 20 kg ha⁻¹ N and 40 kg P ha⁻¹. Rathore and Patel (1991) noticed that application of 18 kg N along with 46 kg P ha⁻¹ increased seed yield of chickpea by 28.7% over no N application.

Khan *et al.* (1992) also reported that application of N and P increased grain yield of chickpea significantly over no N and P application. The application of 20 kg N + 50 kg P₂O₅ ha⁻¹ resulted with significant increase in the chickpea yield. Dahiya *et al.* (1993) noticed higher seed yield in chickpea over control while using N and P at rate of 18-27 and 46-69 kg ha⁻¹ respectively.

Patel *et al.* (1992) conducted a field trial to evaluate the response of mungbean to sulphur fertilization under different levels of nitrogen and phosphorus. Greengram cv. Gujrarat 2 and K 851 were given 10 kg N + 20 kg P ha⁻¹, 20kg N + 40 kg P ha⁻¹

and 0, 10, 20 or 30 kg S ha⁻¹ as gypsum. Seed yield was 1.2 and 1.24 t ha⁻¹ in Gujarat 2 K 851 respectively 20 kg N + 40 kg P ha⁻¹.

Chaudhari *et al.* (1998) conducted a field trial with chickpea grain with different rates of N and P fertilizer. They found a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on the growth and yield of chickpea.

Reddy and Ahlawat (1998) noticed that application 18 kg N, 46 kg P and 5.25 kg Zn ha⁻¹ increased grain and straw yield of chickpea. They also found increase in nitrogen, phosphorus and zinc uptake by plants leading to increase in protein yield.

Karle and Pawar (1998) examined the effect of varying levels of N and P fertilizers on summer mungbean. They reported higher seed yield in mungbean with the application of 15 kg N ha⁻¹ and 40 kg P₂O₅ ha⁻¹.

A field experiment was carried out by Sharma and Sharma (1999) during summer seasons at Golaghat, Assam, India. Mungbean was grown using farmers practices (no fertilizer) or using a combinations of fertilizer application (30 kg N + 35 kg P₂O₅ ha⁻¹). Seed yield was 0.40 t ha⁻¹ with farmer's practices, while the highest yield was obtained by the fertilizer application (0.77 t ha⁻¹).

Mahboob and Asghar (2002) studied the effect of seed inoculation at different nitrogen levels on mungbean at the agronomic research station, Farooqabad in Pakistan. They reported that seed inoculation +50-50-0 NPK kg ha⁻¹ exhibited superior performance in respect of seed yield (955 kg ha⁻¹).

The effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) on mungbean cultivars MH 85-111 and T44 were determined in a field experiment conducted by Rajender *et al.* (2002) in Hisar, Haryana, India during the summer and reported that grain yield increased with increasing rates of up to 40 kg N ha⁻¹ only.

Rajender *et al.* (2003) investigated the effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) fertilizer rates on mungbean genotypes MH 85-111 and T₄₄. Grain yield increased with increasing N rates up to 20 kg ha⁻¹. Further increase in N did not affect yield.

Malik *et al.* (2003) conducted an experiment to determine the effect of varying levels of nitrogen (0, 25, and 50 kg ha⁻¹) and phosphorus (0, 50, 75 and 100 kg ha⁻¹) on the yield and quality of mungbean cv. NM-98. They observed that a fertilizer combination of 25 kg N + 75 P kg ha⁻¹ resulted with maximum seed yield (1112.96 kg ha⁻¹).

Nadeem *et al.* (2004) studied the response of mungbean cv. NM-98 to seed inoculation and different levels of fertilizer (0-0, 15-30, 30-60 and 45-90 kg N-P₂O₅ ha⁻¹) under field conditions. Application of fertilizer significantly increased the yield and the maximum seed yield was obtained when 30 kg N ha⁻¹ was applied.

Oad and Buriro (2005) conducted a field experiment to determine the effect of different NPK levels (0-0-0, 10-20-20, 10-30-30, 10-30-40 and 10-40-40 kg ha⁻¹) on the growth and yield of mungbean in Tandojam, Pakistan. The different NPK levels significantly affected the crop parameters. The 10-30-30 kg NPK ha⁻¹ was the best treatment, recording the highest seed yield of 1205.2 kg ha⁻¹.

A field study conducted by Sharma and Sharma (2006) for two years at the Indian Agricultural Research Institute, New Delhi on a sandy clay loam soil showed that the application of NP increased the total grain production of a rice-wheat-mungbean cropping system by 0.5-0.6 t ha⁻¹, NK by 0.3-0.5 t ha⁻¹ and NPK by 0.8-0.9 t ha⁻¹ compared to N alone, indicating that the balanced use of primary nutrients was more advantageous than their imbalanced application.

A field experiment was conducted by Raman and Venkataramana (2006) in Annamalainagar, Tamil Nadu, India to investigate the effect of foliar nutrition on crop nutrient uptake and yield of greengram (*V. radiata*). There were 10 foliar

spray treatments, consisting of water spray, 2% diammonium phosphate at 30 and 45 days after sowing, 0.01% Penshibao, 0.125% Zn chelate, 30 ppm NAA. Crop nutrient uptake, yield and its attributes (number of pods per plant and number of seeds per pod) of greengram augmented significantly due to foliar nutrition. The foliar application of urea + NAA + Penshibao was significantly superior to other treatments in increasing the values of yield. The highest grain yield of 1529 kg ha⁻¹ was recorded with this treatment.

Tickoo *et al.* (2006) carried out an experiment on mungbean cultivars Pusa 105 and Pusa Vishal were sown at 22.5 and 30 m spacing and supplied with 36-46 and 58-46 kg NP ha⁻¹ in a field experiment conducted in Delhi, India during the kharif season of 2000. Cultivar Pusa Vishal recorded higher grain yield (1.63 t ha⁻¹) compared to cv. Pusa 105.

To evaluate the effect of N application time as basal urea spray and use of plant growth regulator (NAA at 40 ppm) on the yield and yield components of greengram cv. K-851 a study was conducted by Nigamananda and Elamathi (2007) in Uttar Pradesh, India. The recommended rate of N:P:K (20:50:20 kg ha⁻¹) as basal was used as a control. Treatments were as included: 1/2 basal N + foliar N as urea at 25 or 35 days after sowing (DAS); 1/2 basal N + 1/4 at 25 DAS + 1/4 at 35 DAS as urea; and 1/2 basal N + 1/2 foliar spraying as urea + 40 ppm NAA. 2% foliar spray of urea and NAA, applied at 35 DAS, resulted in the highest grain yield (9.66 q ha⁻¹).

An experiment was conducted by Abbasi *et al.* (2013) with nitrogen rates at four levels (0, 25, 50 and 75 kg urea ha⁻¹) and reported that grain yield were significantly affected by nitrogen rates and seed inoculation. Means comparison showed that maximum grain yield (1276.78 kg ha⁻¹) for 75 kg urea ha⁻¹.

Physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, was affected by rhizobium and inorganic nitrogen fertilization reported by Namvar *et al.* (2013) when an investigation was conducted with mineral nitrogen

fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with Rhizobium bacteria as sub plots and found that grain were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

Farzana (2013) conducted an experiment with different forms of urea (prilled urea, PU and urea super granules, USG) for BARI Masur-6 cultivation and observed that seed yield (2.11 t ha⁻¹) were statistically maximum with USG placed at 20 cm distance (avoid one row) and minimum values were recorded from USG placed at 40 cm distance (avoid three rows).

Rumana (2013) carried out an experiment with prilled urea and urea super granules in chickpea cultivation and found the highest seed yield (1.98 t ha⁻¹) from USG placed at 20 cm distance (avoid two rows), while the lowest seed yield (1.27 t ha⁻¹) was observed from USG placed at 40 cm distance (avoid three rows).

2.1.8 Stover yield

Vadavia *et al.* (1991) found that application of 20 kg ha⁻¹ N and 40 P ha⁻¹ increased significant straw yield of chickpea. Subba Rao *et al.* (1986) also reported that the rate of kg N ha⁻¹ was most effective in increasing straw yield.

Srinivas *et al.* (2002) conducted an experiment on the performance of mungbean at 0, 25 and 60 kg N ha⁻¹) and 0, 25, 50 and 60 kg P ha⁻¹) were tested and the experimented results they stated that the stover yield increased with increasing N up to 40 kg ha⁻¹.

The effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) on mung bean cultivars MH 85-111 and T44 were determined in a field experiment conducted by Rajender *et al.* (2002) in Hisar, Haryana, India during the summer and reported that straw yield increased with increasing N rates.

Rajender *et al.* (2003) investigated the effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) fertilizer rates on mungbean genotypes MH 85-111.

Stover yield increased with increasing N rates up to 20 kg ha⁻¹. Further increase in N did not affect yield.

An experiment was conducted by Abbasi *et al.* (2013) with nitrogen rates at four levels (0, 25, 50 and 75 kg urea ha⁻¹) the result reveal that stover yield were significantly affected by nitrogen rates and seed inoculation.

Rumana (2013) carried out an experiment with prilled urea and urea super granules in chickpea cultivation and found the highest stover yield (2.87 t ha⁻¹) was recorded from USG placed at 20 cm distance (avoid two rows), while the lowest stover yield (1.90 t ha⁻¹) was observed from USG placed at 40 cm distance (avoid three rows).

2.1.9 Biological yield

Results of an experiment conducted by Sardana and Verma (1987) in Delhi, India and stated that the application of nitrogen, phosphorus and potassium fertilizers in combination resulted in the significant increase in biological yield of mungbean.

Khan *et al.* (1992) reported from his study that biological yield of chickpea increased significantly with 20 kg N + 50 kg P₂O₅ ha⁻¹.

Karadavut and Ozdemir (2001) stated the application of *Rhizobium sp.* And 30 kg N ha⁻¹ on 3 chickpea cultivars in the winter season significantly increased pods plant⁻¹.

Rajender *et al.* (2003) investigated the effects of N (0, 10, 20 and 30 kg ha⁻¹) and P (0, 20, 40 and 60 kg ha⁻¹) fertilizer rates on mungbean genotypes MH 85-111 and T₄₄. Biological yield increased with increasing N rates up to 20 kg ha⁻¹. Further increase in N did not affect biological yield.

Tickoo *et al.* (2006) carried out an experiment on mungbean cultivars Pusa 105 and Pusa Vishal were sown at 22.5 and 30 m spacing and supplied with 36-46 and 58-46 kg NP ha⁻¹ in a field experiment conducted in Delhi, India. Cultivar Pusa Vishal recorded higher biological (3.66 1.63 t ha⁻¹) compared to cv. Pusa 105.

Some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, investigated by Namvar *et al.* (2013) with mineral nitrogen fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with *Rhizobium* bacteria as sub plots and found that biological yield were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

2.1.10 Harvest index

Harvest index may be influenced by N fertilization. Chaudhari *et al.* (1998) found that application of 20-40 kg N ha⁻¹ significantly influenced harvest index of chickpea.

Islam (2003) found a significant increase in harvest index in bush bean due to application of N. The lowest HI was in control and the maximum was at 36.8 was 36.8 kg N ha⁻¹.

Rajender *et al.* (2003) find out the effects of N (0, 10, 20 and 30 kg ha⁻¹) and P fertilizer rates on mungbean genotypes MH 85-111 and T₄₄. Harvest index increased with increasing N rates up to 20 kg ha⁻¹.

Mukesh (2006) conducted a field experiment to study the impact of starter doses of nitrogen (0, 15 and 30 kg ha⁻¹) on nodulation and yield of different cultivars (Radhey, Avarodhi and K-850) of chickpea under irrigated condition and reported that harvest index were highest in the crop treated with 30 kg N ha⁻¹.

Rhizobium and inorganic nitrogen fertilization on some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) cv. ILC 482, investigated by Namvar *et al.* (2013) with mineral nitrogen fertilizer at four levels (0, 50, 75 and 100 kg urea ha⁻¹) in the main plots, and two levels of inoculation with *Rhizobium* bacteria as sub plots and found that harvest index were obtained from the highest level of nitrogen fertilizer (100 kg urea ha⁻¹).

2.2 Effect of urea super granules on different crops

Yamada *et al.* (1981) stated that in trials in several countries rice was given 0-176 kg ha⁻¹ at each of three growth stages at urea or USG applied broadcast, incorporated or deep placed. Recovery of N was generally higher with USG than with urea and comparatively less amount USG was needed than urea to give the same grain yield. The superior performance of USG in increasing rice yield. The product is 40-50% more efficient than conventional urea.

In trials in several countries rice was given 0-176 to kg N ha⁻¹ at each of three growth stages as urea or USG applied broadcast, incorporated or deep placed. Recovery of N was generally higher with USG than with urea and less than urea was needed to give the same grain yield (Yamada *et al.*, 1981). Umar *et al.* (1981) conducted field trials at Maros, Langrang and Sulawesc and found that grain yield of low land rice was not affected by N application date and did not differ when N was applied as mudballs, urea briquets or USG.

Rambabu *et al.* (1983) reported that of various forms and methods of application of N fertilizers to rice grown under flooded conditions, placement of N as USG (1 g sizes) in the root zone at transplanting was most effective in increasing DM production, rice yield, followed by SCU incorporated before transplanting. Yield and N recovery were lowest with urea applied as basal drilling.

Kumar and Singh (1983) carried out an experiment with rice cv. Hindham grown by applying 29-116 kg N ha⁻¹ under flooded condition and stated that 87 kg N ha⁻¹ in the form of USG gave the highest yield which was 14.4% more compared to split application of urea.

Singh and Singh (1984) conducted a rainfed trial with rice on a calcareous silt-loam soil and reported that 29-87 kg N ha⁻¹ as USG and SCU before transplanting gave rice yields of 3.7 and 3.69 t ha⁻¹, respectively compared with 3.56 t with urea in three split dressings, 3.45 t with urea at transplanting only.

Rajagopalan and Palaniasami (1985) carried out an experiment with rice cv. TK 43 giving 50 or 75 kg N ha⁻¹ as neem-cake coated urea, coal coated urea and USG in the kharif season. They found that tallest plant (83 cm), number of panicles hill⁻¹ (10.00), panicle length (21 cm) and number of filled grains panicle⁻¹ (85) were obtained with 75 kg N as USG ha⁻¹.

Patel and Chandrawanshi (1986) conducted an experiment with rice cv. Sunuidhi (R-23.84) growing without N, or with 40 kg N ha⁻¹ as urea broadcast and incorporated as a basal dose before sowing, USG applied in rows and seeds drilled in alternate rows, urea or USG and seed drilled in the same furrow. They reported that the treatments did not significantly affect the number of panicles m⁻² but yield was highest (2.4 t ha⁻¹) in 40 kg N ha⁻¹ treatments.

Singh and Singh (1986) worked with different levels of nitrogen as urea, super granules (USG), sulphur coated urea and prilled urea @ 27, 54 and 87 kg ha⁻¹. They reported that number of tillers m⁻² increased with increasing nitrogen fertilizer. The number of tillers m⁻² was significantly greater in urea super granules than prilled urea in all levels of nitrogen.

Sudhakara and Prasad (1986) conducted a field experiment in rainy season in 1982-83 at the Indian agricultural Research Institute, New Delhi to study the relative efficiency of prilled urea, urea super granule and neem coated urea. They observed that performance of rice is more or less linear to USG and neem coated USG which were better than prilled urea.

Raja *et al.* (1987) conducted an experiment with rice and six different forms of nitrogen and mentioned that nitrogen as USG gave the highest average yield of 5.44 t compared with 4.64-4.92 t for nitrogen in five other forms. The USG at 75 kg N ha⁻¹ gave the highest yield of 7.2 t ha⁻¹. Juang and Wang (1987) stated that nitrogen rates were insignificant for both the broadcasting and deep placement.

Sarder *et al.* (1988) conducted a field experiment at Mymensingh on wetland rice applying urea, USG and sulphur coated urea (SCU) at 23.7, 47.4 and 94.8 kg N

ha⁻¹. Urea was broadcast in three split applications, USG was point placed at transplanting and SCU was broadcast and incorporated at transplanting. SCU was more efficient than the conventional method of urea application and point placement of USG only at the highest N rate, giving higher grain yield and greater plant height, panicle length and total number of grains panicle⁻¹ than the other N sources. At lower N rate, the crop's response was similar. USG did not show any advantage over conventional urea application.

Lal *et al.* (1988) conducted a field experiment on a silty loam soil during the rainy season of 1981-83 to evaluate the performance of USG and SCU in transplanted rice. They reported that placement of N as USG and broadcast incorporation of SCU were superior to prilled urea (applied in three split surface dressings) at 29, 58 and 87 kg N ha⁻¹ but not at 116 kg N ha⁻¹. SCU gave the highest response followed by USG and both maintained superiority over PU upto 87 kg N ha⁻¹.

Chakravorti *et al.* (1989) reported that applying 37.5 75.0 and 112.5 kg N ha⁻¹ as USG to rice gave rice yield of 3.85, 5.22 & 5.48 t ha⁻¹, respectively compared with 3.10, 4.29 and 4.97 t respectively, with N as urea and 1.95 without N. Rao and Ghai (1989) reported that SCU did not influence grain yield, LCU decreased yields and USG increased yield significantly.

Chauhan and Mishra (1989) conducted field experiments at pantnagar with rice applying 40, 80 or 120 kg N ha⁻¹ as five different forms of urea. They reported that USG point placed one week after transplanting gave the highest mean DM yield and PU gave the lowest grain yield while deep placed USG.

Rama *et al.* (1989) mentioned that the number of panicles m⁻² increased significantly when nitrogen level increased from 40 to 120 kg N ha⁻¹ as different modified materials. Urea super granules (USG) produced significantly higher number of panicles m⁻² and grains panicle⁻¹ than split application of prilled urea. Patra and Padhi (1989) stated that USG recorded the lowest number of tiller hill⁻¹, panicles hill⁻¹ and shortest panicle. Jee and Mahapatra (1989) also observed that

number of panicles m^{-2} were significantly higher @ 90 kg N ha^{-1} as deep placed urea super granules (USG) than split application of urea.

Narayanan and Thangamuthu (1991) carried out field experiments on rice cv. TKM9 and IR 20 at combatore, Tamil Nadu in 1984-85, N was applied at 30, 60 or 90 kg ha^{-1} using USG placed at a depth of 10 cm in the main plot. They noted that maximum yields of grain and straw were obtained from 90 kg N ha^{-1} , while the lowest was under the control treatment.

Reddy *et al.* (1991) carried out a field experiment in 1984 to study the effects of different N sources on rice cv. Jaya and Mangala. They found that the highest grain yield of 5863 kg ha^{-1} was gained from cv. Jaya treated with 112 kg ha^{-1} of urea super granules (USG) placed in the root zone.

Singh and Mishra (1993) conducted a field experiment on a calcareous alluvial sandy loam soil to see the comparative effect of prilled urea and urea super granules on yield and quality of Sugarcane. Use of USG up to 75% of the recommended dose of N increased sugar and sugarcane yields significantly as comparison with prilled urea. Cane juice quality was not affected significantly with these materials. USG was found stimulating ammonification significantly in the soil than prilled urea and reduced nitrification significantly.

Bhale and Salunke (1993) conducted a field trial to study the response of upland irrigated rice to nitrogen applied through urea and USG. They found that grain yield increased with up to 120 kg urea and 100 kg USG . Singh *et al.* (1993) opined that grain yield and N uptakes increased with increased rate of N application and were the highest with deep placed USG. Bhardwaj and Singh (1993) observed that placement of 84 kg N as USG produced a grain yield of 6.8 t ha^{-1} which was similar to placing 112 kg USG and significantly greater than other nitrogen sources and rates.

Quayum and Prasad (1994) showed that application of N up to 112.5 kg ha^{-1} increased grain (4.37 t ha^{-1}) and straw (5.49 t ha^{-1}) yields with fertile grain

panicle⁻¹ being the highest at this N rate. N applied as USG gave the best yield and yield attributes. It is reported that the slow release fertilizers were effective for rainfed lowland rice.

Patel and Mishra (1994) carried out an experiment on rice applied with 0, 30, 60 or 90 kg N ha⁻¹ as Mussorie rock phosphate-coated urea, neem cake-coated urea, gypsum coated urea, USG or prilled urea. The coated materials were incorporated before transplanting, urea super granules were placed 5-10 cm deep a week after transplanting and urea was applied in 3 split doses. They showed that N rate had no significant effect on panicle length, percent sterility and harvest index.

Mishra *et al.* (1994) carried out a field experiment with rice cv. sita giving 0 or 80 kg N ha⁻¹ as urea, USG and neem, lac, rock phosphate or karanj coated urea and showed that the highest grain yield was (3.39 t ha⁻¹) obtained by urea in three split applications. Das and Singh (1994) reported that grain yield and N use efficiency by rice were greater for deep placed USG than for USG broadcast and incorporated or three split applications of PU.

Subbaiah *et al.* (1994) reported that the highest grain yield (6.12 t ha⁻¹) was obtained with USG, 4.76 with PU + SSP and the lowest 2.89 t ha⁻¹ with the control. Grain yield, N use efficiency and apparent N recovery are consistently higher particularly during the *boro* season when N as USG is deep placed. The efficiency is further improved if hole is properly closed immediately after deep point placement of USG.

Panday and Tiwari (1996) observed that the grain yield of rice was highest with N applied as a basal dose of USG or mussoorie rock phosphate urea, (MRPU) applied in two split applications.

Vijaya and Subbaiah (1997) carried out an experiment with rice cv. IET 1444 treated with fertilizer @ 90 kg N ha⁻¹ as prilled urea, large granular urea or urea super granules (USG) and 70 kg P₂O₅ ha⁻¹ as single superphosphate or large diammonium phosphate, both applied by broadcasting or placement methods.

They showed that plant height, number of tillers, root length, number and weight of panicles, dry matter and grain yield of rice increased with increasing urea super granule size and were greater with the deep placement of nitrogen.

Haque (1998) conducted a field experiment on potato at BAU farm and farmers field at Madhupur. Four treatment were used (1) Control (without any fertilizer), (2) 100 Kg N ha⁻¹ from prilled urea, (3) USG, 2; 73 kg N ha⁻¹ from USG (4) USG-3; 109 kg N ha⁻¹ from USG point placement . USG greatly increased yield of potato tubers both at BAU farm and farmers field at Madhupur. Maximum potato tubers yield at BAU farm was from USG-3 treatment which was 24.16 t ha⁻¹ compared to 17.52 t ha⁻¹ from prilled urea application. Maximum yield of potato tubers at farmer's field was obtained from USG-3 treatment which was 26.50 t ha⁻¹ compared to 20.78 t ha⁻¹ from same dose of prilled urea application.

Department of Agricultural Extension (DAE) conducted 432 demonstrations of the effect of USG on boro rice in 72 thanas of 31 districts. It was reported that USG plots, on an average, produced higher yields than the PU plots while applying 30 to 40% less urea in the form of USG (Islam and Black, 1998).

Alam *et al.* (2000) carried out a socio economic study in two rice production environment (Gazipur and Tangail) to assess the comparative advantages of using urea super granule (USG) over prilled urea (PU) in modern rice production and to examine the differences in producers technical efficiency between USG urea and non user in crop management. Results revealed that comparatively low amount (36%) of urea was needed in modern boro rice production using USG instead of PU though 15% more labour was needed while weeding cost was a bit lower in USG using plots. Results also indicated additional yield of 0.87 t ha⁻¹ by using USG and this yield gain additional benefit of TK. 11,506 ha⁻¹.

Saikia *et al.* (2002) reported the response of Indian mustard cv. Pusa giant grown in New Delhi, India during rabi season. They found that those varieties response positively to the use of USG than prilled urea and neem coated urea.

Two field experiments were conducted by Rao *et al.* (2004) in Joypur India to study the effect of prilled urea, neem coated urea and urea super granules on a perennial aromatic herb, geranium (*Pelargonium graveolens* L Her.) grown on a sandy loam soil. They reported that, application of nitrogen increased the biomass and essential oil yields. USG and neem coated urea significantly increased the yields over prilled urea. The concentration and quality of essential oil were not influenced either by levels or carrier of N.

Talukder *et al.* (2004) conducted an experiment with two forms of urea i.e. prilled urea and USG were applied. Two years results revealed that USG had significant positive effect on the yield of tomato as compared to prilled urea. The recommended dose of N (150 kg/ha) from USG gave highest marketable fruit yield of 79.13 t/ha in the 1st year and 73.60 t/ha in the 2nd year. They also observed that when 150 kg and 135 kg/ha of N applied from USG gave 11% and 2% higher yield of tomato than that of using 150 kg/ha N from prilled urea, respectively.

Humphreys *et al.* (2006) reported that recovery of nitrogen from point placement of urea of super granule was 49% higher than prilled urea. The increase in plant nitrogen recovery consequently increases plant height, number of leaf, tillering and a yield or agronomic efficiency of rice plant.

Khalil *et al.* (2006) conducted a field experiment to know the recovery of spring wheat with urea super granules. They observed that the translocation of N from vegetative part to grain portion during grain filling stage (67-116 DAA) was 34.9% with the USG and 28.7% with PU, resulting in (711 kg ha⁻¹) more grain yield with the former than the later.

Singh and Singh (2006) conducted two field experiments for two crop cycles each for two years on an entities over *Citronella Java*. They found that the oil yields were 9% higher in USG than that of PU and N recovery of USG and PU were 31 and 21% respectively.

Hussain *et al.* (2010) conducted a number of experiments for three consecutive years at Farming System Research and development (FSRD) site at Tangail to evaluate the efficiency of USG application over prilled urea on the yield of cabbage and recorded the highest head yield (78.1 t ha⁻¹) was obtained with recommended dose of N as USG.

Ahmed *et al.* (2010) conducted a field experiment in farmers field at the MLT site at Madhupur and Ghatail upazila with four treatment on the production of hybrid maize and reported that the recommended dose of N as prilled urea gave yield of 9.2 t ha⁻¹. They also reported that the highest grain yield (10.30 t ha⁻¹) was obtained from the plot treated with recommended dose of N as USG which was similar to that of plots treated with 10% less than recommended dose of N as USG (9.44 t ha⁻¹).

A field experiment was conducted by Akter (2010) at Sher-e-Bangla Agricultural University, Dhaka to assess the comparative advantages of using urea super granule (USG) over prilled urea and also the effect of different management of nitrogenous fertilizer on growth, yield and yield attributing characters of mustard and noted that USG reduces 40% use of urea which lowers 20% of total cost in mustard production. He also reported that plant height, number of branches plant⁻¹, number of leaves plant⁻¹, total dry matter, number of siliqua plant⁻¹, siliqua length, number of seeds siliqua⁻¹, 1000-seed weight and seed yield (t ha⁻¹) was found highest when USG was applied as basal dose and all the characters showed lowest value when USG was applied at 25 DAS.

From the above reviews it is clear that nitrogenous fertilizer is an important nutrient for the production of any crop. Nitrogenous fertilizers are applied in the field as basal or foliar spray. On the other hand, the use of USG is emerging as an important method of nitrogen application for better growth and development of different crops reducing nitrogen loss and therefore improved nitrogen use efficiency by crops.

CHAPTER III

MATERIALS AND METHODS

The experiment was conducted during the period from November 2012 to March, 2013. This chapter includes materials and methods i.e. location of experimental site, soil and climate condition of the experimental plot, materials used, design of the experiment, data collection procedure and procedure of data analysis that were used in conducting the experiment are presented below under the following headings:

3.1 Experimental site

The experiment was conducted at the Farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka, Bangladesh. The experimental site is situated between 23^o74'/N latitude and 90^o35'/E longitude (Anon., 1989).

3.2 Climatic condition

The climate of experimental site is subtropical, characterized by three distinct seasons, the monsoon from November to February and the pre-monsoon period or hot season from March to April and the monsoon period from May to October. The monthly average temperature, humidity and rainfall during the crop growing period were collected from Weather Yard, Bangladesh Meteorological Department, and presented in Appendix I. During the experimental period the maximum temperature (31.7^oC) was recorded in the month of March 2013 while the minimum temperature (12.8^oC) in the month of January 2013. The highest humidity (78%) was recorded in the month of November, 2012, whereas the highest rainfall (39 mm) was recorded in the month of February 2013.

3.3 Characteristics of soil

The soil of the experimental field belongs to the Tejgaon series under the Agroecological Zone, Madhupur Tract (AEZ- 28) and the General Soil Type is Deep Red Brown Terrace Soils. A composite sample was made by collecting soil from several spots of the field at a depth of 0-15 cm before the initiation of the

experiment. The collected soil was air-dried, grind and passed through 2 mm sieve and analyzed at Soil Resources Development Institute (SRDI), Khamarbari, Farmgate, Dhaka for some important physical and chemical properties. The results showed that the soil composed of 27% sand, 43% silt, 30% clay and organic matter 0.78%, which have been presented in Appendix II.

3.4 Planting material

The variety BARI Khesari 3 was used as the test crop. The seeds were collected from the Agronomy Division of Bangladesh Agricultural Research Institute, Joydebpur, Gazipur. It was the released varieties of chickpea developed by Pulses Research Centre, which was recommended by the national seed board in the year of 2011. The variety can be grown in *Rabi* season. Life cycle of this variety ranges from 120-125 days. Maximum seed yield is 1.8-2.0 t ha⁻¹.

3.5 Land preparation

The land was irrigated before ploughing for land preparation. After having 'zoe' condition the land was first opened with the tractor drawn disc plough. Ploughed soil was brought into desirable tilth by 4 ploughing and cross-ploughing, harrowing and laddering. The stubble and weeds were removed. The first ploughing and the final land preparation were done on 12 and 20 November 2012, respectively. Experimental land was divided into 42 unit plots following the design of experiment.

3.6 Fertilizer application

Urea, Triple super phosphate (TSP), Muriate of potash (MOP), Gypsum, Zinc sulphate and Boric acid were used as a source of nitrogen, phosphorous, potassium, gypsum, sulphur and boron, respectively. The fertilizers were applied at the rate of 40, 80, 40, 30, 2 and 1.5 kg ha⁻¹, respectively as N, P₂O₅, K₂O, S, Zn and Boron following the Fertilizer Recommendation Guide (BARC, 2005). All of the fertilizers except N were applied during final land preparation and urea and USG were applied as per treatment.

3.7 Treatments of the experiment

The experiment consisted of the following treatments:

T₁ = Prilled urea (PU) broadcasted (control)

T₂ = PU given between two rows

T₃ = USG placed at 10 cm distance (avoid one row)

T₄ = USG placed at 10 cm distance (avoid two rows)

T₅ = USG placed at 10 cm distance (avoid three rows)

T₆ = USG placed at 20 cm distance (avoid one row)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₂ = USG placed at 40 cm distance (avoid one row)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

USG placed at 10 cm depth in each case.

3.8 Experimental design and layout

The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. An area of 23.5 m × 19.0 m was divided into three equal blocks. Each block was divided into 14 plots where 14 treatments were allotted at randomly. There were 42 unit plots altogether in the experiment. The size of the each unit plot was 3.0 m × 2.0 m. The space between two blocks and two plots were 1.0 m and 0.5 m, respectively. The layout of the experiment is shown in Figure 1.

3.9 Sowing of seeds in the field

The seeds of graasspea (variety BARI Khesari 3) were sown on 25 November, 2012. Before sowing seeds were treated with Bavistin to control the seed borne disease. The seeds were sown in solid rows in the furrows having a depth of 2-3 cm. Row to row distance was 30 cm.

3.10 Intercultural operations

3.10.1 Thinning

Seeds started germination of four DAS. Thinning was done two times at 8 and 15 days after germination (DAG) to maintain optimum plant population in each plot.

3.10.2 Irrigation and weeding

Irrigation was done as per requirements. The crop field was weeded before the application of irrigation.

3.10.3 Protection against insect and pest

At early stage of growth few worms (*Agrotis ipsilon*) and virus vectors (jassid) infested the young plants and at later stage of growth pod borer (*Maruca testulalis*) attacked the plant. Dimacron 50EC was sprayed at the rate of 1 litre/ha to control the insects.

3.11 Crop sampling and data collection

Ten plants from each plot were randomly selected and marked with sample card for data recording. Plant height, number of leaves per plant, branches per plant were recorded from selected plants at an interval of 20 days started from 20 DAS and continued upto harvest.

3.12 Harvest and post harvest operations

Harvesting was done when 90% of the pods became brown in color and it was done at 30 March, 2013. The matured pods were collected from a pre demarked area of three linear meter at the center of each plot.

3.13 Data collection

3.13.1 Crop growth characters

3.13.1.1 Plant height

The height of plant was recorded in centimeter (cm) at 30, 50, 70 and 90 days after sowing (DAS) and harvest. Data were recorded as the average of 10 plants selected at random from the inner rows of each plot that were tagged earlier. The height was measured from the ground level to the tip of the plant with the help of a meter scale.

3.13.1.2 Branches plant⁻¹

Total number of branches plant⁻¹ was recorded at 30, 50, 70 and 90 DAS and harvest. Data were recorded by counting branches from each plant and as the average of 10 plants selected at random from the inner rows of each plot.

3.13.1.3 Dry matter plant⁻¹

Data from ten sample plants from each plot were collected and gently washed with tap water, thereafter soaked with paper towel. The sample was oven dried at 70⁰C for a period of 72 hours. Then oven-dried samples were transferred into a desiccator and allowed to cool down to room temperature, thereafter dry weight of plant was taken and expressed in gram. Dry matter plant⁻¹ was recorded at 30, 50, 70 and 90 DAS and harvest.

3.13.1.4 Crop and relative growth rate

Using the data on total dry matter from each specific treatment, Crop Growth Rate (CGR) and Relative Growth Rate (RGR) were determined with following formulae (Hunt, 1978):

Crop Growth Rate (CGR)

Crop growth rate was calculated using the following formula:

$$\text{CGR} = \frac{1}{\text{GA}} \times \frac{W_2 - W_1}{T_2 - T_1} \text{g m}^{-2} \text{day}^{-1}$$

Where,

GA = Ground area (m²)

W₁ = Total dry weight at previous sampling date (T₁)

W₂ = Total dry weight at current sampling date (T₂)

T₁ = Date of previous sampling

T₂ = Date of current sampling

Relative Growth Rate (RGR)

Relative growth rate was calculated using the following formula:

$$\text{RGR} = \frac{\text{Ln}W_2 - \text{Ln}W_1}{T_2 - T_1} \quad (\text{g g}^{-1}\text{day}^{-1})$$

Where,

W₁ = Total dry weight at previous sampling date (time T₁)

W₂ = Total dry weight at current sampling date (time T₂)

T₁ = Date of previous sampling

T₂ = Date of current sampling

Ln = Natural logarithm

3.13.2 Yield contributing characters and yield of grasspea

3.13.2.1 Days from sowing to flowering

Days to sowing to flowering were measured by counting the number of days required to initiation of flower.

3.13.2.2 Pods plant⁻¹

Number of total pods of selected plants from each plot was counted and the mean numbers were expressed as plant⁻¹ basis. Data were recorded as the average of 10 plants selected at random from the inner rows of each plot.

3.13.2.3 Seeds pod⁻¹

Seeds pod⁻¹ was recorded randomly from selected plants at the time of harvest. Data were recorded as the average of 10 pods selected at random from the inner rows of each plot.

3.13.2.4 Weight of 1000 seeds

One thousand cleaned, dried seeds were counted from each harvest sample and weighed by using a digital electric balance and weight was expressed in gram (g).

3.13.2.5 Seed yield ha⁻¹

The seeds collected from 3 m² (3 m×1 m) square meter area of each plot were sun dried properly. The weight of seeds was taken and converted the yield in t ha⁻¹.

3.13.2.6 Stover yield ha⁻¹

The stover collected from 3 m² (3 m×1 m) square meter area of each plot was sun dried properly. The weight of stover was taken and converted the yield in t ha⁻¹.

3.13.2.7 Biological yield ha⁻¹

Seed yield and stover yield together were regarded as biological yield. The biological yield was calculated with the following formula:

$$\text{Biological yield} = \text{Seed yield} + \text{Stover yield.}$$

3.13.2.8 Harvest index

Harvest index was calculated from the seed yield and stover yield of grasspea for each plot and expressed in percentage.

$$\text{HI (\%)} = \frac{\text{Economic yield (seed weight)}}{\text{Biological yield (Total dry weight)}} \times 100$$

3.14 Statistical analysis

The data obtained for different parameters were statistically analyzed to find out the significant difference to the response of grasspea to the application of prilled urea and urea super granules with use MSTAT C statistical package. The mean values of all the characters were calculated and analysis of variance was performed by the 'F' (variance ratio) test. The significance of the difference among the treatment means was estimated by the Duncan's Multiple Range Test (DMRT) at 5% level of probability (Gomez and Gomez, 1984).

CHAPTER IV

RESULTS AND DISCUSSION

The experiment was conducted to study the response of grasspea to the application of prilled urea and urea super granules. Data on different crop growth characters, yield contributing characters and yield of grasspea were recorded and found significant differences among the recorded characters. The analyses of variance (ANOVA) of the data on different parameters are presented in Appendix III-VII. The results have been discussed with the help of table and graphs and possible interpretations given under the following headings:

4.1 Crop growth characters of grasspea

4.1.1 Plant height

Plant height of grasspea showed statistically significant variation at 30, 50, 70, 90 DAS and at harvest due to the application of prilled urea and urea super granules (Appendix III). At 30 DAS, the tallest plant (34.64 cm) was observed from T₇ [USG placed at 20 cm distance (avoid two rows)] which was statistically similar (32.72 cm and 31.67 cm, respectively) with T₅ [USG placed at 10 cm distance (avoid three rows)], T₁ [Prilled urea (PU) broadcasted] and closely followed (30.85 cm and 29.81 cm, respectively) by T₆ [USG placed at 20 cm distance (avoid one row)] and T₉ [USG placed at 30 cm distance (avoid one row)]. On the contrary, the shortest plant (19.34 cm) was recorded with T₁₄ [USG placed at 40 cm distance (avoid three rows)] which was statistically similar (21.5 cm) with T₁₂ [USG placed at 10 cm depth at 40 cm distance (avoid one row) (Table 1). At 50 DAS, T₇ (40.30 cm) treatment gave the longest plant which was statistically similar with T₉ (39.44 cm), T₅ (38.56 cm), T₆ (37.98 cm) and T₁ (37.18 cm), while the shortest plant was recorded from T₁₄ (27.30 cm) followed by T₁₂ (31.38 cm) and T₁₀ (37.90 cm) treatments. At 70 DAS, the tallest plant was observed from T₇ (47.35 cm) which was statistically similar with T₅ (46.07 cm), T₆ (45.49 cm), T₉ (45.35 cm) and T₁ (44.96 cm) treatments, while the shortest plant was recorded

from T₁₄ (30.55 cm) treatment followed by T₁₂ (36.16 cm) treatment. At 90 DAS, the highest plant height was observed from T₇ (59.32 cm) which was statistically similar with T₅ (57.01 cm) and T₁ (56.14 cm), whereas the lowest plant height was recorded from T₁₄ (38.11 cm) followed by T₁₂ (43.66 cm). At harvest, the longest plant was found from T₇ (65.49 cm) and it was statistically identical with T₅ (65.14 cm), T₁ (64.36 cm), T₈ (63.99 cm), T₆ (63.75 cm), T₉ (62.38 cm) and T₁₃ (61.72 cm) treatments, whereas the shortest plant was recorded from T₁₄ (57.01 cm) which was similar with T₁₂ (57.84 cm). Among the different treatments T₇ was found very effective to had maximum taller plants as plants were able to uptake nitrogen efficiently from placed urea but not replacing prilled urea application. Chaudhari *et al.* (1998) found a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on increased in chickpea plant height. Aliloo *et al.* (2012) reported that that the effect of urea treatment on plant height was notable the highest plant height was obtained by application of 20 kg ha⁻¹ urea in soil. Humphreys *et al.* (2006) reported that recovery of nitrogen from point placement of urea of super granule was 49% higher than prilled urea. The increase in plant nitrogen recovery consequently increases plant height. Talukder *et al.* (2004), Rao *et al.* (2004), Saikia *et al.* (2002), Alam *et al.* (2000), Haque (1998) and many other researchers recorded similar results in different crops from their earlier experiment. Farzana (2013) reported that the tallest plant of lentil was observed with USG placed at 20 cm distance (avoid one row) and lowest was recorded from USG placed at 40 cm distance (avoid three rows). Rumana (2013) also reported the tallest plant (14.37, 27.28, 45.07, 52.52, 58.76 and 61.23 cm) was observed from USG placed at 20 cm distance (avoid two rows) and the shortest plant (10.35, 23.26, 35.32, 41.52, 47.40 and 49.27 cm) was recorded in USG placed at 40 cm distance (avoid three rows) in chickpea at 20, 40, 60, 80 and 100 days after sowing (DAS) and harvest in same cropping season and the findings was similar to the findings of this study.

Table 1. Effect of application of prilled urea and urea super granule on plant height of grasspea at different growth stages

Treatment	Plant height (cm) at				
	30 DAS	50 DAS	70 DAS	90 DAS	At harvest
T ₁	31.67 a-c	37.18 ab	44.96 a-c	56.14 a-c	64.36 a
T ₂	21.41 gh	33.82 de	38.15 e-g	44.59 f	59.05 cd
T ₃	24.82 fg	32.83 cd	36.95 fg	47.56 ef	60.63 a-d
T ₄	22.63 gh	35.31 bc	39.47 ef	47.19 ef	59.39 b-d
T ₅	32.72 ab	38.56 ab	46.07 a-c	57.01 ab	65.14 a
T ₆	30.85 b-d	37.98 ab	45.49 a-c	54.59 b-d	63.75 ab
T ₇	34.64 a	40.30 a	47.35 a	59.32 a	65.49 a
T ₈	27.40 d-f	37.77 a-c	43.74 b-d	52.54 b-d	63.99 ab
T ₉	29.81 b-e	39.44 ab	45.35 abc	53.45 b-d	62.38 a-c
T ₁₀	28.54 c-e	37.90 e	43.35 cd	52.66 b-d	59.43 b-d
T ₁₁	26.60 ef	33.71 de	37.75 fg	50.44 de	59.22 b-d
T ₁₂	21.59 gh	31.38 e	36.16 g	43.66 f	57.84 cd
T ₁₃	28.97 c-e	37.25 bc	40.98 de	51.84 cd	61.72 a-d
T ₁₄	19.34 h	27.30 f	30.55 h	38.11 g	57.01 d
CV(%)	7.42	5.96	4.86	4.81	4.13

Mean having same or without letter(s) do not differ at 5% level of probability

T₁ = Prilled urea (PU) broadcasted (control)

T₄ = USG placed at 10 cm distance (avoid two rows)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

T₂ = PU given between two rows

T₅ = USG placed at 10 cm distance (avoid three rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

T₃ = USG placed at 10 cm distance (avoid one row)

T₆ = USG placed at 20 cm distance (avoid one row)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₂ = USG placed at 40 cm distance (avoid one row)

USG placed at 10 cm depth of each case

4.1.2 Branches plant⁻¹

Statistically significant variation was recorded for the application of prilled urea and urea super granules in terms of branches plant⁻¹ of grasspea at 30, 50, 70, 90 100 DAS and at harvest (Appendix IV). At 30 DAS, the maximum number of branches plant⁻¹ was recorded from T₇ (4.47) which was statistically similar with T₅ (4.40) and T₁ (4.33), T₁ (4.20), T₉ (4.13), T₁₃ (4.07) and T₁₀ (4.00) treatment. The minimum number was obtained from T₁₄ (2.27) treatment followed by T₁₂ (2.87) treatment. At 50 DAS, the maximum number branches plant⁻¹ was found from T₇ (9.50), which was statistically similar with T₅ (9.46), T₁ (9.35), T₆ (9.21), T₉ (8.87) and T₁₁ (8.85) treatment. The minimum number of branches was recorded from T₁₄ (6.13) which was statistically similar with T₁₂ (6.78) and T₂ (6.80) treatments. At 70 DAS, the maximum number branches plant⁻¹ was observed from T₇ (17.00) which was statistically similar with T₅ (16.08), while the minimum number from T₁₄ (10.37) which was statistically similar with T₁₂ (11.10) and T₂ (11.44). At 90 DAS, the maximum number branches plant⁻¹ was observed from T₇ (22.55), which was similar with T₅ (21.84), T₁ (4.33), T₉ (20.61) and T₁₀ (20.49), while the minimum number of branches plant⁻¹ was recorded from T₁₄ (12.22) followed by T₁₂ (15.09). At harvest, the maximum number branches plant⁻¹ was found from T₇ (25.77) treatment which was statistically similar with T₅ (24.99), T₁ (24.65) and T₆ (24.17) treatments. The minimum number of branches was obtained from T₁₄ (14.56) followed by T₁₂ (17.76). Treatment T₇ was found influential for higher numbers of branches plants⁻¹ probably due to nitrogen uptake efficiently from placed urea as prilled and urea super granules. Chaudhari *et al.* (1998) found a positive effect of nitrogen @ 20 and 40 kg ha⁻¹ on increased number of primary and secondary branches plant⁻¹.

Table 2. Effect of application of prilled urea and urea super granule on branches plant⁻¹ of grasspea at different growth stages

Treatment	Number of branches plant ⁻¹ at				
	30 DAS	50 DAS	70 DAS	90 DAS	At harvest
T ₁	4.33 a	9.35 ab	15.61 bc	21.07 a-c	24.65 ab
T ₂	2.93 d	6.80 fg	11.44 hi	15.48 f	19.11 ef
T ₃	3.20 cd	7.56 ef	12.46 gh	16.26 f	18.31 f
T ₄	3.60 bc	7.54 ef	13.56 e-g	18.81 de	20.82 de
T ₅	4.40 a	9.46 a	16.08 ab	21.84 ab	24.99 ab
T ₆	4.20 a	9.21 a-c	15.38 b-d	20.95 a-d	24.17 ab
T ₇	4.47 a	9.50 a	17.00 a	22.55 a	25.77 a
T ₈	3.00 d	8.27 c-e	13.08 fg	17.10 ef	17.96 f
T ₉	4.13 a	8.87 a-d	14.98 b-e	20.61 a-d	23.56 b
T ₁₀	4.00 ab	8.02 de	14.38 c-f	20.49 a-d	23.13 bc
T ₁₁	3.60 bc	8.85 a-d	13.95 d-f	19.04 c-e	21.26 cd
T ₁₂	2.87 d	6.78 fg	11.10 hi	15.09 f	17.76 f
T ₁₃	4.07 a	8.46 b-e	13.97 d-f	20.31 b-d	23.44 b
T ₁₄	2.27 e	6.13 g	10.37 i	12.22 g	14.56 g
CV(%)	6.74	6.19	5.66	6.16	5.50

Mean having same or without letter(s) do not differ at 5% level of probability

T₁ = Prilled urea (PU) broadcasted (control)

T₄ = USG placed at 10 cm distance (avoid two rows)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

USG placed at 10 cm depth of each case

T₂ = PU given between two rows

T₅ = USG placed at 10 cm distance (avoid three rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

T₃ = USG placed at 10 cm distance (avoid one row)

T₆ = USG placed at 20 cm distance (avoid one row)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₂ = USG placed at 40 cm distance (avoid one row)

4.1.3 Dry matter plant⁻¹

Application of prilled urea and urea super granules varied significantly at 30, 50, 70 and 90 DAS for dry matter plant⁻¹ of grasspea (Appendix V). At 30 DAS, the highest dry matter plant⁻¹ was recorded from T₇ (8.64 g), which was statistically similar to T₅ (8.52 g), T₁ (8.48 g), T₆ (8.15 g) and T₁₀ (8.01 g) treatments. The lowest dry matter plant⁻¹ was observed from T₁₄ (4.51 g) and it was statistically similar to T₁₃ (5.25 g) treatment (Table 3). At 50 DAS, the highest dry matter plant⁻¹ was recorded from T₇ (11.65 g), which was statistically identical with T₁₃ (8.00 g), T₁ (11.46 g) and T₆ (10.90 g), while the lowest dry matter plant⁻¹ was observed from T₁₄ (6.27 g) which was statistically similar with T₁₃ (8.10 g). At 70 DAS, the highest dry matter plant⁻¹ was recorded from T₇ (20.45 g) and it was statistically identical with T₁ (20.34 g), T₅ (20.17 g), T₆ (18.22 g), T₁₀ (18.18 g) and T₁₁ (17.28 g) treatments. The lowest dry matter plant⁻¹ was observed from T₁₄ (13.63 g) which was statistically similar with T₁₃ (15.14 g). At 90 DAS, the highest dry matter plant⁻¹ was recorded from T₇ (40.55 g) which was statistically similar with T₁ (40.37 g), T₅ (36.78 g) and T₁₂ (339.10 g), while the lowest dry matter plant⁻¹ was observed from T₁₄ (24.58 g) which was statistically similar with T₁₃ (27.89 g). Chaudhari *et al.* (1998) found a positive effect of nitrogen at the rate of 20 and 40 kg ha⁻¹ on increased in grasspea dry matter plant⁻¹.

Table 3. Effect of application of prilled urea and urea super granule on dry matter plant⁻¹ of grasspea at different growth stages

Treatment	Dry matter plant ⁻¹ (g)			
	30 DAS	50 DAS	70 DAS	At harvest
T ₁	8.47 ab	11.46 ab	20.34 a	40.37 a
T ₂	6.76 a-e	8.12 c-f	16.28 a-c	27.78 bc
T ₃	6.28 c-f	8.39 d-g	13.41 c	25.95 c
T ₄	6.72 a-e	8.81 c-f	15.73 a-c	30.92 a-c
T ₅	8.52 a	11.37 ab	20.17 ab	36.78 ab
T ₆	8.15 a-c	10.90 a-c	18.22 a-c	33.01 a-c
T ₇	8.64 a	11.65 a	20.45 a	40.55 a
T ₈	5.49 ef	7.44 fg	13.67 c	24.305 c
T ₉	7.49 a-d	10.17 a-e	14.24 c	27.62 bc
T ₁₀	8.01 a-c	10.71 a-d	18.18 a-c	33.49 a-c
T ₁₁	6.78 a-e	9.24 a-f	17.28 a-c	30.14 bc
T ₁₂	6.54 b-e	9.15 b-f	16.72 a-c	33.91 a-c
T ₁₃	5.75 d-f	8.00 e-g	15.40 bc	27.89 bc
T ₁₄	4.51 f	6.27 g	13.63 c	24.58 c
CV(%)	14.49	13.56	5.72	7.10

Mean having same or without letter(s) do not differ at 5% level of probability

T₁ = Prilled urea (PU) broadcasted (control)

T₄ = USG placed at 10 cm distance (avoid two rows)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

USG placed at 10 cm depth of each case

T₂ = PU given between two rows

T₅ = USG placed at 10 cm distance (avoid three rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

T₃ = USG placed at 10 cm distance (avoid one row)

T₆ = USG placed at 20 cm distance (avoid one row)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₂ = USG placed at 40 cm distance (avoid one row)

4.1.4 Crop growth rate

Crop growth rate (CGR) of grasspea showed significant variation at 30-50, 50-70 and 70 DAS-at maturity for the application of prilled urea and urea super granules (Appendix VI). At 30-50 DAS, the highest CGR was attained from T₇ (6.54 g m⁻²day⁻¹), which was similar with T₁ (6.53 g m⁻²day⁻¹), while the lowest CGR was found from T₅ (3.65 g m⁻²day⁻¹) which was similar to T₁₂ (4.36 g m⁻²day⁻¹) (Table 4). At 50-70 DAS, the highest CGR was attained from T₁ (9.87 g m⁻² day⁻¹) which was similar with T₇ (9.79 g m⁻²day⁻¹), whereas the lowest CGR was found from T₉ (4.52 g m⁻²day⁻¹). At 70 DAS-at maturity, the highest CGR value (20.09 g m⁻²day⁻¹) was attained from T₇, which was similar with T₁ (20.03 g m⁻²day⁻¹) and T₁₃ (17.19 g m⁻²day⁻¹), while the lowest (10.62 g m⁻²day⁻¹) from T₈ treatment, which was similar with T₂ (11.50 g m⁻²day⁻¹) and T₃ (12.54 g m⁻²day⁻¹).

4.1.5 Relative growth rate

Statistically significant variation was recorded for relative growth rate (RGR) of grasspea at 30-50, 50-70 and 70 DAS-at maturity for the application of prilled urea and urea super granules (Appendix VI). At 30-50 DAS, the highest RGR value was found from T₇ (0.148 g g⁻¹ day⁻¹) and it was statistically similar with T₁ (0.147 g g⁻¹ day⁻¹), whereas the lowest RGR was found from T₁₄ (0.124 g g⁻¹ day⁻¹) (Table 5). At 50-70 DAS, the highest RGR was found from T₃ (0.195 g g⁻¹ day⁻¹), whereas the lowest RGR was recorded from T₉ (0.159 g g⁻¹ day⁻¹). At 70 DAS-at maturity, the highest RGR value was found from T₇ (0.231 g g⁻¹ day⁻¹), whereas the lowest RGR was found from T₈ (0.203 g g⁻¹ day⁻¹) treatment.

4.1.6 Days from emergence to flowering

Days from emergence to flowering of grasspea showed statistically significant variation for the application of prilled urea and urea super granules (Appendix VII). Treatment T₇ took minimum days (45.33 days) to flowering and it was statistically identical with T₅ (47.00 days), T₆ (48.00 days) and T₁ (49.67 days), while the maximum days from emergence to flowering were found from T₁₄ (57.67 days) which was similar with T₁₂ (57.33 days) (Figure 2). Results indicated that T₇ treatment matured by 26% days earlier than T₁₄ treatment.

Table 4. Effect of application of prilled urea and urea super granule on crop growth rate of grasspea

Treatment	Crop Growth Rate (g m ⁻² day ⁻¹)		
	30 DAS-50 DAS	50 DAS-70 DAS	70 DAS-At maturity
T ₁	6.53 a	9.87 a	20.03 a
T ₂	4.59 b-d	8.19 a-c	11.50 b
T ₃	3.97 cd	5.56 bc	12.54 b
T ₄	4.91 a-d	7.69 a-c	15.19 ab
T ₅	5.82 ab	9.78 a	16.61 ab
T ₆	5.57 a-c	8.13 a-c	14.79 ab
T ₇	6.54 a	9.79 a	20.09 a
T ₈	4.36 b-d	6.92 a-c	10.62 b
T ₉	5.25 a-d	4.52 c	13.38 b
T ₁₀	5.60 a-c	8.30 a-c	15.31 ab
T ₁₁	5.25 a-d	8.92 ab	12.86 b
T ₁₂	4.36 b-d	7.93 a-c	12.75 b
T ₁₃	5.62 a-c	8.42 a-c	17.19 ab
T ₁₄	3.65 d	8.18 a-c	10.95 b
CV(%)	7.08	9.98	7.24

Mean having same or without letter(s) do not differ at 5% level of probability

T₁ = Prilled urea (PU) broadcasted (control)

T₄ = USG placed at 10 cm distance (avoid two rows)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

USG placed at 10 cm depth of each case

T₂ = PU given between two rows

T₅ = USG placed at 10 cm distance (avoid three rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

T₃ = USG placed at 10 cm distance (avoid one row)

T₆ = USG placed at 20 cm distance (avoid one row)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₂ = USG placed at 40 cm distance (avoid one row)

Table 5. Effect of application of prilled urea and urea super granule on relative growth rate of grasspea

Treatment	Relative growth rate (g g ⁻¹ day ⁻¹)		
	30 DAS-50 DAS	50 DAS-70 DAS	70 DAS-At maturity
T ₁	0.147 a	0.194 a	0.230 a
T ₂	0.133 ab	0.186 ab	0.205 b
T ₃	0.132 ab	0.169 bc	0.209 b
T ₄	0.131 ab	0.182 ab	0.217 ab
T ₅	0.145 a	0.194 a	0.222 ab
T ₆	0.144 ab	0.186 ab	0.216 ab
T ₇	0.148 a	0.195 a	0.231 a
T ₈	0.129 ab	0.179 ab	0.203 b
T ₉	0.143 ab	0.159 c	0.212 ab
T ₁₀	0.143 ab	0.186 ab	0.218 ab
T ₁₁	0.139 ab	0.190 a	0.211 ab
T ₁₂	0.135 ab	0.182 ab	0.210 ab
T ₁₃	0.140 ab	0.186 ab	0.220 ab
T ₁₄	0.124 b	0.186 ab	0.204 b
CV(%)	4.74	6.71	4.29

Mean having same or without letter(s) do not differ at 5% level of probability

T₁ = Prilled urea (PU) broadcasted

T₄ = USG placed at 10 cm distance (avoid two rows)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

USG placed at 10 cm depth of each case

T₂ = PU given between two rows

T₅ = USG placed at 10 cm distance (avoid three rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

T₃ = USG placed at 10 cm distance (avoid one row)

T₆ = USG placed at 20 cm distance (avoid one row)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₂ = USG placed at 40 cm distance (avoid one row)




Figure 2. Effect of application of prilled urea and urea super granules on days from semergence to flowering of grasspea ($S_x = 1.345$)

4.2 Yield contributing characters and yield of grasspea

4.2.1 Pods plant⁻¹

Statistically significant variation was recorded in terms of number of pods plant⁻¹ of grasspea due to the application of prilled urea and urea super granules (Appendix VII). The highest number of pods plant⁻¹ was recorded from T₇ (54.48) which was similar with T₅ (54.05), T₁ (52.55) and T₆ (51.75) treatments. The lowest number was observed from T₁₄ (45.09) which was similar with T₁₂ (45.26) and T₈ (54.48) (Table 6). Results indicated that T₇ treatment produced 28.78% higher pods over T₁₄ treatment. Application of prilled urea and urea super granules supported plant with maximum dry matter which eventually produced maximum number of pods plant⁻¹. Abbasi *et al.* (2013) reported that number of pod plant⁻¹ were significantly affected by nitrogen rates and number of pod plant⁻¹ (32.48) was obtained in the higher nitrogen rates. Farzana (2013) reported that pods plant⁻¹ (47.96) was maximum with USG placed at 20 cm distance (avoid one row) and minimum values were recorded from USG placed at 40 cm distance (avoid three rows) in lentil. Rumana (2013) also found the highest pods plant⁻¹ (68.60) from USG placed at 20 cm distance (avoid two rows), while the lowest number of pods plant⁻¹ (53.27) from USG placed at 40 cm distance (avoid three rows) in chickpea.

4.2.2 Seeds pod⁻¹

Application of prilled urea and urea super granules showed statistically significant variation for number of seeds pods⁻¹ of grasspea (Appendix VII). The highest number of seeds pods⁻¹ was recorded from T₇ (4.87) which was statistically similar with T₅ (4.67), T₁ (4.61) and T₆ (4.57), while the lowest number from T₁₄ (3.43) which was similar with T₁₂ (3.63) (Table 6). Patra and Padhi (1989) noticed in grasspea increased number of seeds pod⁻¹ over control with 20 kg N ha⁻¹.

4.2.3 Weight of 1000 seeds

Weight of 1000 seeds showed statistically significant variation due to the application of prilled urea and urea super granules (Appendix VII). The maximum weight of 1000 seeds was found from T₇ (40.57 g) treatment which was

statistically similar with T₅ (40.20 g), T₁ (39.67 g), T₉ (38.24 g) and T₈ (38.21 g) treatments, The lowest weight of 1000 seeds was recorded from T₁₄ (34.79 g) which was similar with T₁₂ (35.76 g) treatment (Table 6). Results revealed that T₇ treatment produced 23% more weight of 1000 seeds than that of T₁₄ treatment. Akter (2010) reported that 1000 seeds weight was the highest when USG was applied as basal dose and lowest when USG was applied at 25 DAS in mustard. Vadavia *et al.* (1991) found that seed weight increased following application of 20 kg N ha⁻¹ and 40 kg P ha⁻¹ of chickpea.

4.2.4 Seed yield ha⁻¹

Statistically significant variation was recorded for seed yield hectare⁻¹ of grasspea due to the application of prilled urea and urea super granules (Appendix VII). The highest seed yield (1.94 t ha⁻¹) was recorded from T₇ treatment which was statistically similar with T₅ (1.87 t ha⁻¹), T₁ (1.85 t ha⁻¹), T₆ (1.76 t ha⁻¹) and T₃ (1.74 t ha⁻¹) treatments, while the lowest seed yield was observed from T₁₄ (1.19 t ha⁻¹) treatment (Table 6). The maximum seed yield was harvested due to maximum pods plant⁻¹, seed pods⁻¹, which had greater number of seeds and highest 1000 grains weight. Nadeem *et al.* (2004) reported that the application of fertilizer significantly increased the yield and the maximum seed yield was obtained when 30 kg N ha⁻¹ was applied. Patel *et al.* (1984) reported that application of nitrogenous fertilizers becomes helpful in increasing the yield. The placement of USG at 8-10 cm depth of soil can save 30% nitrogen than prilled urea, increases nutrient absorption, improves soil health and ultimately increases the crop yields (Singh *et al.*, 1993; Singh and Singh, 1986). Farzana (2013) observed that seed yield of lentil (2.11 t ha⁻¹) was maximum with USG placed at 20 cm distance (avoid one row) and minimum values were recorded from USG placed at 40 cm distance (avoid three rows). Rumana (2013) also found that the highest seed yield (1.98 t ha⁻¹) from USG placed at 20 cm distance (avoid two rows), while the lowest seed yield (1.27 t ha⁻¹) was observed from USG placed at 40 cm distance (avoid three rows).

Table 6. Effect of application of prilled urea and urea super granule on yield contributing characters and yield of grasspea

Treatment	Number of pods plant ⁻¹	Number of seeds pod ⁻¹	Weight of 1000 seeds (g)	Seed yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Harvest index (%)
T ₁	52.55 a-c	4.61 ab	39.67 ab	1.85 ab	2.81 ab	39.70 ab
T ₂	49.81 b-e	4.33 b	37.85 a-c	1.55 cd	2.68 a-c	36.68 a-d
T ₃	46.61 d-f	3.90 c	36.08 cd	1.74 a-c	2.76 a-c	38.60 ab
T ₄	49.07 c-f	4.37 b	35.78 cd	1.32 de	2.33 de	36.17 b-d
T ₅	54.05 ab	4.67 ab	40.20 a	1.87 ab	2.85 ab	39.66 ab
T ₆	51.75 a-c	4.57 ab	38.49 a-c	1.76 a-c	2.80 a-c	38.69 ab
T ₇	54.48 a	4.87 a	40.57 a	1.94 a	2.92 a	39.97 a
T ₈	45.57 ef	4.43 b	38.21 a-c	1.33 de	2.50 cd	34.74 cd
T ₉	51.72 a-c	4.50 b	38.24 a-c	1.69 a-c	2.69 a-c	38.25 a-c
T ₁₀	49.85 b-e	3.80 c	37.09 b-d	1.65 bc	2.61 a-d	38.78 ab
T ₁₁	45.73 ef	4.40 b	37.95 a-c	1.53 cd	2.65 a-c	36.60 a-d
T ₁₂	45.26 f	3.63 cd	35.76 cd	1.30 de	2.51 cd	34.12 cd
T ₁₃	50.50 a-d	4.37 b	36.66 cd	1.32 de	2.54 b-d	34.12 d
T ₁₄	45.09 f	3.43 d	34.79 d	1.19 e	2.11 e	36.02 b-d
CV(%)	8.34	4.56	5.80	9.40	4.68	5.98

Mean having same or without letter(s) do not differ at 5% level of probability

T₁ = Prilled urea (PU) broadcasted

T₄ = USG placed at 10 cm distance (avoid two rows)

T₇ = USG placed at 20 cm distance (avoid two rows)

T₁₀ = USG placed at 30 cm distance (avoid two rows)

T₁₃ = USG placed at 40 cm distance (avoid two rows)

USG placed at 10 cm depth of each case

T₂ = PU given between two rows

T₅ = USG placed at 10 cm distance (avoid three rows)

T₈ = USG placed at 20 cm distance (avoid three rows)

T₁₁ = USG placed at 30 cm distance (avoid three rows)

T₁₄ = USG placed at 40 cm distance (avoid three rows)

T₃ = USG placed at 10 cm distance (avoid one row)

T₆ = USG placed at 20 cm distance (avoid one row)

T₉ = USG placed at 30 cm distance (avoid one row)

T₁₂ = USG placed at 40 cm distance (avoid one row)

4.2.5 Stover yield ha⁻¹

Stover yield ha⁻¹ of grasspea varied significantly due to the application of prilled urea and urea super granules (Appendix VII). The highest stover yield was recorded from T₇ (2.92 t ha⁻¹) treatment, which was statistically similar with T₅ (2.85 t ha⁻¹), T₁ (2.81 t ha⁻¹), T₆ (2.80 t ha⁻¹), T₃ (2.76 t ha⁻¹) and T₉ (2.69 t ha⁻¹) treatment. The lowest stover yield was observed from T₁₄ (2.11 t ha⁻¹), which was similar with T₁₂ (2.51 t ha⁻¹) treatment (Table 6). Treatment T₇ produced 51.05% higher seed yield over T₁₄ treatments. Srinivas *et al.* (2002) observed that stover yield of green gram increased with increasing rates of N up to 40 kg ha⁻¹.

4.2.6 Biological yield ha⁻¹

Application of prilled urea and urea super granules showed statistically significant variation for biological yield ha⁻¹ of grasspea (Appendix VII). The highest biological yield was found from T₇ (4.86 t ha⁻¹), which was statistically similar with T₅ (4.72 t ha⁻¹), T₁ (4.66 t ha⁻¹), T₆ (4.56 t ha⁻¹), T₃ (4.49 t ha⁻¹) and T₉ (4.38 t ha⁻¹), whereas the lowest biological yield was observed from T₁₄ (3.29 t ha⁻¹) which was similar with T₁₂ (3.81 t ha⁻¹) (Figure 3). Srinivas *et al.* (2002) observed that stover yield increased with increasing rates of N up to 40 kg ha⁻¹. Vadavia *et al.* (1991) found that application of 20 kg ha⁻¹ N and 40 P ha⁻¹ increased significant straw yield of grasspea.

4.2.7 Harvest index

Non-significant variation was recorded for harvest index (HI) due to the application of prilled urea and urea super granules (Appendix VII). HI differ from 34.14 to 39.97%. The highest harvest index was recorded from T₇ (39.97%) treatment followed by T₁, T₂, T₃, T₅, T₆, T₉, T₁₀ and T₁₁ treatments. Higher HI in T₇ treatment was due to increased dry matter accumulation resulting from physiological capacity and mobilization or photosynthates of treatments. The lowest harvest index (34.12%) was obtained from T₁₃. Chaudhari *et al.* (1998) found that application of 20-40 kg N ha⁻¹ significantly influenced harvest index of grasspea.




Figure 3. Effect of application of prilled urea and urea super granules on biological yield of grasspea ($S_x = 0.228$)

CHAPTER V

SUMMARY AND CONCLUSION

The experiment was conducted at the farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka during the period from November, 2012 to March, 2013 to study the response of grasspea to the application of prilled urea and urea super granule. The variety BARI Grasspea 3 was used as test crop. The experiment consisted of the following treatments: T₁= Prilled urea (PU) broadcasted; T₂ = PU given between two rows; T₃ = Urea super granule (USG) placed at 10 cm distance (avoid one row); T₄ = USG placed at 10 cm distance (avoid two rows); T₅ = USG placed at 10 cm distance (avoid three rows); T₆ = USG placed at 20 cm distance (avoid one row); T₇ = USG placed at 20 cm distance (avoid two rows); T₈ = USG placed at 20 cm distance (avoid three rows); T₉= USG placed at 30 cm distance (avoid one row); T₁₀ = USG placed at 30 cm distance (avoid two rows); T₁₁ = USG placed at 30 cm distance (avoid three rows); T₁₂ = USG placed at 40 cm distance (avoid one row); T₁₃ = USG placed at 40 cm distance (avoid two rows) and T₁₄ = USG placed at 40 cm distance (avoid three rows). USG placed at 10 cm depth of each case. The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. Data on different crop growth characters, yield contributing characters and yield of grasspea were recorded and found significant differences for treatment variables.

At 30, 50, 70, 90 DAS and at harvest, the tallest plant (34.64, 40.30, 47.35, 59.32 and 65.49 cm) was observed from T₇ [USG placed at 20 cm distance (avoid two rows)] and the shortest plant (19.34, 27.30, 30.55, 38.11 and 57.01 cm) was recorded T₁₄ [USG placed at 40 cm distance (avoid three rows)]. At 30, 50, 70, 90 DAS and at harvest, the maximum branches plant⁻¹ (4.47, 9.50, 17.00, 22.55 and 25.77) was recorded from T₇ treatment, while the minimum number (2.27, 6.13, 10.37, 12.22 and 14.56) was recorded from T₁₄ treatment. At 30, 50, 70 and 90 DAS the highest dry matter content plant⁻¹ (8.64, 11.65, 20.45 and 40.55 g) was

recorded from T₇, whereas the lowest dry matter content plant⁻¹ (4.51, 6.27, 13.63 and 24.58 g) from T₁₄.

The highest CGR value was attained from T₇ (6.54 g m⁻²day⁻¹) at 30-50 DAS while the lowest CGR was found from T₅ (3.65 g m⁻²day⁻¹) treatment. At 50-70 DAS, the highest CGR was attained from T₁ (9.87 g m⁻² day⁻¹) whereas the lowest CGR was found from T₉ (4.52 g m⁻²day⁻¹). At 70-90 DAS, the highest CGR was attained from T₇ (20.09 g m⁻²day⁻¹), while the lowest from T₈ (10.62 g m⁻²day⁻¹), which was similar with T₂ (11.50 g m⁻²day⁻¹) and T₃ (12.54 g m⁻²day⁻¹). At 30-50 DAS, the highest RGR was found from T₇ (0.148 g g⁻¹ day⁻¹), whereas the lowest RGR was found from T₁₄ (0.124 g g⁻¹ day⁻¹). At 50-70 DAS, the highest RGR was found from T₃ (0.195 g g⁻¹ day⁻¹), whereas the lowest RGR was recorded from T₉ (0.159 g g⁻¹ day⁻¹). At 70-90 DAS, the highest RGR was found from T₇ (0.231 g g⁻¹ day⁻¹), whereas the lowest RGR was found from T₈ (0.203 g g⁻¹ day⁻¹).

The minimum days from emergence to flowering (45.33 days) was recorded from T₇ treatment while the maximum days (57.67 days) was found from T₁₄. The highest pods plant⁻¹ (54.48) was recorded from T₇, while the lowest pods plant⁻¹ (45.09) was observed from T₁₄. The highest seeds pods⁻¹ (4.87) was counted from T₇ treatment while the lowest number (3.43) from T₁₄ treatment. The maximum weight of 1000-seeds (40.57 g) was found from T₇, while the lowest weight of 1000-seeds (34.79 g) from T₁₄. The highest seed yield (1.94 t ha⁻¹) was recorded from T₇ treatment and the lowest (1.19 t ha⁻¹) from T₁₄ treatment. The highest stover yield (2.92 t ha⁻¹) was measured from T₇ and the lowest stover yield (2.11 t ha⁻¹) was observed from T₁₄. The highest biological yield (4.86 t ha⁻¹) was found from T₇ and the lowest (3.29 t ha⁻¹) from T₁₄ treatment. The highest harvest index (39.97%) was recorded from T₇, while the lowest harvest index (34.12%) was attained from T₁₃ treatment.

It may be concluded that grasspea plant gave maximum seed yield (1.94 t ha^{-1}) with USG placed at 10 cm depth at 20 cm distance (avoid two rows) which was similar to that of broadcasted prilled urea (1.85 t ha^{-1}) but 4.87% higher than broadcasted prilled urea. This USG treatment facilitated plant to uptake maximum nitrogen for better growth, development and yield of grasspea.

Such experiment could be tested further in different grasspea cultivated areas of Bangladesh to justify the present findings under different growing condition as this is a very new approach of using USG in grasspea crops.

REFERENCES

- Abbasi, A., Jafari, D. and Sharifi, R. S. (2013). Nitrogen rates effects and seed inoculation with *Rhizobium leguminosarum* and plant growth promoting rhizobacteria (PGPR) on yield and total dry matter of chickpea (*Cicer arietinum* L.). *Tec. J. Eng. Appl. Sci.*, **3**(23): 3275-3280.
- Ahmed, S., Mahboob, A. and Latif, M. (2010). Effect of USG as a source of urea on the grain yield of hybrid maize. *Asian J. Pl. Sci.*, **9**(1): 44-49.
- Akter, R. (2010). Response of mustard varieties to different nitrogen management. MS Thesis, Dept. Agron., Sher-e-Bangla Agril. Univ., Dhaka. p. 1-41.
- Alam, B. M. R. (2002). Effect of different level of Urea Super Granule on growth and yield of three variety of boro rice. MS thesis, Dept. Agron., Bangladesh Agril. Univ, Mymensingh. p. 119.
- Alam, M. Shahe, Islam, M. Saiful and Islam, M. A. (2000). Farmers' efficiency enhancement through input management. The issue of USG application in modern rice. *Bangladesh J. Agril. Res.*, **36**(1): 129-141.
- Aliloo, A. A., Khorsandy, H., Mustafavi, S. H. (2012). Response of chickpea (*Cicer arietinum* L.) cultivars to nitrogen applications at vegetative and reproductive stages. *Cereal Agron. Moldova*. **45**(4): 49-59.
- Anonymous. (1989). Annual Report 1987-88. Bangladesh Agricultural Research Institute. Joydebpur, Gazipur. p. 133.
- Arvadia, M. K. and Patel, Z. G. (1988). Influence of date of sowing on the growth and yield of chickpea (*Cicer arietinum* L.) under different fertility levels. *Gujrat Agric. Univ. Res. J.*, **13**(2): 65-66.

- Arya, M. P. S. and Kalra, G. S. (1988). Effect of phosphorus doses on growth, yield and quality of summer mungbean (*Vigna radiata*) and soil nitrogen. *Indian J. Agric. Res.*, **22**(1): 23-30.
- Bali, A. S., Sing, K. N., Shah, M. H. and Khandey, B. A. (1991). Effect of nitrogen and phosphorus fertilizer on yield and plant characters of mungbean (*Vigna radiata*) under the late sown condition of Kasmir valey. *Fert. Newsl.*, **36**(7): 59-61.
- BARC (Bangladesh Agricultural Research Council). (2005). Fertilizer Recommendation Guide. Bangladesh Agriculture Research Council, Farmgate, Dhaka. p. 219.
- BARC. (2005). Fertilizer Recommendation Guide. Bangladesh Agriculture Research Council (BARC), Farmgate, Dhaka. 218 p.
- BBS (Bangladesh Bureau of Statistics). (2013). Statistical Yearbook of Bangladesh. Bangladesh Bureau of Statistics. Statistics Division, Ministry of Planning, Govt. of Peoples Republic of Bangladesh. Dhaka. Bangladesh. p. 54.
- Bhale, V. M. and Salunke, V. D. (1993). Response of upland irrigated rice to nitrogen applied through urea, and urea, super granules, *Mysore J. Agril. Sci.*, **27**(1): 1-4.
- Bhardwaj, P. and Singh, S. R. (1993). Influence of USG on the growth and yield mungbean. *J. Indian Soc. Soil Sci.*, **12**(1): 56-62.
- Bhopal, S. and Singh, B. (1990). Note on response of gardenpea to N and P application in North Hills. *Indian J. Hort.*, **47**(1): 107-108.
- Chakavorti, S. P., Chalam, A. B. and Mohanty, S. K. (1989). Comparative efficiency of prilled urea and urea super granule for wetland rice. *J. Indian Soc. Soil. Sci.*, **37**(1): 177-179.

- Chaudhari, R. K., Patel, T. D., Patel, J. B. and Patel R. H. (1998). Response of chickpea cultivars to irrigation, nitrogen and phosphorus on sandy loam soil. *Intl. Chickpea Newsl.*, **5**: 24-26.
- Chauhan, P. P. and Mishra, R. C. (1989). Effect of nitrogen on dry matter and nutrient accumulation pattern in wheat (*Triticum aestivum*) under different sowing dates. *Indian J. Agron.*, **12**(1): 123-141.
- Dahiya, S. S., Faroda, A. S., Hooda, I. S. and Singh, D. P. (1989). Fertilizer requirement of gram under rainfed condition. *Haryana J. Agron.*, **5**(1): 65-66.
- Dahiya, S. S., Singh, M. and Singh, M. (1993). Relative growth performance of chickpea genotypes to irrigation and fertilizers application. *Haryana J. Agron.*, **9**(2): 172-175.
- Das, S. and Singh, T. A. (1994). Nitrogen use efficiency by rice and flood water parameters as affected by fertilizer placement techniques. *J. Indian Soc. Soil Sci.*, **42**(1): 46-50.
- Deolankar, K. P. (2005). Effect of fertigation on growth and yield of chickpea. *J. Maharashtra Agril. Uni.*, **30**(2): 170-172.
- FAO (Food and Agricultural Organization). (2006). Yearly yield of grasspea 2005. FAO Stat. Citation, 2006. Pp.34-38.
- Farzana, A. (2013). Growth and yield variations in lentil fertilized with different form of urea. MS thesis, Dept. Agron., Sher-e Bangla Agril. Univ, Mymensingh. p. 76.
- Gomez, K. A. and Gomez, A. A. (1984). Statistical procedures for Agricultural Research. Jhon Wiley and Sons, New York. P. 83.
- Haque, S. A. (1998). Case study on USG in three villages of Tangail district, ATDP/IFDC, Dhaka. p. 1-40.

- Hossain M. Q. (2008). Effect of Urea Super Granule on the yield and quality of T–Aman rice. MS thesis, Dept. Agron., Sher-e Bangla Arril. Univ., Dhaka. P. 87.
- Humphreys, E., Chalk, P. M., Muirhead, W. A. and White, R. J. G. (2006). Nitrogen fertilization of dry-seeded rice in South-east-Australia. *J. Earth and Env. Sci.*, **31**(2): 221-234.
- Hunt, R. (1978). Plant growth analysis. The institute of Biology's studies in Biology No. 96. Edward Arnold (Publishers) Limited, London, UK. p. 23.
- Hussain, N., Khan, M. B. and Ahmad, R. (2010). Improving wheat productivity in calcareous soils through band replacement of farmyard manure with phosphorus. *Intl. J. Agri Biol.*, **10**: 709-714.
- Islam, M. M. (2003). Effect of irrigation and N management on the performance of bushbean (*Phaseolus vulgaris* L.). M.S. thesis. Bangabondhu Shiekh Mujibor Rahman Agril. Uni., Gazipur-1706.
- Islam, M. M. and Black, R. P. (1998). Urea super granule technology - impact and action plan for 1988-89. Paper presented at the National Workshop on urea super granule (USG) Technology, held at BARC, Dhaka, Bangladesh. p. 29.
- Javiya, J. J., Ahlawat, R. P. S., Patel, J. C., Kaneria B. B. and Tank, D. A. (1989). Response of gram to irrigation under varying levels of nitrogen and phosphorus. *Indian J. Agron.*, **34**(4): 439-441.
- Jee, L. P. and Mahapatra, Y. K. (1989). Effect of rates, time and method of nitrogen application of rice. *Indian J. Agron.*, **33**(4): 422-432.
- Juang, J. and Wang, P. (1987). Nitrogen in rice both the broadcasting and deep placement. *J. Indian Soc. Soil. Sci.*, **34**(1): 123-125.

- Karadavut, U. and Ozdemir, S. (2001). Effect of rhizobium inoculation and nitrogen application on yield and yield characters of chickpea. *Tarla Bitkileri Bolumu*, **11**(1): 14-22.
- Karle, A. S. and Pawar, G. G. (1998). Effect of legume residue incorporation and fertilizer in mungbean-safflower cropping system. *J. Maharashtra Agril. Univ.*, **23**(3): 333-334.
- Khalil, M. I., Schmidhalter, U. and Inubushi, K. (2006). Fertilizer N movement and recovery by Spring wheat with Urea super granules point placed at different soil depth . 18th World Congress of Soil Science, July 9-15, 2006-Philadelphia, Pennsylvania, USA. pp. 56-59.
- Khan, H., Haqqani, A. M., Khan, M. A. and Malik, B. A. (1992). Biological and Chemical fertilizer studies in chickpea grown under arid conditions of Pakistan. *Sarhad J. Agric.*, **8**(3): 321-327.
- Khokar, R. K. and Warsi, A. S. (1987). Fertilizer response studies in gram. *Indian J. Agron.*, **32**(2): 362-364.
- Kumar, N. and Singh, C. M. (1983). Response of transplanted flooded rice to slow release forms of nitrogen in Kangra Valley of Himachal Pradesh. *Oryza*. **20**(2&3): 100-103.
- Lal, P., Gautam, R. C., Bisht, P. S. and Pandey, P. C. (1988). Agronomic and economic evaluation of urea super granule and sulphur coated urea in transplanted rice. *Indian J. Agron.*, **33**(2): 186-190.
- Mahadkar, U. V. and Saraf, C. S. (1988). Input response of the growth and yield performance of mungbean (*Vigna radiata* L., Wilezek) production. *Minia J. Agril. Res Dev. Egypt.*, **10**(1): 247-255.

- Mahboob, A. and Asghar, M. (2002). Effect of seed inoculation and different nitrogen levels on the grain yield of mungbean. *Asian J. Pl. Sci.*, **1**(4): 314-315.
- Malik, M. A., Saleem, M. F., Asghar, A. and Ijaz, M. (2003). Effect of nitrogen and phosphorus application on growth, yield and quality of mungbean (*Vigna radiate* L.). *Pakistan J. Agril. Sci.*, **40**(3): 133-136.
- Mansoor, M. (2007). Evaluation of various agronomic management practices for increased productivity of mungbean (*Vigna radiate* L.). Ph. D. Thesis, Depart. Agron., Faculty of Agric., Gomal University, D.I. Khan.
- Mishra, P. C., Sarker, A. K. and Mishra, B. (1994). Relative efficiency of different nitrogen sources in transplanted rice of Chotanagpur region. *J. Indian Soc. Soil Sci.*, **42**(2): 333-335.
- Mukesh, K. (2006). Impact of the starter doses of nitrogen on nodulation, yield and yield attributes of chickpea under irrigated conditions. *Intl. J. Agril. Sci.*, **2**(1): 253-255.
- Nadeem, M. A., Ahmad, R. and Ahmad, M. S. (2004). Effect of seed inoculation and different fertilizer levels on the growth and yield of mungbean (*Vigna radiata* L.). *Indian J. Agron.*, **3**(1): 40-42.
- Namvar, A., Sharifi, R. S., Khandan, T. and Moghadam, M. J. (2013). Seed inoculation and inorganic nitrogen fertilization effects on some physiological and agronomical traits of chickpea (*Cicer arietinum* L.) in irrigated condition. *J. Central European Agric.*, **14**(3): 28-40.
- Narayanan, A. M. and Thangamuthu, G. S. (1991). Effect of nitrogen management and nursery on grain and straw yields of rice. *Madras Agric. J.*, **78**(9-12): 310-312.

- Negi, S. C. (1992). Effect of nitrogen and phosphorus in temperate hill grown vegetable pea (*Psam sativum*). *Indian J. Agron.*, **37**(4): 772-774.
- Nigamananda, B. and Elamathi, S. (2007). Studies on the time of nitrogen, application foliar spray of DAP, and growth regulator on yield attributes, yield and economics of green gram (*Vigna radiata* L.). *Intl. J. Agril. Sci.*, **3**(1): 168-169.
- Oad, F. C. and Buriro, U. A. (2005). Influence of different NPK levels on the growth and yield of mungbean. *Indian J. Plant Sci.*, **4**(4): 474-478.
- Pandey, A. and Tiwari, K. L. (1996). Effect of prilled urea, modified urea and coted urea on transplanted rice. *Adv. Agric. Res. India*. **5**: 43-88.
- Patel, J. C., Malavia, D. D., Vyas, M. N. and Patel, K. P. (1989). Response of chickpea to irrigation and fertilizers. *Indian J. Agron.*, **34**(1): 67-69.
- Patel, J. S and Parmar, M. T. (1986). Response of green gram to varing levels of nitrogen and phosphorus. *Madras Agril. J.*, **73**(6): 355-356.
- Patel, L. R., Salvi, N. M. and Patel, R. H. (1992). Response of greengram (*Phaseolus vulgaris*) varieties to sulphur fertilization under different levels of nitrogen and phosphorus. *Indian J. Agron.*, **37**(4): 831-833.
- Patel, R. G., Palel, M. P., Palel, H. C. and Palel, R. B. (1984). Effect of graded levels of nitrogen and phosphorus on growth, yield and economics of summer mungbean. *Indian J. Agron.*, **29**(3): 42-44.
- Patel, S. R. and Chandrawanshi, B. R. (1986). Sources and methods of N application for drilled, rainfed lowland rice. *Intl. Rice. Res. Newsl.*, **11**(1): 26-32.
- Patel, S. R. and Mishra, V. N. (1994). Effect of different forms of urea and levels of nitrogen on the yield and nitrogen uptake of rice. *Adv. Pl. Sci.*, **7**(2): 29-34.

- Patra, S. K. and Padhi, A. K. (1989). Response of rice to sources, methods and levels of N. *Intl. Rice Res. Newsl.*, **14**(6): 20-27.
- Pongkao, V. C. and Inthong, M. A. (1988). Effect of nitrogen fertilizer on mungbean. *J. Agric. Sci.*, **33**(3): 14-19.
- Quayum, A. and Prasad, K. (1994). Performance of modified urea materials in rainfed lowland rice. *J. Res.*, **6**(2): 131-134.
- Raja, R. A., Hussain, M. M. and Reddy, M. N. (1987). Relative efficiency of modified urea materials for lowland rice. *Indian J. Agron.*, **32**(4): 46-53.
- Rajagopalan, S. and Palanisami, S. (1985). Effect of prilled urea and urea super granules in rapidly percolating soil. *Intl. Rice Res. Newsl.*, **14**(2): 28-29.
- Rajender, K., Sing, V. P., Sing, R. C. and Kumar, R. (2003). Monetary analysis on mungbean during summer season. *Annal Biol.*, **19**(2): 123-127.
- Rajender, K., Singh, V. P., Singh, R. C. (2002). Effect of N and P fertilization on summer planted mungbean (*Vigna radiata* L.). *Crop Res. Hisar.*, **24**(3): 467-470.
- Rama, S., Reddy, G. and Reddy, K. (1989). Effect of levels and source of nitrogen on rice. *Indian J. Agron.*, **34**(3): 364-366.
- Raman, R. and Venkataramana, K. (2006). Effect of foliar nutrition on NPK uptake, yield attributes and yield of greengram (*Vigna radiata* L.) *Crop Res. Hisar.*, **32**(1): 21-23.
- Rambabu, P., Pillai, K. G. and Reddy, S. N. (1983). Effect of modified urea materials and their methods of application on dry matter production, grain yield and nitrogen uptake in rice. *Oryza*. **20**(2&3): 86-90.
- Rao, B. R., Rajeswara, P., Singh Kailash, Bhattacharya, A. K. and Naqvi, A. A. (2004). Effect of prilled urea and modified urea materials on yield and

- quality of geranium (*Pelargonium graveolens* L. (Her). *J. Earth Env. Sci.*, **23**(2): 81-85.
- Rao, D. L. N. and Ghai, S. K. (1989). Slow release urea fertilizers in sodic soils. *Intl. Rice Res. Newsl.*, **12**(1): 32-39.
- Rathore, A. L. and Patel, S. L. (1991). Response of late sown chickpea to method of sowing, seed rate and fertilizer. *Indian J. Agron.* **36**(2): 180-183.
- Reddy, G. R. S., Reddy, G. B., Ramaiah, N. V. and Reddy, G. V. (1991). Effect of different levels of nitrogen and form of urea on growth and yield of wetland rice. *Indian J. Agron.*, **31**(2): 195-197.
- Reddy, N. R. N. and Ahlawat, I. P. S. (1998). Response of chickpea (*Cicer arietinum*) genotypes to irrigation and fertilizers under late-sown condition. *Indian J. Agron.*, **43**(1): 95-101.
- Rumana, A. (2013). Performance of prilled urea and urea super granules in chickpea cultivation. MS thesis, Dept. Agron., Sher-e Bangla Agril. Univ, Mymensingh. p. 71.
- Saikia, U. S., Chopra, U. K., Singh, A. K. and Goswami, B. (2002). Simulation of biomass and seed yield of Indian mustard (*B. Juncea*) under different levels of nitrogen. *Ann. Agric. Res.*, **23**(4): 685-691.
- Sardana, H. R. and Verma, S. (1987). Combined effect of insecticide and fertilizers on the growth and yield of mungbean (*Vigna radiata* (L.) Wilczek). *Indian J. Entom.*, **49**(1): 64-68.
- Sarder, N. A., Shamusuddin, A. M. and Khan, N. H. (1988). Yield and yield component of wetland rice under various sources and levels of nitrogen. *Philippine J. Crop Sci.*, **13**(3): 155-158.
- Sarkar, R. K. and Banik, P. (1991). Response of mungbean (*Vigna radiata*) to nitrogen, phosphorus and molybdeum. *Indian J. Agron.*, **36**(1): 91-94.

- Shamim, S. and Naimat, A. (1987). Effect of seed inoculation with Rhizobium and NP fertilizer levels on the yield of gram. *Pakistan J. Agril. Res.*, **8**(4): 383-386.
- Sharma, C. K. and Sharma, H. K. (1999). Effect of different production factors on growth, yield and economics of mungbean (*Vigna radiata* L. Wilezeck). *Hill Farming.*, **12**(1&2): 29-31.
- Sharma, S. K. and Sharma, S. N. (2006). Effect of different combinations of inorganic nutrients and farmyard manure on the sustainability of a rice-wheat-mungbean cropping system. *Acta Agronomica Hungarica.* **54**(1): 93-99.
- Singh, B. K. and Singh, R. P. (1986). Effect of modified urea materials on rainfed low land transplanted rice and their residual effect on succeeding wheat crop. *Indian J. Agron.*, **31**(2): 198-200.
- Singh, B. K. and Singh, R. R. (1984). Relative efficiency of forms of urea and fates of nitrogen in rainfed low land rice on calcareous soil. *Indian J. Agron.*, **29**(4): 566-568.
- Singh, G., Singh, O. P., Yadav, R. A., Singh, R. S. and Singh, B. B. (1993). Effect of sources and levels of nitrogen on grain yield, yield attributes, N uptake, recovery and response by rice in deep water conditions (100 cm). *Crop Res.*, **6**(2): 213-216.
- Singh, K. and Singh, D. V. (2006). Effect of rates and sources of nitrogen application on yield and nutrient uptake of citronella Java. *J. Earth Env. Sci.*, **33**(3): 187-191.
- Singh, R. U. and Mishra, S. P. S. (1993). Response of Indian mustard varieties to nitrogen. *Indian J. Agron.*, **41**(2): 338-344.

- Srinivas, M., Shaik, M. and Mohammad, S. (2002). Performance of greengram (*Vigna radiata* L. Wilczek) and response functions as influenced by different levels of nitrogen and phosphorus. *Crop Res. Hisar.*, **24**(3): 458-462.
- Subba Rao, N. S., Tilak, K. V. B. R. and Singh, C. S. (1986). Dual inoculation with *Rhizobium* sp. And *Glamus fasciculatum* enhances nodulation, yield and nitrogen fixation in chickpea. *Plant and Soil.*, **59**(6): 402-404.
- Subbaiah, S. V., Kumar, R. M. and Pillai, K. G. (1994). Relative performance of granular fertilizers in irrigated rice (*Oriza sativa*). *Indian J. Agril. Sci.*, **64**(4): 255-256.
- Sudhakara, K. and Prasad, R. (1986). Relative efficiency of prilled urea, Urea super granules (USG) and USG coated with neem cake or DCD for direct seeded rice. *J. Agrc. Sci.*, **106**: 185-190.
- Tadesse, S. L. (1977). Response of mungbean cultivars to soil moisture stress at different growth phases. In: Proceeding of the first international Symposium, Mungbean. p. 122-127.
- Talukder, M. A. H., Mannaf M. A., Jabber, S. M. A., Islam, M. B., Kamal, S. M. A. M. and Shaha, A. K. (2004). Effect of Urea super granule as a source of Nitrogen on the growth and yield of tomato. *Pakistan J. Biol. Sci.*, **7**: 2078-2081.
- Tank, U. N., Damor, U. M., Patel, J. C. and Chauhan, D. S. (1992). Response of summer mungbean (*Vigna radiata*) to irrigation, nitrogen and phosphorus. *Indian J. Agron.*, **37**(4): 833-835.
- Tickoo, J. L., Naresh, C., Gangaiah, B. and Dikshit, H. K. (2006). Performance of mungbean (*Vigna radiata*) varieties at different row spacings and nitrogen-phosphorus fertilizer levels. *Indian J. Agric. Sci.*, **76**(9): 564-565.

- Tomar, N. K. and Sharma, J. C. (1985). Response and economic of N, P and K application in gram (*Cicer arietinum*). *Legume Res.*, **8**(2): 73-76.
- Trung, B. C. and Yoshida, S. (1983). Significance and nitrogen nutrition on the productivity of mungbean (*Vigna radiata* L. Wilczek). *Japanese J. Crop Sci.*, **52**(4): 493-499.
- Umar, A. P., Mamaril, C. P., Wahid, A. S. and Momuat, C. J. S. (1981). Methods of increasing efficiency of nitrogen fertilization in lowland rice. *In* : Proc. first Symp. on the role of food crops research findings to the agricultural development in Indonesia, Bogor, *Indonesia Agric. Sci.*, p. 42.
- Vadavia, A. T., Kalaria, K. K., Patel, J. C. and Baldha, N. M. (1991). Influence of organic, inorganic and biofertilizers on growth, yield and nodulation of chickpea. *Indian J. Agron.*, **36**(2): 263-264.
- Vijai, B., Singh, T. and Bahadur, V. (1990). Yield and growth response of gardenpea (*Pisum sativum*) to nitrogen and phosphorus application. *Indian J. Veg. Sci.*, **17**(2): 205-209.
- Vijaya, D. and Subbaiah, S. V. (1997). Effect of methods of application of granular forms of fertilizers on growth, nutrient uptake and yield of paddy. *Annal. Agril. Res.*, **18**(3): 361-364.
- Yakadri, M., Thatikunta, R. and Rao, L. M., Thatikunta, R. (2002). Effect of nitrogen and phosphorus on growth and yield of greengram (*Vigna radiata* L. Wilczek). *Legume Res.*, **25**(2): 139-141.
- Yamada, Y., Ahamed, S., Alcantara, A. and Khan, N. H. (1981). Nitrogen efficiency study under flooded paddy conditions: a review of inputs study. China, Inst. Soil. Sci., Academia Sinica, Nanjing: *Proc. Symp. on paddy Soil*. p. 588-596.

APPENDICES

Appendix I. Monthly average of air temperature, relative humidity and total rainfall of the experimental site during the period from November, 2012 to March, 2013

Month	Air temperature (°C)		Relative humidity (%)	Rainfall (mm) (total)
	Maximum	Minimum		
November, 2012	25.8	16.0	78	00
December, 2012	22.4	13.5	74	00
January, 2013	25.2	12.8	69	00
February, 2013	27.3	16.9	66	39
March, 2013	31.7	19.2	57	23

Source: Bangladesh Meteorological Department (Climate & weather division) Agargoan, Dhaka - 1212

Appendix II. Characteristics of soil of experimental field

A. Morphological characteristics of the experimental field

Morphological features	Characteristics
Location	Agronomy field , SAU, Dhaka
AEZ	Madhupur Tract (28)
General Soil Type	Shallow red brown terrace soil
Land type	High land
Soil series	Tejgaon
Topography	Fairly leveled

B. Physical and chemical properties of the initial soil

Characteristics	Value
% Sand	27
% Silt	43
% clay	30
Textural class	silty-clay
pH	5.6
Organic matter (%)	0.78
Total N (%)	0.03
Available P (ppm)	20.00
Exchangeable K (me/100 g soil)	0.10
Available S (ppm)	45

Source: Soil Resources Development Institute (SRDI), Khamarbari, Farmgate, Dhaka

Appendix III. Analysis of variance of the data on plant height of grasspea as influenced by application of prilled urea and urea super granule

Source of variation	Degrees of freedom	Mean square				
		Plant height (cm)				
		30 DAS	50 DAS	70 DAS	90 DAS	At harvest
Replication	2	1.991	0.343	1.890	0.526	1.197
Treatment	13	172.77 **	95.397**	203.432**	253.53**	54.399**
Error	26	2.013	2.123	5.673	1.189	6.409

** : Significant at 0.01 level of probability; * : Significant at 0.05 level of probability

Appendix IV. Analysis of variance of the data on number of branches plant⁻¹ of grasspea as influenced by application of prilled urea and urea super granule

Source of variation	Degrees of freedom	Mean square				
		Number of branches plant ⁻¹				
		30 DAS	50 DAS	70 DAS	90 DAS	At harvest
Replication	2	0.164	0.399	0.027	0.026	0.055
Treatment	13	8.920**	28.267**	74.772**	101.442**	5.087**
Error	26	0.254	0.602	1.315	1.378	0.0393

** : Significant at 0.01 level of probability; * : Significant at 0.05 level of probability

Appendix V. Analysis of variance of the data on dry matter content plant⁻¹ of grasspea as influenced by application of prilled urea and urea super granule

Source of variation	Degrees of freedom	Mean square			
		Dry matter content plant ⁻¹			
		30 DAS	50 DAS	70 DAS	90 DAS
Replication	2	33.365	43.441	131.406	868.285
Treatment	13	477.520**	807.143**	1921.268**	8636.724**
Error	26	103.115	164.677	687.457	2851.914

** : Significant at 0.01 level of probability

Appendix VI. Analysis of variance of the data on crop growth rate and relative growth rate of grasspea as influenced by application of prilled urea and urea super granule

Source of variation	Degrees of freedom	Mean square					
		Crop Growth Rate (g m ⁻² day ⁻¹)			Relative growth rate (g g ⁻¹ day ⁻¹)		
		30-50 DAS	50-70 DAS	70-90 DAS	30-50 DAS	50-70 DAS	70-90 DAS
Replication	2	0.120	0.803	5.163	0.0001	0.0001	0.0001
Treatment	13	2.347**	7.007*	28.196**	0.0002**	0.0001	0.00015*
Error	26	0.772	4.008	11.450	0.0001	0.0001	0.0001

** : Significant at 0.01 level of probability; * : Significant at 0.05 level of probability

Appendix VII. Analysis of variance of the data on yield contributing characters, yields and harvest index of grasspea as influenced by application of prilled urea and urea super granule

Source of variation	Degrees of freedom	Mean square							
		Days from emergence to flowering	Pods plant ⁻¹	Seeds pod ⁻¹	Weight of 1000 seeds (g)	Seed yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)	Biological yield (t ha ⁻¹)	Harvest index (%)
Replication	2	1.234	0.781	0.013	1.824	0.005	0.003	0.011	1.262
Treatment	13	13.983**	6.891**	1.314*	14.892**	0.535**	0.456*	1.979**	21.987**
Error	26	1.872	1.901	0.342	3.891	0.023	0.027	0.075	3.793

** : Significant at 0.01 level of probability;

* : Significant at 0.05 level of probability