


COMBINING ABILITY AND HETEROSIS IN RAPE SEED



BY

BASHIR AHMED BHUIYAN

REGISTRATION NO.: 06-01902

A Thesis

*Submitted to the Faculty of Agriculture,
Sher-e-Bangla Agricultural University, Dhaka,
in partial fulfillment of the requirements
for the degree of*

MASTER OF SCIENCE

IN

GENETICS AND PLANT BREEDING

SEMESTER: JULY - DECEMBER, 2012

Approved by:

.....
(*Dr. Md. Shahidur Rashid Bhuiyan*)

**Professor
Supervisor**

.....
(*Dr. Jamilur Rahman*)

**Associate Professor
Co-supervisor**

.....
(*Associate Prof. Dr. Mohammad Saiful Islam*)

**Chairman
Examination Committee**



Dr. Md. Shahidur Rashid Bhuiyan
Professor

Department of Genetics and Plant Breeding
Sher-e-Bangla Agricultural University
Dhaka-1207, Bangladesh


Mob: +88-01552467945

CERTIFICATE

This is to certify that thesis entitled, "*Combining Ability and Heterosis in Rape Seed*" submitted to the Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka, in partial fulfillment of the requirements for the degree of **MASTER OF SCIENCE IN GENETICS AND PLANT BREEDING**, embodies the result of a piece of bonafide research work carried out by **Bashir Ahmed Bhuiyan**, Registration No. **06-01902** under my supervision and guidance. No part of the thesis has been submitted for any other degree or diploma.

I further certify that such help or source of information, as has been availed of during the course of this investigation has duly been acknowledged.

Dated: December, 2012
Place: Dhaka, Bangladesh


(Dr. Md. Shahidur Rashid Bhuiyan)
Professor
Supervisor



**DEDICATED
TO
MY BELOVED PARENTS**

ACKNOWLEDGEMENTS

All praises to Almighty and Kindfull trust on to "Allah" for his never-ending blessing, it is a great pleasure to express profound thankfulness to my respected father and mother, who entiled much hardship inspiring for prosecuting my studies, thereby receiving proper education.

I would like to express my heartiest respect, my deep sense of gratitude and sincere, profound appreciation to my supervisor, Prof. Dr. Md. Shahidur Rashid Bhuiyan, Honorable Pro-Vice Chancellor, Sher-e-Bangla Agricultural University, Dhaka for his sincere guidance, scholastic supervision, constructive criticism and constant inspiration throughout the course and in preparation of the manuscript of the thesis.

I would like to express my heartiest respect and profound appreciation to my Co-supervisor, Dr. Jamilur Rahman, Associate Professor, Department of Genetics and Plant Breeding, Sher-e-Bangla Agricultural University, Dhaka for his utmost co-operation and constructive suggestions to conduct the research work as well as preparation of the thesis.

I am grateful to Prof. Md. Shadat Ulla, Honorable Vice Chancellor, Sher-e-Bangla Agricultural University, Dhaka for providing me with all possible help during my studies.

I am highly grateful to my honorable teacher Dr. Mohammad Saiful Islam , Chairman and Associate Professor, Department of Genetics and Plant Breeding, Sher-e-Bangla Agricultural University, Dhaka for his scholarly suggestions, constructive criticism, support and encouragement during the course of studies and for providing unforgettable help at the time of preparing the thesis.

I express my sincere respect to the teachers of Prof. Abu Akher Mia, Prof. Dr. Md. Sarowar Hossain, Prof. Dr. Naheed Zeba, Prof. Dr. Firoz Mahmud, Assit., Prof. Dr. Md. Ashaduzzaman Siddiquee, Assit. Prof. Md. Harun-Ur-Rashid and Lecturer Shahanaz Parveen Department of Genetics and Plant Breeding, Sher-e-Bangla Agricultural University, Dhaka for providing the facilities to conduct the experiment and for their valuable advice and sympathetic consideration in connection with the study.

I am specially thankful to Dr. Md. Motiar Rahman, Senior Scientific Officer, Plant Breeding Division, BARI, Gazipur for his helpful co-operation in compiling data and valuable suggestion to analysis the data and for giving technical support to prepare this thesis paper.

I am pleased to all staff and workers of Genetics and Plant Breeding Department and all farm labors and staff of Sher-e-Bangla Agricultural University, for their valuable and sincere help in carrying out the research work.

I feel much pleasure to convey the profound thanks to my Friends and Brothers specially, Sauda, Safin, Kamrul, Akkas, and all other friends and all well wishers for their active encouragement and inspiration. There are many others who helped and supported me in various ways. I sincerely thank to all of them and request their forgiveness for not mentioning here by name.

Mere diction is not enough to express my profound gratitude and deepest appreciation to my father (Abdul Momen Bhuiyan), mother (Begum Ayesha), brother Nasir Ahmed Bhuiyan and friends for their ever ending prayer, encouragement, sacrifice dedicated efforts to educate me to this level.

***Dated: December, 2012
Place: Dhaka, Bangladesh***

The Author



CONTENTS

CHAPTER	TITLE	PAGE NO.
	ACKNOWLEDGEMENT	i-ii
	CONTENTS	iii
	LIST OF TABLES	iv
	LIST OF FIGURES	v
	LIST OF PLATES	vi
	LIST OF APPENDICES	vii
	LIST OF ABBREVIATIONS AND SYMBOLS	viii-ix
	ABSTRACT	x
I	INTRODUCTION	1-3
II	REVIEW OF LITERATURE	04-35
	2.1 COMBINING ABILITY	04-19
	2.2 HETEROSIS	20-35
III	MATERIALS AND METHODS	36-49
	3.1 Experimental site and duration	36
	3.2 Plant Materials	36
	3.3 Land preparation and fertilizer application	39
	3.4 Experimental design and layout Methods	39
	3.5 Irrigation and drainage	39
	3.6 Intercultural operation, Insect and disease control	39
	3.7 Harvesting of sample plants	40
	3.8 Data recorded	43
	3.9 Statistical analysis	44-49
IV	RESULTS AND DISCUSSION	50-85
	4.1 Mean performance	50-52
	4.2 Combining ability	53-55
	4.2.1 General combining ability (GCA) effects	56-59
	4.2.2 Specific combining ability (GCA) effects	60-63
	4.3 Heterosis	64-75
	4.4 Vr-Wr Graph	76-85
V	SUMMARY AND CONCLUSION	86-87
	REFERENCES	88-99
	APPENDICES	100-101

LIST OF TABLES

Table No.	Title of the Tables	Page No.
1	Cross combinations in half diallel system of five varieties in <i>Brassica rapa</i> L.	38
2	Mean performance for 9 different characters in 5 parents and their 10 F ₁ 's in <i>Brassica rapa</i> L.	51
3	Analysis of variances (MS values) for seed yield per plant and its component characters in <i>Brassica rapa</i> L.	54
4	Analysis of variances (MS values) for GCA and SCA for seed yield and yield contributing components in <i>Brassica rapa</i> L	55
5	General combining ability (GCA) effects for 5 parents in 5x5 half diallel crosses of <i>Brassica rapa</i> L	57
6	Specific combining ability (SCA) effects for 5 parents in 5x5 half diallel crosses of <i>Brassica rapa</i> L	61
7	Percent heterosis over mid-parent and better parent for 9 different characters in intervarietal hybrids of oleiferous <i>Brassica rapa</i> L.	65-67

LIST OF FIGURES

Figure No.	Title of the figures	Page No.
1	Location of the experimental field	37
2	V _r -W _r graph for plant height in <i>Brassica rapa</i>	76
3	V _r -W _r graph for days to maturity in <i>Brassica rapa</i>	77
4	V _r -W _r graph for primary branches per plant in <i>Brassica rapa</i>	78
5	V _r -W _r graph for secondary branches per plant in <i>Brassica rapa</i>	79
6	V _r -W _r graph for siliqua per plant in <i>Brassica rapa</i>	80
7	V _r -W _r graph for length siliquae in <i>Brassica rapa</i>	81
8	V _r -W _r graph for seed/siliquae in <i>Brassica rapa</i>	82
9	V _r -W _r graph for seed yield per plant in <i>Brassica rapa</i>	83
10	V _r -W _r graph for 1000 seed weight in <i>Brassica rapa</i>	84



LIST OF PLATES

Plate No.	Title of the Plates	Page No
1	Photograph showing a field view of experimental site at seedling stage at SAU farm	41
2	Photograph showing a field view of experimental site at SAU farm	41
3	Photograph showing a field view of experimental site at flowering stage at SAU farm	42
4	Photograph showing a field view of experimental site at maturity stage at SAU farm	42
5	Photograph showing variation for plant height between SAU sarisha X BARI sarisha 15 and Tori 7 X SAU sarisha 2 of <i>Brassica rapa</i> genotype.	68
6	Photograph showing variation between early (SAU sarisha 3 X BARI sarisha 6) and late maturity (SAU sarisha 2 X BARI 15) line of <i>Brassica rapa</i> genotype	69
7	Photograph showing variation for number of primary branches / plant between SAU sarisha 2 X BARI sarisha 6 (lowest) and SAU sarisha 3 X BARI sarisha 15 (highest) of <i>Brassica rapa</i> genotype	70
8	Photograph showing variation for number of secondary branches / plant between SAU sarisha 2 X SAU sarisha 3 (lowest) and Tori 7 X BARI sarisha 15 (highest) of <i>Brassica rapa</i> genotype	71
9	Photograph showing variation for number of siliquae / plant between Tori 7 X BARI sarisha 15 (highest) and SAU sarisha 2 X SAU sarisha 3 (lowest) of <i>Brassica rapa</i> genotype	72
10	Photograph showing variation for siliqua length between SAU sarisha 2 X BARI sarisha 15 (highest) and Tori 7 X SAU sarisha 2 (lowest) of <i>Brassica rapa</i> genotype	73

LIST OF APPENDICES

Appendix No.	Title of the appendix	Page No.
I.	Morphological, physical and chemical characteristics of initial soil (0-15 cm depth) of the experimental site	100
II.	Monthly average temperature, relative humidity and total rainfall of the experimental site during the period from October, 2011 to March, 2012	101



SOME COMMONLY USED ABBREVIATIONS AND SYMBOLS

Abbreviations	=	Full word
°C	=	Degree Celsius
@	=	At the rate
σ_p^2	=	Phenotypic variance
σ_e^2	=	Environmental variance
σ_g^2	=	Genotypic variance
h_b^2	=	Heritability in broad sense
AEZ	=	Agro-Ecological Zone
Agric.	=	Agriculture
Agril.	=	Agricultural
Agron.	=	Agronomy
ANOVA	=	Analysis of variance
BARI	=	Bangladesh Agricultural Research Institute
BBS	=	Bangladesh Bureau of Statistics
BD	=	Bangladesh
BSMRAU	=	Bangabundhu Sheikh Mujibur Rahaman Agricultural University
CEC	=	Cation Exchange Capacity
CV%	=	Percentage of Coefficient of Variation
CVA	=	Canonical Variate Analysis
cv.	=	Cultivar (s)
D 50% F	=	Days to 50% flowering
DAS	=	Days After Sowing
df	=	Degrees of Freedom
DM	=	Days to maturity
Dm	=	Dry-matter
DMRT	=	Duncan's Multiple Range Test
EC	=	Emulsifiable Concentrate
<i>et al.</i>	=	And others
etc.	=	Etcetera
F ₈	=	The eight generation of a cross between two dissimilar homozygous parents
FAO	=	Food and Agricultural Organization
G	=	Genotype
GA	=	Genetic advance
GCV	=	Genotypic Coefficient of Variation
GN.	=	Genotype Number
HI	=	Harvest Index

Abbreviations		Full word
hr.	=	Hour (s)
HSW	=	Hundred seed weight
IARI	=	Indian Agricultural Research Institute
ICARDA	=	International Centre for Agricultural Research in Dry Areas
j.	=	Journal
kg	=	kilogram (s)
m	=	Meter
M.P.	=	Muriate of Potash
m ²	=	Square meter
MOA	=	Ministry of Agriculture
MSG	=	Mean square of the genotypes
MSE	=	Mean square of the error
NARS	=	National Agricultural Research System
No.	=	Number
NPB/P	=	Number of primary branches per plant
NSB/P	=	Number of secondary branches per plant
NS	=	Not Significant
NSP	=	Number of siliquae per plant
PCA	=	Principal Component Analysis
PCO	=	Principal Coordinate Analysis
PCV	=	Phenotypic Coefficient of Variation
PH	=	Plant height
ppm	=	Parts Per Million
R	=	Residual effect
RCBD	=	Randomized Complete Block Design
Rep.	=	Replication
Res.	=	Research
SAU	=	Sher-e-Bangla Agricultural University
Sci.	=	Science
SE	=	Standard Error
SL	=	Silique length
S/S	=	Seeds per silique
T.S.P.	=	Triple Super Phosphate
t/ha	=	Tones per hectare
Univ.	=	University
var.	=	Variety
Via	=	By way of
Viz	=	Namely
WP	=	Wetable powder
YPP	=	Yield per plant



COMBINING ABILITY AND HETEROSIS IN RAPE SEED

BY


BASHIR AHMED BHUIYAN

ABSTRACT

An experiment on oleiferous *Brassica rapa* L. was conducted to evaluate the combining ability and heterosis for nine different characters. The parent BARI sarisha 6 was the best general combiner for plant height, early maturity, number of primary branches, number of siliqua per plant, seed yield per plant and thousand seed weight. The parent BARI sarisha 15 was the best general combiner for number of primary branches per plant and number of seeds per siliqua. The parent Tori 7 was good general combiner for number of secondary branches per plant. The parent SAU sarisha 2 was the best for seed per siliqua. On the basis of average score and rank position, Tori 7 X BARI sarisha 15 was the best cross combination for plant height, no. of secondary branches per plant, and no. of siliquae per plant. The combination SAU sarisha 2 X BARI sarisha 15 was the best for days to early maturity, Siliqua length and seed per siliqua. The hybrid SAU sarisha 3 X BARI sarisha 15 was the best for no. of primary branches per plant and thousand seed weight. For seed yield per plant the cross BARI sarisha 6 X BARI sarisha 15 was the best. The higher magnitude of GCA variance was observed than that of SCA variance for plant height, days to maturity, no. of primary branches/ plant, no. of secondary branches/ plant, no. of siliquae/ plant, seed yield/ plant and 1000 seed weight which indicated the preponderance of additive component in their expression. The Vr-Wr graph indicates over dominance for days to maturity, primary branches per plant, secondary branches per plant, siliquae per plant, and seeds per siliqua. Partial dominance was observed for plant height, length of siliqua, seed yield per plant and thousand seed weight. The graphical analysis also indicated wide genetic diversity among the parents. Out of ten crosses, the hybrids SAU sarisha 2 X BARI sarisha 6 showed desirable negative heterosis for the character of short plant height. The hybrids SAU sarisha 3 X BARI sarisha 6 were best for early maturity. The hybrids Tori 7 X BARI sarisha 15, SAU sarisha 3 X BARI sarisha 15, Tori 7 X SAU sarisha 2 were found to be heterotic for no. of primary branches per plant, no. of secondary branches per plant and no. of siliquae per plant. While the cross Tori 7 X BARI sarisha 15 produced maximum no. of secondary branches and no. of siliquae per plant. Most of the crosses showed significant and negative performance for length of siliqua and no. of seeds per siliqua. The crosses SAU sarisha 2 X BARI sarisha 15 were best for siliqua length and the crosses SAU sarisha 3 X BARI sarisha 6 and Tori 7 X BARI sarisha 6 were best for seeds per siliqua. For thousand seed weight the hybrids SAU sarisha 3 X BARI sarisha 15, Tori 7 X BARI sarisha 15, BARI sarisha 6 X BARI sarisha 15 and Tori 7 X SAU sarisha 2 were best. For seed yield per plant the cross SAU sarisha 2 X BARI sarisha 15 was found to be the best.



Chapter 1
Introduction



INTRODUCTION

Rapeseed and mustard (*Brassica sp.*) is an important oil seed crop of Bangladesh belonging to the family Brassicaceae. *Brassica* is an important genus of plant kingdom consisting of over 3200 species with highly diverse morphology. It has great economic and commercial value. They range from nutritious vegetables, condiments and oil producing oleiferous *Brassica*. It is a cross pollinated and annual crop. The seeds of mustard and rapeseed contain 42% oil and 25% protein (Khaleque, 1985). Among the oil seed crops rapeseed and mustard is the third highest source of edible oil in the world after soybean and palm (Anonymous, 2000; Piazza and Foglia, 2001 and Walker and Booth, 2001).

The genus *Brassica* has been divided into three groups viz-rape seed, mustard and cole. The rape seed group includes the diploid *Brassica rapa*, turnip rape (AA, $2n=20$) and amphidiploid *Brassica napus* L, rape (AACC, $2n=38$) while the mustard group includes species like *Brassica juncea* Czern and Coss; *Brassica nigra* Koch and *Brassica carinata* Braun (Yarnell, 1956). All these species have many cultivated varieties suited to different agro-climatic conditions. In the oiliferous *Brassica* group, a considerable variation of genetic nature exists among different species and varieties within each species in respect of different morphological characters (Malik *et al.* 1995; Kakroo and Kumar 1991).

Brassica plays a major role in feeding the world population. The oil of *Brassica* is of variable quality with bad to good for health probably due to high to zero contents of erucic acid (Woyke, 1981) and high to low contents of glucosinolates (Li *et al.* 1990). Morphological traits are unique in nature and maintain speciality in a particular crop variety and their interrelationship has influence on yield.

Oil seed crop covers about 4.04% area of the total cultivable land in Bangladesh and rapeseed and mustard covers about 74.5% of that area (BBS, 2006a). This crop covers the maximum area of the oil yielding crop due to the climate and edaphic factors of Bangladesh are quite favorable for the cultivation of this crop. But the shortage of edible oil has become a chronic problem for the nation. Bangladesh requires 0.29 million tons of oil equivalent to 0.8 million tons of oilseeds for nourishing her people where the oilseed production is about 0.254 million tons, which covers

only 40% of the domestic need (FAO, 2001). As a result, more than 60% of the requirement of oil and oil seed has been imported every year by spending huge amount of foreign currency involving over 317 cores taka (BBS, 2006c). The average yield per hectare of mustard and rapeseed crop is 850-900 kg (BBS, 2006b) in Bangladesh compared to the world average of 1,575kg (FAO, 2003).

Although rape and mustard is most important oil crop in Bangladesh, farmers usually cultivate them in less fertile land followed by low management with least investment. Almost all the cultivars are brown seeded and smaller in size (2-2.5 g/ 1000seeds). Yellow seed contains 2-3% more oil than the same sized brown seeded type due to its thinner seed coat. Bold and yellow seeded rapeseed varieties may increase total edible oil production of Bangladesh. High yielding varieties in late sown condition having early maturity may increase 12-15% area of total edible oil seed of Bangladesh, when it replaces the total rapeseed and mustard grown in the country. The above scenario dictates the major quantitative and agronomic modification of this crop.

Meanwhile, about 26 mustard and rapeseed varieties have been released in Bangladesh, among these 15 from Bangladesh Agricultural Research Institute (BARI), five from Bangladesh Institute of Nuclear Agriculture (BINA), two from Bangladesh Agricultural University (BAU) and two from Sher-e-Bangla Agricultural University (SAU) and two from Bangladesh Agricultural Development Corporation (BADC) but most of them are not popular to the farming community because of their long duration, low to moderate yield and susceptibility to severe biotic and abiotic stresses.

So there is a plenty of scope to increase yield per unit of area through cultivation of short duration high yielding varieties. The production potential of rapeseed and mustard may be well exploited if the varieties can be identified with early maturity, rapid response to high fertility, has large seed size and oil content.

Intra-specific hybridization is a good way of improving the varieties of different natures by combining desired traits followed by selection of desired types. The most important aspects for hybridization are the choice of parents and the selection of best genotypes from hybrid progenies.

Availability of heterotic germplasm is prerequisite for successful hybrid breeding work of rape seed. Though mustard and rapeseed contribute major portion of oil requirement in our country we have a limitation to accelerate total yield due to lack of good performance of our existing varieties. So we have ample scope to improve our existing varieties by exploiting heterosis for different yield contributing characters. Thus heterosis should lead to increase in yield, adaptability, disease and insect resistance, general vigor, quality etc.

Combining ability studies are reliable as they provide useful information for the selection of parents in terms of performance of the hybrids and elucidate the nature and magnitude of various types of gene action involved in the expression of quantitative traits. Genetic information helps in the selection of suitable parents for hybridization and in isolating the promising early generation hybrids for further exploitation in breeding programs.

Information on heterosis and combining ability of hybrid progenies at its early generation are very useful for the purpose of selection criteria. So the present research was undertaken with the following objectives:

- To estimate the nature and extent of gene action controlling the traits in rapeseed.
- To study the extent of heterosis in genotype of *Brassica rapa*.



Chapter 2

Review of literature

REVIEW OF LITERATURE

In the field of *Brassica* breeding, many researchers have conducted research works on heterosis over mid parental values or better parental values and combining ability, a large volume of literature is available on topics. However, attempt has been made to review some of the literatures relevant to the present study on mustard in this chapter.

2.1 COMBINING ABILITY

General combining ability is the average performance of a given genotype in a series of hybrid combinations, while the specific combining ability is expressed through the performance of a parent in a specific cross in relation to the genotype. For the characters studied, both significant and insignificant results were noted in the literatures discussed in this chapter.

Yadav *et al.* (2005) found significant differences due to parents vs. crosses indicating the presence of heterosis in the crosses through conducted an experiment during the rabi seasons of 1998-2000 to study the nature of combining ability for seed yield and other yield-attributing characters through line \times tester analysis in rape (*Brassica napus*) [*B. napus* var. *oleifera*]). They derived forty-five F_1 from the crosses of two cytoplasmic male sterile lines (Ogura, ISN-706a) and one normal fertile line (NDBN-1) used as females and 15 testers (Westar, FM-27, GSL-6267, GSL-8814, EC129120, PBN 9501, NRCG-7, GSL-6067, HNS-4, GSL-1, GSL-406, NRCG-2, GSL-6303, NRCG-13 and NRCG-14) as males. Among lines, they observed significant differences for plant height and number of secondary branches per plant. Higher magnitude of variances due to testers compared to lines were observed for seed yield per plant, plant height, primary branches per plant, days to flower initiation, days to maturity and oil content. They also found that the estimates of SCA variances were higher than GCA (average) for all the characters studied, indicating the preponderance of non-additive type of gene action in the inheritance of these traits and the cross Ogura \times NRCG-13 showed high SCA effects for yield per plant which involved both good combining parents.

Nair *et al.* (2005) worked on combining ability in mustard [*Brassica juncea*] to identify the better parents (Pusa Bold, Rohini, TM-17, ACN-9 and PCR-7) on the basis of their combining ability and to isolate superior crosses for studying them in further generations. The analysis of variances indicated that variances due to lines were significant for plant height and variances due to the testers were highly significant for all traits except days to maturity indicating significant genetic variation. Rohini was identified as the superior parent for the improvement of siliquae number per plant and hence, may be used in breeding programmers for the improvement of this trait. The cross Seeta × Rohini was identified as the promising cross for yield and contributing characters.

Heterosis for seedling, physiological and morphological traits in three rape crosses derived from four genotypes (Ester, Rainbow, Range and Shiralee) and grown under irrigated and non-irrigated condition was determined in experiments conducted by Cheema *et al.* (2004) in Pakistan during 1999-2002. High heterosis for shoot length and fresh root weight of the crosses over the mid- and better parents was recorded under irrigated and non-irrigated conditions. The highest positive and significant heterosis for water potential over the better parent was recorded in Range × Ester under normal and drought conditions. Heterosis over the mid parent for chlorophyll was recorded in Range × Shiralee grown under normal and drought conditions. Range × Shiralee recorded high heterosis over the mid and better parent under drought conditions and high heterosis for yield over the mid parent under normal conditions.

Chowdhury *et al.* (2004) studied the nature and magnitude of combining ability of parents and crosses (F_{1s}) were estimated in a 7×7 diallel cross analysis in turnip rape for seed yield, its different contributing characters and oil content. Higher magnitudes of GCA variances were observed than those of sca variances for all the characters except siliquae per plant, seeds per siliqua and seed yield per plant. Majority of the crosses showed high SCA effects for seed yield involving high × low, average × average and average × low GCA parents.

Pietka *et al.* (2003) proposed that the general combining ability (GCA) values in terms of individual glucosinolates are important in breeding. Eleven inbred lines of winter oilseed rape (*B. napus*[var. *oleifers*]) characterized by very low glucosinolate contents were studied by them. These lines were crossed with five cultivars used as testers. Hybrids were grown in the field and

statistical analyses of GCA values were performed separately for particular glucosinolates, as well as F₁ and F₂ generations. Heritabilities of regressions were estimated by determining the coefficients between both generations. Most of the coefficients were significant at alpha =0.01 or 0.05, providing that the GCA estimation used in the experiments was satisfactorily reproducible.

Prasad *et al.* (2002) evaluated combining ability of 21 F₁ hybrids derived from a diallel cross of seven Indian cultivars along with the parents in a field experiment. The general and specific combining ability were significant for all the traits examined. The cultivar Varuna recorded high general combining ability for most of the characters and *per se* performance. The specific combining ability for early maturity, length of main raceme and yield per plant were observed in the crosses involving high × low GCA parents.

Liu *et al.* (2001) combining ability and heritability of eight main agronomic characters of the crosses obtained by crossing four double-low male sterile lines of rapeseed with glucosinolate lower than 30 micro mol/g and erucic acid lower than 1% with four good restorer lines based on North Carolina II design. They observed sterile line 121A, known as the sterile line of Shanyou 6, was shown to be most outstanding, with high general combining ability of many yield-contributing characters, thus having relatively high yield potential.

Matho and Haider (2001) worked with the magnitude of specific combining ability (SCA) effects was much higher than the general combining ability (GCA) effects for all the characters studied, except for number of secondary branches per plant. In most of the cases, the crosses showing high SCA effects also exhibited high heterosis.

Pietka *et al.* (2001) conducted an experiment to establish the relationship of general (GCA) and specific combining ability (SCA) with glucosinolate content in seeds collected from F₁ and F₂ hybrids generations of winter double row rapeseed. They examined that hybrids produced by crossing cultivars Mar, Polo, Silvia, Lirajet, and Wotan with inbred lines extremely low in glucosinolate content. They also found the calculated GCA values which showed that both inbred lines and cultivars were highly and significantly differentiated in terms of glucosinolate content and composition. They also suggested that an effective selection for low glucosinolate

content is possible for segregating hybrid populations and the possibility of using SCA in improving glucosinolate content was smaller than that of GCA.

Tak and Khan (2000) conducted an experiment to estimate the combining ability, magnitude of variability and gene effect of the available germplasm resources of 15 Indian mustard (*B. juncea*) lines crossed to three genetically different testers. Estimates of genetic variance revealed that the days to flowering was predominantly governed by a non-additive gene action. However both additive and non-additive gene actions were important in the inheritance of most of the characters studied. The line KS-216 showed significant general combining ability effect for earliness, whereas KS-240 and KS-181 were superior general combiners for seed yield.

Goffman and Becker (2001) stated that because of the nutritional and antioxidative properties, tocopherol production is an interesting trait for the lipid quality of oil crops. Total tocopherol content in rapeseed (*Brassica napus L.*) is medium to low, and therefore, higher levels of tocopherol are desirable in this species. The objective of the present study was to determine the inheritance of alpha-, gamma-, and total tocopherol content and the alpha -/ gamma -tocopherol ratio in seed of rapeseed. Two diallel mating designs with six parents each were used. In Diallel I, the parents selected were high or low for total tocopherol content and in Diallel II, the parents were high or low for the alpha -/ gamma -tocopherol ratio. Parents and F₁ hybrids were tested in a screenhouse in 1998 and under field conditions in 1999 by means of a completely randomized design with two replications. In addition, 10 selected F₂ populations were grown along with their respective parents. Compared with the parents, the F₁ hybrids showed a significantly higher gamma -tocopherol content of about 6 mg kg⁻¹ seed for Diallel I and 24 mg kg⁻¹ seed for Diallel II. General combining ability effects in both diallels were highly significant (P<0.01) and much larger than specific combining ability effects for all traits studied. Reciprocal effects were not statistically significant. Gamma-Tocopherol was not correlated with alpha -tocopherol. The results indicate that tocopherol content and composition inheritances are strongly associated with additive gene action in rapeseed.



Wos *et al.* (1999) presented general combining ability (GCA) and specific combining ability (SCA) for 23 cytoplasmic male sterility (CMS) ogura lines. Field trials were executed in four localities (Malyszyn, Marwice, Borowo and Bakow) in Poland. The seed yield of hybrids, GCA and SCA of CMS lines and GCA of pollinators were significant. 23 CMS ogura lines were crossed using three pollinator cultivars Kana, Marita and MAH 1592. Obtained results were used to find the best combinations for hybrid production.

Krzymaniński *et al.* (1999) examined combining ability and heterosis for selected eleven winter double low rape inbred lines (PN 3181/95, PN 3451/95, PN 3455/95, PN 3462/95, PN 3707/95, PN 3710/95, PN 3734/95, PN 3999/95, PN 4043/95, PN 4272/95 AND PN 4297/95) with extremely low glucosinolate content. Three foreign cultivars, Lirajet, Silvia, and Wotan, and two Polish cultivars, Mar and Polo, were used as testers. Crosses were made in both directions. The results of calculations made for the F₁ generation concern general and specific combining abilities with regard to parental form and 55 hybrid combinations and reciprocal effects. The results enabled the determination of the best combination of crosses. It was also proved that combining effects depend in some combinations on the direction of crossing.

Krzymaniński *et al.* (1999) made diallel (13x13) crossings of double low oilseed rape cultivars and strains. Parental forms and F₁ combinations of diallel were compared in field trials in Poland. Two cultivars and four strains were the parental forms that most frequently occurred in F₁ combinations yielding considerably above the standard cultivar (Bor), two strains gave combinations of the highest fat contents, considerably differing from the standard. The yields oscillated between 126.5 and 209.1% of the standard (38.2 q/ha) and the fat content between 103 and 108% of the standard (47%). Calculations were made to estimate the expected values of seed yield of synthetic varieties, which could be obtained from tested cultivars and strains. Two or three component synthetics composed from the best combining cultivars and strains were taken into account by them.

Wos *et al.* (2000) presented the results of the breeding studies on the development of winter and spring oilseed cytoplasmically male sterile (CMS) lines, restorers and composite hybrids performed at the Plant Breeding Station in Malyszyn (Poland) in collaboration with the Oil Crop

Department of Plant Breeding and Acclimatization Institute in Poznan. Some breeding aspects of the CMS lines, restorers and composite hybrids, including general combining ability and specific combining ability, contents of glucosinolates and erucic acid, winter hardiness and yield, are analysed. The results obtained so far have allowed the introduction of eight winter and four spring composite hybrids of oilseed rape to the State Official Trials. In 1999, the first Polish-French composite hybrid of spring rape named Margo was listed on the Polish Variety List.

Verma (2000) studied combining ability analysis of yield and its components through diallel crosses in indica coiza (*Brassica juncea* L.) Czern & Coss. the variance due to general (GCA) and specific combining ability (SCA) were estimated to assess the additive and non-additive gene action involved in the inheritance of nine characters in eight parents and F₁ hybrids of *Brassica juncea*. The parents RC 870, RC 759, RC 751, and RC 792 have shown higher GCA effects for seed yield and other characters. The best five crosses are RC 832 × RC 788, RC 827 × RC 870, RC 827 × RC 751, RC 837 × RC 870 and RC 832 × RC 870. These crosses are likely to give better sergeants in future generations.

Katiyar *et al.* (2000b) studied on heterosis for seed yield in Indian mustard (*Brassica juncea* (L) Czren. and Coss.). Six varieties and 16 lines of *B.juncea* in a tester mating design, and the resulting 96 crosses were evaluated for yield components. Seven combinations exhibited > 30% heterosis and eleven crosses showed 31.2-71.3% heterosis. It is concluded that there is adequate genetic divergence among Indian mustard lines to support a successful hybrid programme.

Huang *et al.* (2000) studied three rapeseed (*Brassica napus*) genotypes tolerant of resistant to *Sclerotinia sclerotiorum* and three susceptible genotypes differing in origin were used in reciprocal or complete diallel crosses and found that resistant genotype from China, 018, had the highest general combining ability (4.46) while the French variety Cobra had the lowest general combining ability (-10.54). They also found optimum cross combination in this study was Cobra 018, with high specific combining ability (10.41) and desirable agronomic characters.



Singh *et al.* (2000) worked with genetic analysis in yellow sarson, *Brassica campestris* L. They found significant differences for both SCA and GCA among the genotypes for all the characters indicating there by that both additive and non additive components were involving in the expression of all the traits. The parents with high GCA was showed good general combining ability for seed yield, days to maturity and siliqua per plant in both F₁ and F₂ generation and for primary and secondary branches per plant in F₂ generation only. The cross with high × low GCA effects showed significant SCA for seed yield.

Singh *et al.* (1999) studied the combining ability in *Brassica campestris* L. Comparison of SCA effects in relation of GCA effects of the respective parental lines indicated that crosses with high SCA effects involved low × high, high × low and low × low general combiners.

Sheikh and Singh (1998) analysis combining ability in 10 × 10 half-diallel (excluding reciprocals) of Indian mustard for ten characters and found preponderance of non additive gene action for most of the characters including seed yield and oil content. They also observed that Additive genetic variance was more important for plant height and length of silliqua. Majority of the crosses showed high SCA effects for seed yield involved high × low GCA parents.

Wos *et al.* (1998) presented the results of investigated general combining ability of 64 inbred lines and heterosis effects of winter oilseed rape F₁ hybrids. General combining ability was estimated by test topcrosses. Field experiments were designed in lattice design, in two replications (four rows per plot, three msuperscript two plot and sowing rate of 100 seeds per 1 msuperscript 2). The experiment was carried out in 1996-97. General combining ability (GCA) was significant for seed yield, 1000 seed weight, winter hardiness, beginning and end of flowering, oil and protein content. However, it has been proved that GCA was not significant for plant height. Results of these studies revealed: nine hybrids with significant higher yielding than tester (check) cv. Lirajet, 19 hybrids with significant better winter hardiness than tester, 35 hybrids with significant earlier beginning of flowering in comparison with Lirajet, 22 hybrids with significant earlier ending of flowering, three hybrids with significant higher 1000-seed weight, two hybrids with significant shorter plants than tester, 13 hybrids with significant higher oil content than tester Lirajet. The best hybrids out yielded about 40% higher than tester Lirajet.

Nevertheless the average effect of heterosis with respect to the seed yield was 16% in comparison with the tester Lirajet. Moreover, Spearman coefficients of correlation between estimated traits were calculated. Positive significant correlations at $P \leq 0.01$ Spearman coefficient of correlation $r_s = 0.48^{**}$ was calculated between winter hardiness and yielding. Moreover, negative Spearman coefficients of correlation between winter hardiness as well as beginning and ending of flowering was noted.

Satwinder *et al.* (1997) evaluated diallel crosses involving eight varieties of *Brassica napus* for seed oil yield and seven related components and they found high variation for SCA and GCA for all traits, suggesting both additive and non-additive gene effects. They also found combinations of varieties with high \times low or high \times average oil contents had high SCA effects.

Pietka *et al.* (1998) reported that winter hardiness of winter oilseed rape cultivars became very important trait after two strong winters which destroyed many plantations of this crop in Poland. These two winters gave rape breeders an opportunity to estimate winter hardiness of breeding materials and to make effective selections. A field trial with an F_2 generation of a diallel cross (7 \times 7) and with an F_1 generation of diallel cross (10 \times 10) were sown in autumn 1996. Winter losses of plants on the plots differentiated the hybrids significantly, allowing more sophisticated analysis. Seeds used for sowing the first trial were harvested from F_1 plants which survived the severe 1995-96 winter. The second trial was sown with seeds obtained by hand pollination after removing the anthers. The trials were made in a complete randomized block design with standard plots distributed systematically. Interblock variability was reduced with covariance analysis. The hybrids of both generations were examined in trials without parents. The number of plants which survived the winter were estimated in spring. Diallel analysis on transformed values was done according to Griffing method III. Effects of general (GCA) and specific combining abilities (SCA) and effects of reciprocal (RE) crosses were calculated. All effects except of reciprocal effects in F_1 generation are highly significant. Winter hardiness was shown to be a complicated character whose genetic control depends on additive effects of parent, interaction of parental genotypes and maternal cytoplasm.

Pu (1998) stated that a cytoplasmically male sterile line Ning A3 (MICMS), a *Brassica napus* line with a high level of sinaptic acid, was used as the basic breeding stock. The maintainer line Ning B3 was crossed with an elite cultivar with double low and fertile cytoplasm. Ning A6 and the maintainer line Ning B6 were bred after six generations of breeding. The combining ability of Ning A6 is high and the hybrids showed obvious heterotic vigour. Some hybrid combinations gave good performance in both yield and low content of sinaptic acid. The content of sinaptic acid in Ning A6 is 0.38% $\mu\text{mol per g DW}$.

Wos *et al.* (1997) studied in the combining ability of 55 inbred lines of rape (*Brassica napus*) and heterosis effects of their 62 F_1 hybrids. GCA was significant for seed yield, 1000-seed weight, time to flowering and fat content. They found that some 24 hybrids had higher yields, 14 earlier onset of flowering, three shorter plants, 14 higher fat content and three had higher protein content than control Global. Average yield increase over Global was 10%. There was a significant positive correlation of seed protein content with 1000-seed weight, and a negative correlation with seed fat content.

Kudla (1997) stated that inbred lines T1170, T1162, T1148 and T1166 were crossed in a factorial design with cultivars Maxol, Mandarin and Silex. Parental forms and 12 F_1 hybrids were evaluated in 1994-95 in a field trial. GCA of inbred lines and cultivars was significant for height to first branch, number of primary branches, siliqua length, seeds/siliqua and 1000-seed weight. T1170 and T1166 transferred some high-yield traits to their progeny. Significant differentiation of SCA was found for height to first branch. Dominance effects appeared high and positive for seed yield/plant and plant height. Additive gene action played a predominant role in the inheritance of height to first branch and seeds/siliqua. Relation of additive and non-additive gene action was generally similar in the inheritance of number of primary branches, siliqua length and 1000-seed weight. F_1 hybrids showed positive heterosis, averaging 14% for seed yield/plant.

Thakur *et al.* (1997) found that GSL8809, HPNI, GSL1501 and HNS8803 were good combiners for seed yield and some of its components and for oil content. They evaluated nine diverse inbreeds and their 36 F_1 hybrids from a diallel cross for yield and its components and for oil content. Mean squares due to general and specific combining ability were significant for all the

traits studied, suggesting the importance of both additive and dominance components of variation.

In a study of 8×8 diallel analysis (excluding reciprocals) Yadav *et al.* (1996) reported that the presence of both additive and dominance genetic components for seed yield and yield components in Toria (*Brassica campestris* L. var. Toria). But the magnitude of dominance component was larger than the additive component for all the traits including seed yield. Heritability estimates were higher for days to maturity and 1000 seed weight.

Krzymani *et al.* (1994) compared F_1 and F_2 generations from a diallel set of crosses between ten best strains. SCA for seed yield was significant in the first generation, but not in the second.

Kudla (1996) investigated the combining ability of winter oilseed rape (*Brassica napus*) inbred lines, and heterosis effects of F_1 and F_2 hybrids in the growing season of 1994-95. Analysis of variance showed that non-additive gene action had an advantage over additive gene action in the inheritance of plant height and number of primary branches. The significant effects of dominance genes in the F_1 for siliqua length, seeds/siliqua, seed yield/plant and 1000-seed weight did not occur in the F_2 . The differentiation of GCA of inbred lines, based on F_1 hybrids, was significant for siliqua length, seeds/siliqua, seed yield/plant and 1000-seed weight. GCA based on the F_2 was significant for pod length and seeds/siliqua. Inbred lines T1056 and T1150 were good components for crossing to increase seed yield in the F_1 . Both lines can be used for breeding high yielding oilseed rape hybrids varieties. In most of the F_1 and F_2 hybrids, significant positive effects of heterosis were found for plant height. F_1 of T1056 x Wotan showed the highest and significant heterotic effect (24.5%) for seed yield/plant. The mean heterotic effect in F_1 hybrids was 10% for seed yield, decreasing to 2% in the F_2 generation.

Patel *et al.* (1996) provided information that combining ability was derived from data on nine yield components in four parental genotypes (*Brassica juncea* cultivars Pusa Bold and TM17, *B. carinata* and *B. napus*) and their 12 F_1 hybrids grown during 1994-95. Variance due to GCA and SCA were significant for all the characters, except number of seeds/silique for GCA variance and 1000-seed weight for SCA variance. Non-additive gene action appeared to predominate for

all characters except days to maturity, which was governed by additive gene action. *B. carinata* was the best general combiner for plant height, number of branches/plant, number of siliquae/plant and oil percentage. Among the hybrids, *B. napus* x Pusa Bold was the best specific combination, followed by the reciprocal.

Krzymanski *et al.* (1995) evaluated seed glucosinolate content in hybrids from a diallel set of crosses involving ten *Brassica napus* strains. Only three of the strains showed significant GCA effects for total content of aliphatic glucosinolates but their values were low. SCA effects for the trait were significant only for three of the 45 crosses and heterosis only for two, but their values were high. Most strains appeared to have the same alleles that controlled low glucosinolate content. Heterosis for content of glucosinolates was not correlated with heterosis for seed yield.

Barua and Hazarika (1993) conducted a study during 1993 with five varieties representing two *Brassica napus* types and *Brassica campestris* var toria along with their hybrids from a half diallel set of crosses. According to them, heterosis mainly due to non-additive gene effect was important for dry matter and seed yield/plant. The important heterotic crosses were BSH1 × M27, B9 × PT303 and PK × M27.

Habetinek (1993) worked on *Brassica napus* and found higher GCA effects than SCA effects for all characters except seed weight/ plant. Darmor had the highest GCA for number of seeds/siliqua, siliqua length and 1000 seed weight, while Sonata had the highest GCA for oil content. SCA for seed weight/plant was highest in Sonata × SL2502.

Krzymanski (1993) studied yield and oil quality in ten parental and their 45 hybrids. Significant GCA and SCA effects were found for all 19 traits.

Kudla (1993) studied nine maternal lines (5S3 and 4S4), their pollinator (tester) Toplider and 9 F₁ hybrids derived by top crossing. Additive gene effects were most important in control of 1000-seed weight and the number of seed/siliqua, but non-additive effects predominated in control of number of primary branches, seed yield/plant, plant height and siliqua length. Differences in GCA between parents were significant for all characters except siliqua length. The inbred lines

T1057 and T6237 transmitted to the progeny high yield potential and T1057 had a good effect also on 1000 seed weight in the hybrids, but reduced seed/siliqua (which was increased by T6237). Favorable GCA effects were shown by T1080, T1097 and T1039 for seed/siliqua, T1097 for number of primary branches and T996 and T1039 for plant height.

Pszczola (1993) inter crossed the varieties Bolko, Tor, Diadem, Arabeke, Panter and Libravo in one set of diallel crosses and the varieties BOH 1491 (Bor), Falcon, Tapidor, Ofello and Lircus in another set. The characters evaluated were seed yield, 1000 seed weight, and others of importance. There was significant SCA effect in some crosses for all traits. Maternal (cytoplasmic) effect was apparent for all characters.

Rawat (1992) studied the reciprocal differences in the inheritance of eight yield traits in progeny from a diallel set of cross involving 12 lines of *Brassica juncea*. GCA effects predominated in the control of all the traits. Reciprocal effects were more pronounced than SCA effects, though the later were significant for all traits. The most promising parent lines on the basis of *per se* performance and of combining ability and F_1 performance were BICI624, BICI3S2, BICI439, BICI114 and BICI702. There was only one cross (BICI382 \times BICI702) in which reciprocal effects acted in a favorable direction for all traits. This allowed the selection of a maternal parent, which was capable of enhancing beneficial non-additive effects in a specific cross. The parents of this cross also showed high GCA for most of the traits, allowing the exploitation also of beneficial additive effects.

Singh *et al.* (1992) determined combining ability from data on 12 quantitative characters in the parents and F_1 hybrids from a 10 line \times 4 tester cross of Ethiopian mustard. Several of the lines were identified as being good general combiners. These are HC1, BC2 and BCIDI for maturity traits. FC5 for seed attributes and CAJR4-3, BCIDI, CAR3 and CARS for seed yield and several other desirable traits. The best specific combinations for yield improvement were CAR3 \times BC2 and BCIDI \times BC2 for using a pedigree selection programme.

Yadav *et al.* (1992) evaluated 45 F₁ hybrids of Indian mustard together with ten parents for combining ability with respect to seed yield and its component characters. Veruna, Kranti, RIC1359 and RLCI357 were identified as good combiners for seed yield, earliness, siliqua length, number of seeds/siliqua and 1000 seed weight. The following varieties or parents ECI26743, ECI26745 and ECI26746-1 have emerged as good combiners for plant height, primary branch and secondary branch.

Tamber *et al.* (1991) crossed 23 morphologically diverse *Brassica juncea* lines with four broad-based testers in 1987-88. The resulting 92 F₁ and parents and F₂ and parents were sown in 1988-89 and 1989-90, respectively. Data were recorded on number of days to first flowering and maturity. Analysis of variance of combining ability in both generations revealed that GCA variance due to lines and testers were significant for all characters except for maturity in the F₁ and additive effects in the F₂ were greater than in the F₁. Among the lines, RSK11 was the best general combining parent and was seen to be a suitable parent for evolving lines having short period of maturity. Among the testers, Varuna was a good general combiner in the F₂ generation and an average general combiner in the F₁ generation.

In tests of up to 210 *Brassica juncea* germplasm lines by Chauhan *et al.* (1990), there was wide variation in yield and its component. When 36 *Brassica juncea* crosses and their 15 parents were tested, there was significant difference in seed yield between genotype. NDRS602, Krishna, Pusa Bold and TM9 showed good general combining ability.

Siddique *et al.* (1990) studied a complete diallel cross involving four genotypes of *Brassica campestris* and their F₁'s for nine characters including seed yield/plant. Both additive and non additive gene action was found in the inheritance of characters except days to flower, plant height and primary branches. Preponderance of additive gene action for days to maturity, number of secondary branches/plant, number of siliqua/plant, number seeds/siliqua and non additive gene action for days to flowering, plant height, number of primary branches, siliqua length were found. Among the parents M-27 was the best general combiner for siliqua/plant and seed yield/plant. The hybrids YS-52 × M-27 exhibited highest significant SCA effect for seed yield/plant.

Arya *et al.* (1989) worked on combining ability from data of 12 yield related component characters in parents and F₁ of a 13 line × 3 tester mating design of *Brassica napus*. The varieties Midas, Regent 3-1 and DB054 were identified as good general combiners and DNA 38 × DISNI and N20-1 × Regent as good specific cross combinations.

Singh *et al.* (1989) worked with six *Brassica juncea* parents and their resultant 15 F₁ and 15 F₂ populations. They evaluated 11 quantitative and qualitative characters. GCA and SCA variance were significant for all characters. RLM198 showed good general combining ability for plant height, number of siliqua/plant, and yield. The parents, I RNS12 showed good general combining ability for no. of seeds/siliqua and seed weight. The cross RLM198 × R75-1 showed significant SCA for seed yield in both F₁ and F₂.

Information on combining ability derived from data on seven characters in 23 lines of *Brassica juncea* and their F₁ and F₂ hybrids by Wani and Srivasiava (1989) indicated that parents RK8202, KR5610, RK1418, RH30, V10 and B3U were good general combiners for seed yield.

In another study Thakur *et al.* (1989) studied yield components in 15 *Brassica juncea* lines and three testers and their F₁ hybrids. The lines Gonda-3 and R71-2 have had high GCA for yield.

Varma *et al.* (1989) studied seven yellow sarson (*Brassica campestris*) lines and their hybrids for eleven yield component characters YST151 and PYS6 had high GCA for all characters except 1000 seed weight.

Chawdhury *et al.* (1988) investigated thirteen selected *Brassica juncea* genotypes and their 78 hybrids from a half diallel cross. Data were tabulated on genetic variance and combining ability. RH30, RH785 and Varuna showed good performance and GCA for yield/plant, and its component. KC781 × RH30 and RH7513 × Varuna were the hybrids with best SCA effects and mean performance for yield and its components.

Badwal and Labana (1987) analysed data on seed yield/plant and eight related traits from a 10 × 10 half diallel cross in *Brassica juncea*. They reported that both additive and non-additive components of variance controlled the inheritance of seed yield, number of seeds/silique, plant height, primary branches, silique length; only non-additive variance was significant for secondary branches.

Chaudhury *et al.* (1987) found significant differences for GCA and SCA variances indicating that both additive and non-additive components of gene effects influenced the expression of each character in a trial of *Brassica chinensis* and four genotypes of *Brassica campestris* with their ten possible combinations (excluding reciprocals). The dominance component was greater than the additive component for all characters except seed size and silique length. The best general combiners for yield and its component were BSHI and Pusa Kalyani. The hybrids with the highest *per se* performance and SCA effects were *Brassica chinensis* × Pusa Kalyani and *Brassica chinensis* × Span. The best overall cross for the characters studied was Bell × Pusakalyani.

Chauhan (1987) tabulated genetic variance parameters for yield/plant and eight related traits from a 20 partial diallel cross in *Brassica juncea*. Variance due to GCA and SCA effects were highly significant for all traits. Additive genetic effects appeared predominant for three characters and non-additive effects for the remainder, Varuna, RS3 and Cult47 were good general combiners for yield as was RB85 for days to flowering and maturity.

Gupta *et al.* (1987a) worked with 8 × 8 diallel cross without reciprocals of *Brassica* genotype. GCA and SCA mean squares were significant for all characters studied. Non-additive gene effects appeared to be predominant for number of primary and secondary branches, silique length, number of seed/silique and seed yield, while additive-gene effects were apparently predominant for plant height. The best general combiner for seed yield was RLM198. The best crosses for further selection were RLM822 × Varuna and RLM198 × RH30.

Gupta *et al.* (1987b) performed an analysis in a 13×4 line \times tester cross in *Brassica juncea*. Additive gene effects were relatively more important than non-additive for seed yield/plant and most of the five yield component investigated. Among females, the best general combiners were RLM29 for seed yield, P Rai-1 for plant height, RLM240 for no. of primary and secondary branches. Among males, RLM198 was the best general combiners for seed yield, number of primary branches. Varuna was best for plant height and RL18 for number of secondary branches. The cross PI 1/17 \times RH-30 exhibited high performance for seed yield along with significant SCA for number of primary and secondary branches, RLM24 \times RH30 and RLM82 \times Varuna showed desirable significant SCA effect for seed yield and plant height.

Prakash *et al.* (1987) analyzed data of the F_2 of an eight parent diallel cross and showed that GCA and SCA variances were significant for yield components. SCA variance were higher than GCA variance for number of seeds/siliqua, 1000 seed weight, and seed yield indicating that dominance was possibly the predominant gene action for these traits. The parents DIR146 and RCL1017 were good general combiners for most of the characters studied.

Rawat (1987) observed a line \times tester analysis involving 12 females and five males of *Brassica juncea* of diverse origin. Variance components of GCA and SCA were significant for days of 50% flowering, number of primary branch, plant height, seed weight and seed yield/plant. For secondary branches GCA was important. Pusa Rai 34 and Pusa Rai 45 among the female parents and Pusa Rai 30 among the male parents performed well and were good general combiners. The cross RLM514 \times RLM198, RW336 \times Pusa Rai 30, Pusa Rai 45 \times BR40 and RH7710 \times Pusa Rai30 showed significant SCA for increased seed yield.

Singh and Chauhan (1987) worked with 60 triple test cross families produced by the crossing of $20F_2$ parents as males to the parents and F_1 s. In Varuna \times TM9 additive genetic variance appeared to be predominant for days to maturity, number of primary branch while dominance seemed to be mainly involved in the control of seed yield/plant. In Varuna \times RW75-80-1, additive genetic variance was estimated to be predominant for plant height and dominant for days to maturity, number of seeds/siliqua, 1000 seed weight, yeild/plant.

Singh *et al.* (1987) reported data on yield and eight other agronomic characters from an eight parent diallel cross in yellow sarson to indicate the presence of both additive and non-additive gene action, in the inheritance of all traits, with non-additive gene action being predominant for all traits, except plant height. YSK4 and YSK5 were good general combiners for seed yield/plant while the best combinations were YSK5 × YST151 and K88 × YSK5.

Griffing (1956) proposed a more general procedure for diallel analysis which makes provision for non-allelic interaction. In this approach mean measurement of a cross is partitioned into two major components, a part from a general mean (μ) and an environmental component, (i) the contribution of the parents, the general combining ability (GCA) effect analogous to main effect of a factorial designs, and (ii) the excess over and above the sum of the two GCA effects called the specific combining ability (SCA) effect, analogous to an interaction effect of a factorial design. The diallel approach has been extensively used, in cross pollinated crops. Griffing (1958) emphasized the statistical concepts of general and specific combining ability. Variance for general combining ability involves mostly additive gene effects which variance for specific combining ability depends on dominance.

2.2 HETEROSIS

The term hetrosis refers to the phenomenon in which F_1 population generated by crossing of two dissimilar parents showed increased or decreased vigor over the mid parental values or the better parental values. Both intra and inter-specific crosses showed some heterotic effect and both positive and negative heterosis were found.

Aderfis and Heiko (2005) revealed that heterosis is commercially exploited in rapeseed (*Brassica napus* L.) and its potential use has been demonstrated in turnip rape (*B. rapa* L.) and Indian mustard (*B. juncea* L.). In Ethiopian mustard (*B. carinata* A. Braun), however, information regarding heterosis has not been previously reported. This study, therefore, was conducted to generate information on heterosis and combining ability in *B. carinata*. Nine inbred parents and their 36 F_1 s, obtained by half-diallel cross, were evaluated for 12 traits at three locations in Ethiopia. Analysis of variance showed the presence of significant heterosis for all the traits. Seed yield showed the highest relative mid-parent heterosis that varied from 25 to 145% with a mean

of 67%. Relative high-parent heterosis for seed yield varied from 16 to 124% with a mean of 53%. General combining ability (GCA) effects were predominant in all traits except secondary branches and pods per plant. Specific combining ability (SCA) was significant for days to flowering, secondary branches, pods per plant, pod length, seeds per pod, 1000-seed weight and oil content. Interaction effects of GCA x location were significant for all traits except days to flowering, days to maturity, and oil content. All traits had significant SCA x location interaction effects. GCA effect for seed yield was positively correlated with F_1 performance ($r = 0.77$) and absolute mid-parent heterosis ($r = 0.67$). The presence of high levels of mid- and high-parent heterosis indicates a considerable potential to embark on breeding of hybrid or synthetic cultivars in Ethiopian mustard.

Huq (2007) conducted an experiment on *Brassica rapa* involving 7×7 half diallel cross. Heterosis and combining ability were estimated for seed yield and other related characters such as days to flowering, days to maturity, plant height, number of primary and secondary branches, length of siliquae, seeds per siliqua, seed yield per plant, thousand seed weight. Out of twenty one crosses Agroni x BARIsar-6, Agroni x Tori-7, Shafal x BARIsar-6 and Agroni x Tori-7 showed significant heterosis over mid and better parent. Agroni x Tori-7 best for number of primary branches/plant and siliquae/plant.

Iftikhar *et al.* (2000) studied rape variety Tower and three stable M9 mutants for heterosis of yield components of inter-mutant crosses during 1997-99. F_1 generations expressed significant heterosis for number of primary branches, number and length of primary roots and siliquae, seeds/siliqua, yield/plant and oil content. It is concluded that these mutants are a good source of variation for future breeding programmes.

Qian *et al.* (2005) reported the observation on the inter subgenomic heterosis for seed yield among hybrids between natural *Brassica napus* (AnAnCnCn) and a new type of *B. napus* with introgressions of genomic components of *Brassica rapa* (ArAr). This *B. napus* was selected from the progeny of *B. napus* x *B. rapa* and (*B. napus* x *B. rapa*) x *B. rapa* based on extensive phenotypic and cytological observation. Among the 129 studied partial intersubgenomic hybrids, which were obtained by randomly crossing 13 lines of the new type of *B. napus* to 27 cultivars of *B. napus* from different regions as tester lines, about 90% of combinations exceeded the yield of

their respective tester lines, whereas about 75% and 25% of combinations surpassed two elite Chinese cultivars, respectively. This strong heterosis was further confirmed by reevaluating two out of the 129 combinations in a successive year and by surveying hybrids between 20 lines of the new type of *B. napus* and its parental *B. napus* in two locations. Some DNA segments from *B. rapa* were identified with significant effects on seed yield and yield components of the new type of *B. napus* and intersubgenomic hybrids in positive or negative direction. It seems that the genomic components introgressed from *B. rapa* contributed to improvement of seed yield of rapeseed.

Heterosis over the mid parent, better parent and commercial, check variety pusa bold was estimated for plant height, days to maturity, number of branches per plant, number of siliquae per plant, seed yield per plant (gm) and 1000 seed weight (g) in 17 crosses of *B. juncea* by Patil *et al.* (2005). The crosses ACN-9 × MCN-126 and ACN-9 × MCN-128 were the best performers for seed yield and number of siliquae/ plant. The maximum magnitude of significant positive heterosis for all the three types were also exhibited by these crosses and hence can be exploited for further utilization in a breeding programme.

Shen *et al.* (2005) observed significant differences in seed yield per plant and seed oil content among the F₁ hybrids and between F₁ progenies and their parents of *Brassica campestris*. However, the heterosis for seed yield per plant was much greater than that for seed oil content. Mid parent heterosis and high parent heterosis of seed yield per plant ranged from 5.50 to 64.11% and from -2.81 to 46.02%, while those of seed oil content ranged from -1.55 to 7.44% and -3.61 to 6.55%, respectively.

Katiyar *et al.* (2004) crosses out a study on heterosis for the seed yield in ninety intervarietal crosses of *Brassica campestris*. Twenty one crosses (23.3%) showed significant positive heterosis over better parent while only four crosses (4.4%) were over the best commercial variety (MYSL -203). The crosses, YST -151 × Pusa gold (dwarf), and MYSL -203 × EC -333596 showed highest heterosis up to 150.33 and 43.38 percents over best parent and commercial variety respectively. Line GYSG -1 (female parent) and Pusa gold (dwarf) were the most potential ones for giving largest proportions of crosses with high degree of heterosis.

Wang *et al.* (1999) analysed heterosis and combining abilities of 20 reciprocal cross combinations of five double low rape (*Brassica napus*) cultivars (lines) showing high seed yield. Positive mean heterosis varied among crosses. The positive mean heterosis of siliqua number/plant was 17.6% was highest, followed by seed number/siliqua and 1000-seed weight. Heterosis of F_1 generations were greatest when Zhihu 1 and Zhongyou 220 were used as parents. Liersch *et al.* (1999) conducted a breeding approach known as CMS ogura system of oilseed rape hybrid cultivars in Poland to evaluate yield and yield component variability of F_1 hybrids and their parental lines also heterosis effect, and qualitative traits such as oil and glucosinolate content in seeds. They found that composite hybrid cultivars yielded higher than restored hybrids. They stated that the yield of hybrids and qualitative traits such as oil and glucosinolate content in seeds are significantly dependent on genotypes and environmental conditions.

Ramsay *et al.* (1994) stated a complete diallel set of crosses, including selfs, was produced from eleven inbred lines of swedes and assessed in the field for both components of dry matter yield and neck length at Dundee, UK, during 1987. They found that there was a strong positive heterosis for dry matter yield with high yielding F_1 s showing an improvement of more than 20% above the better parent. Reciprocal differences were also found. Both additive and non-additive genetic variation was found for dry matter yield and other quantitative traits. However a simple additive-dominance model with independence of action and distribution of the genes failed to describe the data adequately. Given the implications for the breeding of inbred or F_1 hybrid swede cultivars, further experiments, using triple test crosses are suggested.

Satyndra *et al.* (2004) evaluated twenty one Indian mustard hybrids and their parents for eight quantitative traits: days to flowering, days to maturity, plant height, number of primary branches, length of the main raceme, seed yield, thousand seed weight and oil content percentage, in an experiment. High heterosis (15.99, 15.51 and 12.37%) was obtained for seed yield in the crosses Basanti \times NDR 8501, Basanti \times Kanti and Basanti \times RH 30, respectively. These hybrids showed high heterosis over the best cultivar. Among the crosses, Basanti \times Kranti may be used for selecting for seed yield and quality traits.

Yadav *et al.* (2004) had undertaken an investigation to estimate heterosis for seed yield and its components in Indian mustard. Hybrids Siifolia × NDRE-4 (-18.5%) and Trachystoma × NRCM-40 (-6.1%) exhibited the highest heterosis for days to flower initiation and days to maturity over better parent, respectively. The magnitude of heterosis was highest for plant height in Trachystoma × SK 93-1 (27.7%) over BP and (25.8%) over SV both. For the number of primary branches per plant Trachystoma × PR 905 showed 106.5 and 100.0% heterosis over BP and SV, respectively. Trachystoma × PHR -1 (125.1%) showed maximum heterosis over BP and Moricandia × NRCM -79 (9.6%) over SV for the number of secondary branches per plant. Siifolia × SM -1 showed 54.1% heterosis over BP and negative heterosis (-9.2%) over SV for seeds per siliqua. The highest heterosis for thousand seed weight was observed in Moricandia × PHR -1 (48.80%), followed by Trachystoma × NRCM 69 (20.6%) over BP and SV, respectively. Significant and positive magnitude of heterosis for oil content was observed in Trachystoma × NDYR -8 (10.1%) over BP and Siifolia × NRCM 79 (8.5%) over SV, respectively. The cross, Moricandia × NRCM 86 exhibited significant and positive heterosis over BP (82.8%) for seed yield per plant, followed by Siifolia × NRCM 86 (76.0%) and Moricandia × NRCM 98 (52.5%).

Goswami *et al.* (2004) conducted an experiment and estimated heterosis for yield and yield components in 30 crosses of Indian mustard. Results showed that the cross RH9404 × RH30 had the maximum heterosis for seed yield per plant (92.88 and 106.23%) during E₁ and E₂ respectively. This cross also showed high heterosis for thousand seed weight. The crosses RH9617 × RWHI and RH9621 × RWHI were selected because of high heterosis for all the parameters tested.

Mahak and lallu (2004) performed an experiment on Indian mustard strains/cultivars Varuna, Shekhar, Vardan, Laha 101, Pusa Bold, RH -30, Pusa Basant, NDR -8501 and Kranti were crossed in a diallel mating design excluding reciprocals. The parents along with 36 F₁s and 36 F₂s were grown data recorded for plant height, branches per plant, siliquae on main raceme, seed yield per plant, thousand seed weight, seed oil content, de-fatted seed content and protein content. The crosses exhibited highly significant heterosis for most of the characters studied.

Mahak *et al.* (2003a) studied heterosis for days to flowering, plant height, number of primary and secondary branches, length of main raceme, days to maturity, thousand seed weight, harvest index, oil content, protein content, and seed yield in 10 Indian mustard cultivars and 45 F₁ and F₂ hybrids. High heterosis for seed yield was observed in Varuna × Rohini (56.74%), Vardan × Rohini (53.43%) Varuna × RK 9501 (52.86%), Vardan × NDR 8501 (36.73%), pusa Bold × Rohini (37.68%), and Varuna × NDA8501 (32.54%).

Qi *et al.* (2003) carried an experiment out in 1997, 66 crosses were made in a diallel design of twelve parental varieties of *Brassica napus* to study heterosis of seed and its components. Twenty-one crosses showed a significant heterosis in seed yield/ plant. The average yield heterosis over their parents was 70.24% (30.70-218.10%). Eight crosses showed better parent heterosis (3.57-20.48%) in 1000-seed weights, while the parent of seven crosses showed low 1000-seed weights. Forty-seven crosses gave on average 28.02% (0.93-97.87%) more siliquae / plant in parents, while thirteen crosses showed 11.67% more seeds/ siliqua in parents. By this experiment they concluded that there was large potential heterosis in seed yield with heterosis of siliquae number/plant making the biggest contribution.

Ghosh *et al.* (2002) carried out a line × tester analysis involving 29 promising female and seven male parents for 10 quantitative traits in Indian mustard. The crosses YSRL-10 × Pusa bold, DBS-10 × Pusa bold showed high heterosis for seed yield and some of the yield contributing traits.

Kumar *et al.* (2002) crossed three lines and twelve testers of Indian mustard and the resulting 36 F₁'s and 15 parents were grown. Physiological data were determined from five plants per entry and the range of heterosis given for all crosses. The five hybrids with the highest heterosis for seed yield were RN-505 × RN-490, RN-505 × PCR-43, RN-393 × RN-481, RN-393 × RN-453 and RN-505 × RN-481, and these crosses offer the best possibilities of further exploitation for the development of high yielding varieties.

Pankaj *et al.* (2002) studied heterosis of parents for seed yield, oil content and protein content in an 8×8 diallel cross in toria (*brassica campestris* var. toria). Trait data were recorded on five plants of each of the 28 F_1 's and 28 reciprocal F_1 's (RF_1 's). 24 F_1 's and 21 RF_1 's showed significant positive heterosis for seed yield over mid parent (MP) and 16 F_1 's and 21 RF_1 's over the better parent (BP).

Zhang *et al.* (2000) crossed three double low cytoplasmically male sterile (CMS) and five double low restorer lines of *Brassica napus* and they analyzed resulting 15 hybrids for eight yield components. In this experiment they found that the CMS F_1 had significant heterosis, particularly for yield, but that predicted for the F_2 was lower. They also suggested that the major yield components, total siliquae number/plant had the highest heterosis and would be of more value in a breeding programme than trying to increase seed number per siliqua or 1000-seed weight.

Lu *et al.* (2001) proposed that heterosis is proportional to genetic divergence between respective parents in many crops. They evaluated heterosis in interspecific hybrids between *Brassica napus* (AACC, $2n=38$) and *Brassica rapa* (*B. campestris*) (AA, $2n=20$) for ten agronomic characteristics and compared to heterosis in hybrids of *B. napus*. They characterized fifteen inter-specific crosses for their cross ability, germination rate, morphology, pollen fertility, and seed production. They found cross ability ranged from 0.8 to 16.7 seeds per flower pollinated, with 7.5 seeds on average; germination of the F_1 seeds varied with combinations from 20.7 to 89.8%; highly significant high-parent heterosis in the number of secondary branches and siliquae number per plant and significant mid-parent heterosis in plant height, length of main inflorescence, and the number of primary branches. They also found that seed number per siliqua in inter-specific hybrid was significantly lower than both parents' and varied with different combinations and inter-specific hybrids showed higher vegetative heterosis than intra-specific hybrids.

Swarnkar *et al.* (2001) carried out heterosis analysis using 36 F_1 hybrids, 36 F_2 generations and parents obtained from 9×9 diallel mating design for 11 quantitative traits, viz. days to flowering, plants height (cm), number of primary branches, number of secondary branches, length of main raceme (cm), number of siliquae on main raceme, days to maturity, yield per

plant (g), thousand seed weight (g), oil content (%) and protein content (%). High economic heterosis for seed yield was observed to be present in four crosses, KR-5610 × PR-15 (58.38%), YRT-3 × PR-15 (54.33%), RK-1467 × T-6342 (52.60%) and KR-5610 × KRV –Tall (36.70%). The hybrids showing high heterosis over best cultivar can be successfully grown up to 2 or 3 early generations, which may prove beneficial for the Indian mustard growers.

Tyagi *et al.* (2001) evaluated forty-five hybrids of Indian mustard obtained from crossing ten cultivars for seed yield and yield components. The relative heterosis was desirable for plant height, number of primary and secondary branches per plant, seeds per siliqua, number of siliquae on main shoots, biological and seed yield, and oil content. Heterobeltiosis was desirable for primary and secondary branches per plant; siliquae on main shoots, and biological and seed yields. Standard heterosis was desirable for the number of primary and secondary branches per plant, siliqua length, and seeds per siliqua, number of siliquae on main shoots, biological and seed yields and oil content. The mean level of heterosis was highest for biological yield. The highest standard heterosis (206.14%) and heterobeltiosis (240.56%) for seed yield per plant was recorded in the cross BIO 772 × Rohini. This cross was the best heterotic combination for all the three types of heterosis for seed yield.

Wu *et al.* (2001) evaluated the heterosis of 80 hybrid combinations from TGMS line 402S and its original parent Xianyou 91S, and the combining ability of 40 test cross lines. The results of identification test showed that among 47 combinations yielding over the control Xianyou 15, seventeen ones with 402S and three ones with Xianyou 91S over yielded more than 20%, reaching the significant level of 1%; and among 51 combinations yielding over their corresponding higher yield parents, 18 ones with 402S and nine ones with Xianyou 91S over yielded at 5 or 1% significant level.

Tyagi *et al.* (2000) reported data on heterosis in intervarietal crosses in mustard (*Brassica juncea* (L.) Czern & cross.). Desirable significant and negative heterosis for plant height was observed in seven crosses, with Varuna × SKNM-90-14 exhibiting the most negative value (-14%). Maximum positive heterosis was recorded for seed yield per plant (-48.0 to 93.3%), with crosses PCR-7 × SKNM90-13, RH-30 × TM18-8 and PCR7 × JM90-12 giving values of 93.3, 81.3 and

77.3%, respectively. In general, positive heterosis for seed yield was accompanied by positive heterosis for siliqua length, seeds per siliqua, 1000-seed weight, biological yield and harvest index.

Katiyar *et al.* (2000a) information on heterosis and combining ability is derived from data on seed yield and three yield components in six lines, 16 testers and their 96 F₁ hybrids from a line × tester mating design. Of the hybrids, 64 and 38 showed heterosis for seed yield over the better parent and standard cv. varuna, respectively.

Qi *et al.* (2000) investigated heterosis in hybrids of six cultivars of *Brassica campestris*. They found that yields of hybrids ranged from 46 to 125kg. Significant heterosis for yield was found some hybrids with highest being 96.4%. Most hybrids showed lower levels of heterosis, with the lowest being 1.4%.

Agarwal and Badwal (1998) studied the extent of heterosis for yield and other characters in 19 F₁ hybrids of *Brassica juncea* and compared to five commercial cultivars. Eighteen hybrids out yielded the best control variety RLM514. Three of them (MS × Plant Rai 1002, MS × RH848 and MS × RLC1047) were superior over the best control in seed yield by 81.19, 50.65 and 64.94%, respectively. Overall heterosis (taking all hybrids and check into account) for seed yield was very high (59.69%). The agronomic superiority of the three hybrids was reflected by 1.5 to 2.0 fold increase in oil yield and one week earliness in flowering as compared to RLM514.

Yadav *et al.* (1998) studied some 27 crosses of female and three male sarson (*Brassica campestris*) parents for seven yield components. Of these, 18 hybrids exhibited significant positive heterosis. Highest heterotic response for seed was observed in DB₁ × Pusa kalyani and BSKI × BSI k₂.

Thakur *et al.* (1997) evaluated nine diverse inbreds and their 36 F₁ hybrids from a diallel cross for yield and its components and oil content. They observed that estimates of heterosis over better parent (BP) for the various traits were significant for seed yield (-14.8 to 82.8%), primary branches (-26.0 to 193.6%) and siliquae per plant (-21.9 to 162.6%). They also observed

unidirectional dominance for most of the traits studied and the cross GSB7027 × HNS8803 gave highest positive heterosis for seed yield per plant.

Varshney and Rao (1997) estimated combining ability, heterosis and inbreeding depression in yellow sarson (*Brassica campestris*) for eleven quantitative characters. The hybrids, which exhibited highest heterosis also showed higher inbreeding depression. Heterosis over better parent was highest for siliquae per plant (162.9%), followed by economic yield per plant (129.4%), Biological yield per plant (118.7%), primary branches per plant (118.7%) and secondary branches per plant (88.1%).

Yadav *et al.* (1997) studied heterosis in toria (*Brassica campestris* var. toria). He used 6 lines and their 15 F₁ hybrids and studied on eight yield components. The cross white flower × TC113 had the highest negative heterosis (being desirable) for plant height. The crosses White flower × TS61, TH68 × TC113, White flower × Sangam and White flower × TS61 were superior for seed yield.

Singh *et al.* (1996) studied heterosis for yield and oil content in *Brassica juncea* L. Heterosis over better parent was recorded in the crosses PR1108 × BJ-679 by 77.6% and BJ-1257 × Glossy mutant by 13.1% for seed yield and oil content, respectively. Oil content was positively associated with thousand seed weight and seed yield indicating the possibility of simultaneous improvement for these characters.

Ali *et al.* (1995) investigated the association between distance and mid-parent heterosis and they found that the correlation between genetic distance and heterosis was positive and highly significant for seed yield, siliquae/plant and seeds/siliqua. They estimated genetic distance among canola [rape] cultivars through multivariate analysis. They analysed thirty cultivars from various sources and clustered into three distinct clusters based upon five morphological characteristics and yield components (crown diameter, branches/plant, siliquae/plant, seeds/siliqua and yield/plant). Two cultivars from each cluster were selected as parents and 15 partial-diallel inter- and intracluster crosses were made between the six selected parents and evaluated at two locations in Michigan, USA in 1990-91.

Hari *et al.* (1995) conducted an experiment to derive information on heterosis from data on eight yield components in seven rape (*Brassica napus*) genotypes and their 21 F₁ hybrids grown during winter 1992 in Haryana. They found that hybrid HNS9002 × N20-7 had high positive heterosis for primary and secondary branches, siliques on main shoot and seeds per silique. They also found another hybrid, HNS9005 × N20-7, exhibited appreciable heterosis over the better parent (HNS9005) for seed yield and oil content. They also proposed that these hybrids were promising for exploitation of heterosis. They informed that parent N20-7 developed from Japanese material Norin 20 was a promising parent for exploitation in the hybrid breeding programme.

Information on heterosis has also been recorded by Rai and Singh (1994) from data on six yield components in eight *Brassica campestris* varieties and their 28 F₁ hybrids. A number of hybrids expressed heterosis for seed yield and its component. The average heterosis over better parent for seed yield was 21.3%. The crosses showed significantly high positive heterosis for seed yield in all cases except had high negative heterosis for yield in DTS × YST151.

Ahmad (1993) worked with parents and F₁ hybrids from crosses between resynthesized lines and improved 00 varieties. F₁ were earlier maturing than resynthesized lines and heterosis was observed for spring regrowth and plant height. In trials, the best resyn. line H128 could only produce 87% of the mean yield of the improved varieties.

Gupta *et al.* (1993) studied 56 hybrids from a half diallel set of crosses involving eight genetic stocks with 28 hybrids being derived from crosses of the initial S₀ population and the rest from crosses of S₁ families from each of the parents. The use of S₁ families generally gave hybrids with a higher degree of commercial heterosis (over the best open pollinated commercial variety) than hybrids using S₀ materials, though the S₀ × S₀ crosses gave high commercial heterosis for yield in many cases.

Gupta and Labana (1995) provided information on combining ability and heterosis for seed, straw and chaff protein contents and nitrogen and protein harvest indexes was derived from data on distribution of nitrogen in plant parts as assessed in 8 *Brassica napus* cultivars and their 28 F₁ hybrids grown at Ludhiana in 1985-86. Protein contents were estimated from nitrogen content values. Topa was the best combiner for seed protein content.

Yu and Tang (1995) studied on seven inbred rape lines and their 21 F₁ hybrids which were compared at the seedling stage for acid phosphatase (APS) isoenzyme patterns by polyacrylamide gel electrophoresis (PAGE) analysis. All hybrids with hybrid band(s) in their zymograms showed heterosis in yield, and those without hybrid bands showed no heterosis. Hybrids with two or three hybrid bands and high APS activity showed great heterosis. Hybrids with 2-3 medium or weak hybrid bands had only moderate heterosis. Hybrids derived from parents with very different zymograms showed high heterosis even though they had only one strong hybrid band. When the parents had similar zymograms and the hybrid showed relatively low APS activity, heterosis was low. Since the isoenzymes of APS in *Brassica napus* appeared to be quite stable, they were recommended to serve as a biochemical indicator of heterosis at the seedling stage (the 2-3 leaf stage).

The agronomic performance of inter-cultivar hybrids of *Brassica rapa* L. was studied in crosses between four *B. rapa* cultivars by Falk *et. al.* (1994). Six reciprocal hybrid combinations were produced by hand pollination in the greenhouse. Hybrids and their parents were tested in replicated yield tests for 3 years. An average, over all hybrid combinations' of 13% heterosis for seed yield was observed. Heterosis for seed yield was greatest in crosses between genetically diverse cultivars which agrees with classical theories on heterosis. There were also significant differences in yield of reciprocal F₁ combinations. Seed oil content was not heterotic. The results of this study indicated that the level of heterosis of seed yield in crosses between *B. rapa* cultivars adapted for production in western Canada is sufficient to warrant the development of a suitable pollination control system for hybrid *B. rapa* production.

Habetinek (1993) determined plant length, siliqua length, no. of seed/siliqua, 1000 seed weight in five varieties of the 00 types and their F₁ hybrids from a diallel set of crosses. The greatest heterosis over the better parent was for seed weight/plant. Sonata × SL502 had the highest heterosis value for seed weight/plant. Kudla (1993) also found high heterosis for seed yield/plant and was shown by all hybrids (10.2- 62% over the better parent) in a study of nine maternal lines (5S₃ and 4S₄) and their pollinator, taplidor and 9 F₁ hybrids derived by top crossing.

Krzymaniński (1993) found significant heterosis for seed yield, oil content and some flowering traits in ten parental strains and their 45 hybrids. The mean heterosis for seed yield over the mid parental mean was 24.71%. The highest heterosis for this trait was seen in the cross of PN2595/91 × PN2870/91 (71.81% relative to the mid parental mean).

Pradhan *et al.* (1993) found from the component character analysis concluded that characters such as no. of primary and secondary branches, number of siliquae/plant and siliqua density contributed significantly to positive heterosis for yield.

Srivastava and Rai (1993) tested heterosis for seed yield and three of its component in hybrids from a half diallel set of 15 crosses involving three Indian and three foreign varieties. The highly heterotic hybrids YST151 × Tobin, YST151 × Torch and PT303 × Torch, each had one Indian and one foreign parent and in general the Indian × foreign hybrids showed a higher degree of heterosis than the Indian × Indian and foreign × Foreign.

Krishnapal and Ghose (1992) investigated the relationship between heterosis and genetic diversity in the F₁ from crosses involving five genotype of rapeseed (*Brassica campestris*) and six mustard (*Brassica juncea*). Cross combinations in genotype having mediums *djk* values (ranging from 2.52 to 7.79) exhibited positive and significant heterosis for most characters in rapeseed but in mustard, heterosis for seed yield was positive and significant in all cross combination regardless of which genotype had high or low *djk* value. In mustard more heterosis for seed yield/plant and 1000 seed weight were observed. However, combination with a medium heterosis for seed yield and some of its component, high heterosis in cross combinations of genotypes of low *djk* value may result from cancellation of the mean of one character by that of

the other characters). Therefore, dissimilarity/ variation between genotypes is not always positively associated with heterosis.

Hirve and Tiwari (1991) evaluated 28 elite *Brassica juncea* genotypes produced 28 F₁ and F₂ progenies together with the parents, for siliquae and seed yield per plant and siliqua length. The highest heterosis for seed yield was obtained in the cross RAU × RPU 18 (161%). RLM 198 × Veruna, RAU RP₄ × Varuna and Tm 7 × Varuna also gave good seed yield heterosis and gave high heterosis for other yield contributing characters. In general, crosses containing Varuna as one parent gave high heterotic values.

Heterosis and epistasis in spring oil seed rape (*Brassica napus*) was analysed by Evggvist and Becker (1991) by comparing generation means for ten agronomic traits. Parents, F₂, F₂ and F₆ generations of four crosses with Swedish French material were investigated. The F₂ was 11% higher in yield, earlier in flowering time and slightly later in maturation when compared with their parents.

A male sterile line, European-Xinping A, a maintainer line European -Xinping B and a – restorer line 74243-6, were developed from a male sterile plant of *Brassica juncea* by Shi *et al.* (1991). The seedling stage of F₁ hybrids showed fairly strong heterosis; there was also heterosis in seed yield. The F₁ hybrids yielded 19.2-34.8% more than CV. Kunming –Gaoke.

Zheng and Fu (1991) worked with eight F₁ hybrids of *Brassica napus* L. They evaluated 17 agronomic traits with four heterosis standard. Of all the traits investigated, seed yield/plant and effective siliqua/plant showed significant heterosis, their mean heterosis (over mean value of the parents) rates being 80.21 and 51.47 percent, respectively.

Kumar *et al.* (1990) evaluated 16 parents and 39 F₁s for six traits. Crosses showing positive heterosis for seed yield also showed positive heterosis for primary branches, secondary branches, siliqua length and number of seeds/siliqua. Highest positive heterosis in secondary branches, siliqua length and number of seeds/siliqua. Highest positive heterosis for seed yield was observed in the cross RLM198 × RH30 and was followed by the crosses RJLMSH × Varuna;

RL18 × Varuna and RS64 × Varuna. RLM198 × RH30 also recorded highest heterobeltiosis for secondary branches.

In a similar experiment conducted by Nasim (1990) with six cultivars of *Brassica campestris* crossed in half diallel fashion M-91 × TS-72 showed highest heterosis over mid parent for seed yield/plant.

In a study of combining ability and heterosis in *Brassica campestris* Siddique *et al.* (1990) found up to 117.21% heterosis over mid parent for seed yield.

Badwal and Labana (1987) studied *Brassica juncea* for seed yield/plant and other eight related characters. In F₁, they found positive and significant heterosis for almost all traits. In a study for heterosis and cytoplasmic-genetic male sterility in oil seed rape (*Brassica napus* L.) through diallel cross of six Canadian and European cultivars.

Grants (1985) found heterosis for seed yield up to 72% over better parents.

Lefort *et al.* (1987a) while studying *Brassica napus* of Asian and European parental lines and their hybrids, reported that plant height and seed yield showed positive heterosis in the hybrids.

Banga and Labana (1984) reported several important findings on heterosis of Indian mustard (*Brassica juncea*). They studied 139 F₁ of two groups Indian and European lines. The greatest heterosis over better parent was estimated for seed yield/plant. High heterosis was also estimated for number of secondary branches.

Lefort (1982) studied 140 F₁ hybrids of winter oil seed rape (*Brassica napus* L.) and found that for seed yield average hybrids vigour was 23.5% on the basis of the mid parent. In a few cross combinations the value reached up to 50% in relation to the best parent value. This emphasizes the interest of hybrids varieties for improving yield.

Schuster *et al.* (1978) reported heterosis of 203% for seed yield, 211% for seed no./ siliqua and 187% of no. of siliqua/plant in crosses between diverse lines in each generation of black mustard (*Brassica nigra* L.). There was lower heterosis for 1000 seed weight.

Zuberi and Ahmed (1973) studies six crosses of four strains of *Brassica campestris* var Toria for yeild and its component characters. They estimated heterosis for different charaters. According to them heterosis for different characters varied widely due to cross combination.



Chapter 3

Materials and Methods

MATERIALS AND METHODS


To conduct the experiment five selected cultivars were used as parents and these were Tori 7, SAU sarisha 2, SAU sarisha 3, BARI sarisha 6 and BARI sarisha 15. Ten crosses were done among parents in Rabi season 2010-2011. In 2011-2012 Rabi season, the parents and F_1 's were grown in the experimental farm.

3.1 Experimental site and duration

The research work was conducted at the experimental farm of Department of Genetics and Plant Breeding, Sher-e-Bangla Agricultural University (SAU), Dhaka-1207, Bangladesh, during the period from October 2010 to February 2011 and October 2011 to February 2012. The soil of the experimental plots were clay loam, land was medium high with medium fertility level (Appendix I). The site was situated in the subtropical climatic zone, wet summer and dry winter is the general climatic feature of this region (Figure 1). During the rabi season the rainfall generally is scant and temperature moderate with short day length. Meteorological data on rainfall, temperature, relative humidity from October 2011 to February 2012 were obtained from the Department of Metrological Centre, Dhaka-1207, Bangladesh (Appendix II).

3.2 Plant Materials

Five parental genotypes and there ten intervarietal hybrids were used in the experiment. The present status, source of the materials and characteristics of the parents used in the intra-specific crosses and the attempted cross combinations are presented in Table 1. The parents were crossed in half diallel to produce 10 F_1 's during winter 2010 at the experimental farm of SAU, Dhaka, Bangladesh.



The experimental site under study

Figure 1: Location of the experimental field

Table 1: Cross combinations in half diallel system of five varieties in *Brassica napus* L.

Parents Parents	Tori 7	SAU Sarisha 2	SAU Sarisha 3	BARI Sarisha 6	BARI Sarisha 15
Tori 7	Tori 7	Tori 7 X SAU Sarisha 2	Tori 7 X SAU Sarisha 3	Tori 7 X BARI Sarisha 6	Tori 7 X BARI Sarisha 15
SAU Sarisha 2		SAU Sarisha 2	SAU Sarisha 2 X SAU Sarisha 3	SAU Sarisha 2 X BARI Sarisha 6	SAU Sarisha 2 X BARI Sarisha 15
SAU Sarisha 3			SAU Sarisha 3	SAU Sarisha 3 X BARI Sarisha 6	SAU Sarisha 3 X BARI Sarisha 15
BARI Sarisha 6				BARI Sarisha 6	BARI Sarisha 6 X BARI Sarisha 15
BARI Sarisha 15					BARI Sarisha 15

3.3 Land preparation and fertilizer application

The land was ploughed well by power tiller followed by laddering. The stubbles and weeds were removed carefully. Chemical fertilizers were applied at the rate of 220-140-80-150-5 kg/ha of Urea, Triple Super Phosphate (TSP), Muriate of Potash (MP), Gypsum and Zinc Sulphate respectively. Cowdung was applied at the rate of 5 ton/ha. The whole amount of TSP, MP, Gypsum, Zinc Sulphate and 50% Urea were applied as basal dose. The remaining 50% Urea was applied as top dressing at flower initiation stage.

3.4 Experimental design and layout

The seeds of ten F_1 's and five parents were grown in randomized complete block design (RCBD) with three replications. Each plot consisted of single row of 3 m length spaced 40 cm apart and 10 cm between plants. The seeds were sown in separate line in the experimental field on 3 October 2011 by hand uniformly. The seeds were sown at a soil depth of 2.5 to 3.5 cm. After sowing the seeds were covered with soil carefully so that no clods were on the seeds. Seed germination started after three to four days of sowing. Treatment was distributed in the experimental unit through randomization by using the random number.

3.5 Irrigation and drainage

One post sowing irrigation was given by sprinkler after sowing of seeds to bring proper moisture condition of soil to ensure uniform germination of the seeds. A good drainage system was maintained for immediate release of rain water from the experimental plot during the growing period.

3.6 Intercultural operation, insect and disease control

Necessary intercultural operations were done during the crop period to ensure normal growth and development of the plants. Thinning and first weeding were done after fifteen days of sowing. Top-dressing, weeding and necessary thinning were done after 25 days of sowing. No remarkable disease attack was observed.

3.7 Harvesting of sample plants

When 80% of the plants showed symptoms of maturity i.e. straw color of siliquae, leaves, stem and desirable seed color in the matured siliquae, the crop was assessed to attain maturity. At maturity, ten plants were selected at random from the row of each plot. The sample plants were harvested by uprooting and then they were tagged properly. Data were recorded from these ten plants.





Plate 1: Photograph showing a field view of experimental site at seedling stage at SAU farm



Plate 2: Photograph showing a field view of experimental site at SAU farm



Plate 3: Photograph showing a field view of experimental site at flowering stage at SAU farm



Plate 4: Photograph showing a field view of experimental site at maturity stage at SAU farm

3.8 Data recorded

3.8.1 Days to maturity: Number of days required from sowing to siliquae maturity of 80% plants of each row.

3.8.2 Plant height: During harvesting the plant height was measured in cm from the ground level of the plant to the top of the plant. Mean height of ten plants was recorded.

3.8.3 Number of primary branches per plant: Mean numbers of branches originated from the main stem from ten randomly selected plants from each F_1 's and parents at maturity.

3.8.4 Number of secondary branches per plant: Number of branches originated from the primary branch from ten randomly selected plants from each F_1 's and parents at maturity.

3.8.5 Number of siliquae per plant: Mean number of siliquae obtained from ten randomly selected plants from each F_1 's and parents at maturity

3.8.6 Length of siliqua: Five siliqua was selected at random from every selected plant to measure the length of siliqua. The measurement was in cm. Distance between the end of the peduncle to the starting point of the beak was considered as siliqua length.

3.8.7 Number of seeds per siliqua: All siliqua from the sample plants was collected and 05 siliqua was randomly selected. Seeds obtained from them, were counted and average numbers of seeds per siliquae was recorded.

3.8.8 Thousand-seed weight (g): Weight in grams of 1000-seed was recorded from ten randomly selected plants of each F_1 's and parents.

3.8.9 Seed yield per plant (g): Mean seed weight in grams of ten randomly selected plants from each F_1 's and parents after harvest.

3.9 Statistical analysis

Statistical analyses were done to calculate the analyses of variance and other parameters of the genotypes for the characters tested.

3.9.1 Combining ability in relation to diallel cross

Griffing (1956) proposed four methods of analysis depending on the materials involved. Griffing has also considered Eisenhart's model I (fixed effect) and model II (random effect) situation in the analysis. In the present research work combining ability analysis were done following method 2 (excluding reciprocals) and model-1.

The mathematical model for the analysis was:

$$Y_{ij} = m + g_i + g_j + S_{ij} + 1/bc \sum_{kl} e_{ijkl}$$

Where,

$i, j = 1, 2, \dots, p$

$K = 1, 2, \dots, b$

$L = 1, 2, \dots, c$

$P =$ Number of parents

$B =$ Number of blocks or replications

$c =$ Number of observation in each plot

$Y_{ij} =$ The mean of $i \times j^{\text{th}}$ genotype over K and L

$m =$ The population mean.

$g_i =$ The general combining ability (GCA) effect to i^{th} parent

$g_j =$ The GCA of j^{th} parent

$s_{ij} =$ The SCA effect such that $s_{ij} = s_{ji}$

$1/bc \sum_{kl} e_{ijkl} =$ The mean error effect

The restriction imposed are $\sum g_i = 0$ and $\sum S_{ij} + S_{ji} = 0$ (for each i)

The analysis of variance for combining ability was carried out using replication mean of each entry (diallel family) as follows:

Item	d.f.	Sum of squares	MSS	Expected MSS
GCA	P-1	S_g	M_g	$\sigma_e^2 + (P+2) \frac{1}{(P-1)} \sum g_i^2$
SCA	$P(P-1)/2$	S_s	M_s	$\sigma_e^2 + \frac{2}{P(P-1)} \sum_i \sum_j S_{ij}^2$
Error	$(b-1)(e-1)$	S_e	M_e'	σ_e^2

Where,

GCA = general combining ability

SCA = specific combining ability

p = Number of parents

b = Number of blocks or replications

e = Number of entry (family)

Y_i = Array total of the i^{th} parent

Y_{ij} = Mean value of the i^{th} parent

$Y.$ = Grand total of the $\frac{1}{2} p(p-1)$ crosses and parental lines

Y_{ij} = Progeny mean values in the diallel table

S_e = Sum of square due to error

$$S_g = \frac{1}{(P+2)} \left[\sum_i (Y_i + Y_{ii})^2 - \frac{4}{P} Y..^2 \right]$$

$$S_s = \sum_i \sum_j Y_{ij}^2 - \frac{1}{(P+2)} \sum_i (Y_i + Y_{ii})^2 + \frac{2}{(P+1)(P+2)} Y..^2$$



The GCA and SCA effects of each character were calculated as follows;

$$g_i = \frac{1}{(P+2)} \left[\sum_i (Y_i + Y_{ii})^2 - \frac{2}{p} Y_{..} \right]$$

$$s_{ij} = Y_{ij} - \frac{1}{(P+2)} \sum (y_i + y_{ii} + y_j + y_{ji}) + \frac{2}{(p+1)(p+2)} y_{..}$$

The variance of GCA and SCA were,

$$\text{Var}(g_i) = \frac{(p-1)}{p(p+2)} \sigma^2 e$$

$$\text{Var}(s_{ij}) = \frac{2(p-1)}{(p+1)(p+2)} \sigma^2 e (i \neq j)$$

Standard error (SE) of an estimate was calculated the square root of the variance of concerned estimate eg.

j Var(g;) and jVar(s.)

$$\sqrt{\text{Var}(g_i)} \text{ and } \sqrt{\text{Var}(s_{ij})}$$

3.9.2 Estimation of heterosis:

The amount of heterosis in the F_1 's was analysed using the following formulae:

$$\text{Heterosis over better parent \%} = \frac{\overline{F_1} - \overline{BP}}{\overline{BP}} \times 100$$

Here, $\overline{F_1}$ = Mean of F_1 individuals

\overline{BP} = Mean of the better parent values

$$\text{Heterosis over mid parent \%} = \frac{\overline{F_1} - \overline{MP}}{\overline{MP}} \times 100$$

Here, $\overline{F_1}$ = Mean of F_1 individuals

\overline{MP} = Mean of the mid parent values

CD (Critical Difference) values were used for testing significance of heterotic effects.

$$\text{Critical Differences (CD)} = t \times \sqrt{\frac{2 \text{ EMS}}{r}}$$

Here, EMS = Error Mean Sum of square

r = No. of replication

t = Tabulated t value at error df

CD values were compared with the values come from (F_1 -BP) and (F_1 -MP) to test significance of respective heterotic effects.

3.9.3 Graphical diallel analysis

Diallel analysis for the components of genetic variances and W_r - V_r graphs for all the characters studied were done according to Hayman (1954a,b). A diallel table was prepared from the averages over all the three replications and the following statistics were estimated.

V_r = Variance of all the progenies in each parental array (an array is a group of crosses involving a particular parents)

W_r = Covariance between parents and their offspring in each array

V_{OL0} = Variance of parents

V_{OL1} = Variance of the means of array

W_r^2 = The W_r for constructing the limiting parabola

$b_{w_r.v_r}$ = Regression of W_r on V_r

a = The Y- intercept

V_{1L1} = Mean of all the V_r values

W_{OLO1} = Mean of all the W_r values

Y_r' = Standardized mean for each parent

$(W_r + V_r)'$ = Standardized $(W_r + V_r)$ values for each parent

$r_{Y_r'(W_r+V_r)}$ = Correlation between parental order of dominance

$(M_{L1} - M_{LO})^2$ = Dominance relationship

r^2 = Possible limit of selection of parents showing dominance

The validity of Hayman's hypothesis was tested for all the characters studied by the equations.

3.9.3.1 Test of homogeneity of W_r - V_r variances

$$t^2 = \frac{n-2}{4} \left[\frac{(\text{Var}V_r - \text{Var}W_r)^2}{(\text{Var}V_r \times \text{Var}W_r) - \text{Cov}^2(V_r, W_r)} \right]$$

Where,

$\text{Var} V_r$ = Variance of the array variance

$\text{Var} W_r$ = Variance of the parent and array covariance

$\text{Cov}(V_r, W_r)$ = covariance of the variance and covariance

n = Number of parents involved in the diallel crosses

t = equivalent to a F-test with 4 and $(n-2)$ degrees of freedom

3.9.3.2 Test of deviation of regression slope from unity

i. Deviation from 0

$$t_1 = (b-0)/SE, \text{ (at } n-2 \text{ df)}$$


ii. Deviation from unity

$$t^2 = (1-b)/SE \text{ (at } n-2 \text{ df)}$$

Where,

b = regression co-efficient of W_r on V_r

SE_b = standard error of b



Chapter 4
Results and Discussion

RESULTS AND DISCUSSION

4.1 Mean performance

Mean performance of nine agronomic and yield related traits of parents and hybrid combinations are presented in Table 2.

4.1.1 Plant height

For parent, the lowest plant height was observed in BARI sarisha15 (100.77 cm) and for hybrid SAU sarisha 2 X BARI sarisha 15 (98.73 cm). Whereas the parent BARI sarisha 6 (114.33 cm) exhibited the highest plant height. The highest plant height was found from the hybrid Tori 7 X BARI sarisha 15 (113.40 cm). The hybrids were approximately 8-10 cm higher than the parents.

4.1.2 Days to 80% Maturity

Considering earliness, the parent Tori 7 (76 Days) showed the lowest duration for maturation but the parent BARI sarisha 6 (93 Days) had taken the highest duration. On the other hand, the hybrid combination Tori 7 X SAU sarisha 3 (75 Days) matured with the lowest growth duration, which was earlier than its parents.

4.1.3 Number of primary branches per plant

For this character the parents showed the value ranging from 7.62 to 5.67. The parent BARI sarisha 6 (7.67) showed the highest value. In the hybrid, the highest value was provided by the combination SAU sarisha 3 X BARI sarishab15 (7.33) which was higher than SAU sarisha 3 but lower than BARI sarisha 6.

4.1.4 Number of secondary branches per plant

For the number of secondary branches per plant, the parent Tori 7 (7.00) showed the highest value. Similarly in the hybrid, the highest value of number of secondary branches per plant was provided by the combination Tori 7 X SAU sarisha 2 (7.67) which was higher than Tori 7.

Table 2: Mean performance for 9 different characters in 5 parents and their 10 F₁'s of *Brassica rapa* L.

Treatments	Plant height (cm)	Days to 80% maturity	No. of primary branches/plant	No. of secondary branches/plant	No. of siliquae/plant	siliqua length (cm)	Seeds/siliqua	Seed yield/plant (gm)	1000 seed weight (gm)
Tori 7	103.87	76.00	6.67	7.00	151.00	5.37	15.33	5.40	2.27
SAU sarisha 2	104.73	79.00	6.00	6.00	144.67	5.17	17.00	4.20	2.83
SAU sarisha 3	102.50	84.00	6.00	6.00	158.33	5.27	13.00	5.07	2.93
BARI sarisha 6	114.33	93.00	7.67	6.67	163.33	5.07	13.33	5.30	3.07
BARI sarisha 15	100.77	84.00	6.67	0.00	83.00	4.97	22.33	3.93	2.53
Tori 7 x SAU sarisha 2	100.20	78.00	5.67	7.67	137.33	4.67	14.00	4.70	2.87
Tori 7 x SAU sarisha 3	99.50	75.00	6.00	3.67	131.00	5.10	14.33	3.57	2.67
Tori 7 x BARI sarisha 6	108.27	89.00	6.67	4.00	150.67	5.10	16.00	5.07	2.53
Tori 7 x BARI sarisha 15	113.40	82.00	7.33	5.00	164.00	4.90	14.67	5.13	2.73
SAU sarisha 2 x SAU sarisha 3	100.53	76.00	5.67	3.00	125.33	5.10	14.00	4.07	2.60
SAU sarisha 2 x BARI sarisha 6	107.77	84.00	6.00	5.00	156.33	5.07	12.33	4.63	3.00
SAU sarisha 2 x BARI sarisha 15	98.73	91.00	7.00	1.67	127.33	5.37	19.67	4.77	2.53
SAU sarisha 3 x BARI sarisha 6	111.87	78.00	6.33	4.00	140.00	5.00	14.33	4.30	2.53
SAU sarisha 3 x BARI sarisha 15	109.27	80.00	7.33	3.33	133.33	4.80	14.67	4.83	3.37
BARI sarisha 6 x BARI sarisha 15	112.10	91.00	7.00	4.00	160.67	4.97	15.33	5.33	3.10
Grand mean	105.86	83.00	6.53	4.47	141.76	5.06	15.36	4.69	2.77

4.1.5 Number of siliquae per plant

Number of siliquae per plant varied from 83.00 to 163.33 where the parent BARI sarisha 2 produced the highest and BARI sarisha 15 produced the lowest number of siliquae per plant. Considering hybrid performance, it was ranged from 125.33 to 164.00. The hybrid combination Tori 7 X BARI sarisha 15 (164.30) provided the highest number which was higher than its parent.

4.1.6 Siliqua length

Siliqua length also varied from 4.97 to 5.37 cm in parents. The parent BARI sarisha 15 produced the shortest siliqua while the parent Tori 7 produced the longest siliqua. On the other hand, the values varied from 4.67 to 5.37 cm for hybrids. In this regard, the hybrid combination SAU sarisha 2 X BARI sarisha 15 (5.37 cm) exhibited the highest length of siliqua and that was a little bit higher than that of its either parent.

4.1.7 Seeds per siliqua

Seed per siliqua also varied from 13.00 to 22.33 in parents and from 12.33 to 19.67 in hybrids. The hybrid SAU sarisha 2 X BARI sarisha 15 (19.67) produced an excellent number of seeds per siliqua. (Table 2).

4.1.8 Seed yield per plant

Seed yield per plant was found to be diversified in different genotypes including parents and hybrids. Seed yield of the genotypes varied from 3.93 to 5.40 gm in parents and from 3.57 to 5.33 gm in hybrids. The highest seed yield of the parent was found in Tori 7 (5.40) where as lowest in BARI sarisha 15 (3.93 gm). Similarly, the highest seed yield was also observed in the hybrid BARI sarisha 6 X BARI sarisha 15 (5.33 gm) which was almost higher than both of its parents.

4.1.9 Thousand seed weight

Thousand seed weight in *B. rapa* varied with some extent i.e. from 2.27 to 3.07 gm in parent and from 2.53 to 3.37 gm in hybrid. However, the heaviest seeds were produced by the parent BARI sarisha 6 (6.00 gm) and also by the hybrid combination SAU sarisha 3 X BARI sarisha 15 (3.37 gm). The hybrid provided the highest weighted seeds which were higher than its both parents (Table 2).

4.2 COMBINING ABILITY

The analysis of variance for the genotypes, combining ability variances, ratio of GCA and SCA variances, sum of square, estimates of general and specific combining ability effects were presented in Table 3 to Table 6. The analysis of variance carried out for nine characters was presented in Table 3 which indicated that the genotypes differed significantly for all the characters studied. Treatment mean sum of squares (mean of genotypes) were further partitioned into parents, crosses (hybrids) and parent vs crosses. Parents and crosses showed highly significant variances for days to maturity, no. of primary branches/ plant, no. of secondary branches/ plant, no. of siliquae/ plant, siliqua length, seed/ siliqua, seed yield/ plant and 1000 seed weight (Table 3). Variances due to parent vs cross interaction were also observed significant for most of the traits except plant height.

The general and specific combining ability effects are effective genetic parameters in the breeding program. Analysis of variances for yield and yield contributing characters (Table 3) revealed highly significant variation among the parents and hybrids indicating the presence of variability in the material. Variance due to genotypes was significant for most of the traits.

Combining ability analysis of five parents and ten F_1 's in half diallel cross was for nine quantitative traits. The variances due to general and specific combining ability were estimated for assessing the contribution of the additive and non-additive type of gene action involved in the inheritance of different characters. The mean sum of square due to general combining ability (GCA) was significant for all the traits except plant height, no. of primary branches/plant, siliqua length, seed yield/plant and 1000 seed weight indicating that the additive gene action was predominant for the expression of these characters (Table 4). The significant mean sum of square due to specific combining ability (SCA) was also observed for days to maturity and seed/ siliqua indicating that the non-additive gene actions were predominant for the expression of these characters (Table 4). The results showed the agreement with the findings of Malik *et al.* (1995); Thakur and Sagwal (1997) in rape seed. Similar findings were also reported by Tamber *et al.* (1991) in Indian mustard.

The higher magnitude of GCA variance was observed than that of SCA variance for plant height, days to maturity, no. of primary branches/ plant, no. of secondary branches/ plant, no. of siliquae/ plant, seed/siliqua, seed yield/ plant and 1000 seed weight. In an earlier study of Verma (2000), reported that SCA variance was higher than GCA variance (non-additive type) for seed yield per plant.

Table 3. Analysis of variances (MS values) for seed yield per plant and its component characters in *Brassica rapa* L.

Source of Variation	df	Plant height (cm)	Days to maturity	No. of Primary branches/ plant	No. of secondary branches/ Plant	No. of siliquae/ plant	Siliqua length (cm)	Seed/ siliqua	Seed yield/ plant (gm)	1000 seed weight (gm)
Replication	2	45.208	5.756 **	0.067	9.800	984.956	0.365	3.889 **	2.181	0.094
Genotype	14	89.747	112.356 **	1.229	12.705	1310.975	0.111	20.546 **	0.946	0.249
Parent	4	84.262	84.262 **	84.262 **	84.262 **	84.262	84.262 **	84.262 **	84.262 **	84.262 **
Cross	9	101.210	115.348 **	1.278	7.422	608.059	0.110	11.022 **	0.853	0.244
Parent vs Cross	1	8.525	197.795 **	331.349 **	225.983 **	12544.062 **	336.493 **	148.605 **	331.483 **	335.749 **
Error	28	85.343	13.827	1.352	7.276	1538.813	0.089	2.960	2.211	0.193

**p<0.01, *p<0.05

Table 4. Analysis of variances (MS values) for GCA and SCA for seed yield and yield contributing components in *Brassica rapa* L.

Source of Variation	df	Plant height (cm)	Days to maturity	No. of primary branches/plant	No. of secondary branches/Plant	No. of siliquae/plant	Siliqua length (cm)	Seed/siliqua	Seed yield/Plant (gm)	1000 seed weight (gm)
GCA	4	56.467	76.838 **	0.862	8.576 *	653.203 *	0.009	15.259 **	0.305	0.083
SCA	10	9.648	10.849 **	0.114	1.249	175.253	0.024	1.742 *	0.160	0.042
GCA/ SCA	--	5.853	7.083	7.542	6.865	3.727	0.361	8.758	1.909	1.994
Error	28	28.448	4.609	0.451	2.425	512.938	0.030	0.987	0.737	0.064

**p<0.01, *p<0.05



4.2.1 General combining ability (GCA) Effects

The additive nature and magnitude of gene action for a trait could be measured by estimation of GCA effects. Similarly the magnitude and nature of non-additive i.e. dominance and epistasis nature of gene actions could be measured by estimation of SCA effects. A parent with higher significant GCA effects is considered as a good general combiner. A parent showing high GCA and SCA variances is a better parent for creating high yielding specific combination. Parents with significant high GCA effect could be used in conventional breeding programme and crosses with significant high SCA effect could be used in hybrid development. The estimates of GCA effects are presented in Table 5. The magnitude and direction of the significant GCA effects for six parents provided meaningful comparisons and would give a clue to design the future breeding programme. The results of GCA effects of different characters are presented as follows:

4.2.1.1 Plant height:

Out of five parental GCA, four parents showed negative GCA effect. The highest negative GCA effects (-2.633) was provided by SAU sarisha 2. The other parents which represented negative GCA were Tori 7 (-0.862), BARI sarisha 15 (-0.014) and SAU sarisha 3 (-1.281). Those parents with positive and significant GCA effects were considered as good general combiner for the trait aimed to promote desirable plant height in their crosses (Table 5). The parent BARI sarisha 6 (4.790) showed positive GCA effects to promote the plant height in *Brassica rapa*. Chowdhury *et al.* (2004) obtained dwarfness in YSK-8501 in *Brassica campestris* L. Singh *et al.* (1996) observed dwarfness in glossy mutant in *Brassica juncea* L.

4.2.1.2 Days to maturity:

The parent BARI sarisha 6 (4.505) provided the highest significant and positive GCA effects for days to maturity followed by BARI sarisha 15 (2.505) hence the parents were desirable general combiners to promote the earliness in *Brassica rapa* L. (Table 5). Parent SAU sarisha 3 (-2.829) showed the highest negative GCA effects followed by Tori 7 (-2.683) and SAU sarisha 2 (-1.543). Chowdhury *et al.* (2004) observed earliness in Din-2 in *Brassica rapa* L. Singh *et al.* (2000) found earliness in YSC-68 in *Brassica campestris* L.

Table 5: General combining ability (GCA) effects for 5 parents in 5x5 half diallel crosses of *Brassica rapa* L.

Parents	Plant height (cm)	Days to 80% maturity	No of primary branches/ Plant	No. of secondary branches/ plant	No. of siliquae/ plant	Siliqua length (cm)	Seeds/ siliqua	Seed yield/ plant (gm)	1000 seed weight (gm)
Tori 7	-0.862	-2.638	-0.029	1.076	4.924	0.020	-0.352	0.164	-0.185
SAU sarisha 2	-2.633	-1.543	-0.410	0.362	-2.124	0.025	0.267*	-0.222	0.006
SAU sarisha 3	-1.281	-2.829	-0.267	-0.114	-0.600	0.025	-1.257	-0.174	0.058
BARI sarisha 6	4.790 *	4.505 *	0.305	0.505	11.971	-0.013	-1.067 *	0.259	0.096
BARI sarisha 15	-0.014	2.505 *	0.400	-1.829 *	-14.171	-0.056	2.410 *	-0.027	0.025
SE(gl)	2.016	0.811	0.254	0.589	8.560	0.065	0.375	0.324	0.096
SE(sij)	5.980	2.407	0.753	1.746	25.394	0.193	1.114	0.962	0.284

**p<0.01, *p<0.05

4.2.1.3 Number of primary branches per plant:

There was two parent viz. BARI sarisha 15 (0.400) and BARI sarisha 6 (0.305) provided positive GCA effects out of five parents which indicated that this parent was good general combiners for promising primary branches. So thus parent was considered as good to use in the breeding programme for more primary branches (Table 5). Other parents showed negative and insignificant effects. Chowdhury *et al.* (2004) obtained more primary branches on Sampan in *Brassica rapa* L. Singh *et al.* (2000) observed maximum number of primary branches on YSP-842 in *Brassica campestris* L.

4.2.1.4 Number of secondary branches per plant:

For number of secondary branches per plant, the positive GCA effects were observed in Tori 7 (1.076) and considered as the best general combiner for the trait. The parents BARI sarisha 15 showed significant but negative GCA effects and other three demonstrated insignificant GCA effects. Singh *et al.* (1996) obtained the highest secondary branches in BJ-1235 in *Brassica juncea* L. Chowdhury *et al.* (2004a) observed more secondary branches in Din-2 in *Brassica rapa* L.

4.2.1.5 Number of siliquae per plant:

The parent BARI sarisha 6 exhibited the highest (11.971) positive GCA effects for this character. This parent was selected as the best general combiner and desirable to use in hybridization program to improve the number of siliquae per plant in *Brassica rapa* L. (Table 5). On the other hand, the highest negative and insignificant GCA value were provided by BARI sarisha 15 (-14.171) and SAU sarisha 2 (-2.124). Chowdhury *et al.* (2004) found the highest number of silliquae in Din-2 in *Brassica rapa*. Singh and Murty (1980) obtained maximum number of siliquae per plant in SS-1 in *Brassica campestris* L.

4.2.1.6 Siliqua length:

Out of five parents, three parents were exhibited insignificant and positive GCA effect. Other two parents showed insignificant and negative GCA values. None parent could be used to improve this character. Sheikh and Singh (1998) obtained different result (maximum siliqua length) in glossy mutant.

4.2.1.7 Number of seeds per siliqua:

Out of five parents, only BARI sarisha 15 and SAU sarisha 2 were exhibited significant and positive GCA effect (2.410) and (0.267), respectively. So the parents would be considered as general combiner for the character and could be used for hybrid production with more seeds per siliqua development in breeding programme. Other one parents showed significant but negative GCA values. Rest of the parents provided insignificant and negative GCA effects. Chowdhury *et al.* (2004) found maximum seeds per siliqua in Dhali in *Brassica rapa* L. Singh and Murty (1980) obtained more seeds per siliqua in YPS-842 in *Brassica campestris* L.

4.2.1.8 Seed yield per plant:

The highest positive GCA effects were observed in BARI sarisha 6 (0.259). This parent might be selected as promising general combiner for high yield. On the other side, SAU sarisha 2 (-0.222), SAU sarisha 3 (-0.174) and BARI sarisha 15 (-0.027) provided insignificant and negative GCA effects and were not fit for increasing seed yield. Other parents showed insignificant GCA effect for this character (Table 5) indicating that the parents were not good for improving seed yield per plant. Chowdhury *et al.* (2004a) found the highest seed yield per plant in Pt-303 in *Brassica rapa* L.

4.2.1.9 Thousand seed weight

Out of five parents, four parents were exhibited insignificant and positive GCA effect. BARI Sarisha 6 (0.096) showed highest positive GCA effect and could be considered as the best general combiner for this trait. Chowdhury *et al.* (2004a) found the highest seed weight in Dhali in *Braasiea rapa* L.

4.2.2 Specific combining ability (SCA) effects

The specific combining ability effects signify the role of non-additive i.e. dominance and or epistatic gene action in the expression of the characters. It denotes the highly specific combining ability leading to highest performance of some specific cross combinations. For this reason it relates to a particular cross. The specific combining ability effects are also seen in relation to their size. High SCA effects may arise not only on cross involving high × high combinations, but also in those involving low × high and also from low × low. Thus in practice, some of the low combiners should also be accommodated in hybridization programme. The specific combining ability effects of ten crosses for the different characters studied are presented in Table 6. The magnitude and direction of the significant effects for the ten crosses provide meaningful comparisons and would give a clue to the future breeding programme. The results of SCA effects for different characters are given below:

4.2.2.1 Plant height

Out of 10 crosses, all crosses showed the significant and positive SCA effects. The highest positive SCA effects were observed in Tori 7 X BARI sarisha 15 (30.99) for plant height and could be considered as the best combination. (Table 6). Chowdhury *et al.* (2004a) observed dwarfness in PT-303 x Tori-7 in *Brassica rapa*. Acharya and Swain (2004) obtained dwarfness in Varuna x Pusa Bahar in *Brassica juncea*.

4.2.2.2 Days to maturity

The cross combination SAU sarisha 2 X BARI sarisha 15 (25.35) showed the highest significant and positive SCA effects while the significant and positive value for this parameter was obtained from all crosses. Hence, the cross combination SAU sarisha 2 X BARI sarisha 15 was the best for early maturity in *Brassica rapa* (Table 6). Chowdhury *et al.* (2004a) observed earliness in M-27 x Din-2 in *Brassica rapa*. Singh *et al.* (2000) obtained earliness in SS-3 x SS-1 in *Brassica campestris*. Acharya and Swain (2004) found early maturity in JC 26 x Jai kisan in *Brassica juncea*.

Table 6: Specific combining ability (SCA) effects for 10 crosses in 5x5 half diallel crosses of *Brassica rapa* L.

Cross combination	Plant height (cm)	Days to 80% maturity	No. of primary branches /plant	No. of secondary branches/ plant	No. of siliquae/ plant	Siliqua length (cm)	Seeds/ siliqua	Seed yield/ plant (gm)	1000 seed weight (gm)
Tori 7 X SAU sarisha 2	20.09 **	16.18 **	0.92	2.90	23.50	0.65 **	2.01	1.07	0.85 *
Tori 7 X SAU sarisha 3	18.20 *	14.97 **	1.13	-0.68	15.84	1.09 **	3.68 **	-0.11	0.60
Tori 7 X BARI sarisha 6	21.66 **	22.56 **	1.29	-0.89	24.50	1.12 **	5.18 **	1.01	0.43
Tori 7 X BARI sarisha 15	30.99 **	20.31 **	1.88 *	2.15	60.71 *	0.96 **	0.80	1.33	0.70 *
SAU sarisha 2 X SAU sarisha 3	20.79 **	14.68 **	1.13	-0.73	16.34	1.08 **	2.80 *	0.73	0.37
SAU sarisha 2 X BARI sarisha 6	22.71 **	15.93 **	0.96	0.73	36.34	1.08 **	0.97	0.92	0.73 *
SAU sarisha 2 X BARI sarisha 15	17.88 *	25.35 **	1.88 *	-0.56	30.21	1.42 **	5.26 **	1.30	0.33
SAU sarisha 3 X BARI sarisha 6	25.62 **	11.72 **	1.17	0.15	18.67	1.01 **	4.30 **	0.54	0.22
SAU sarisha 3 X BARI sarisha 15	27.23 **	15.14 **	2.08 *	1.52	34.88	0.85 **	1.59	1.33	1.12 **
BARI sarisha 6 X BARI sarisha 15	24.75 **	19.72 **	1.25	1.65	51.21 *	1.05 **	2.09	1.45	0.82 *
SEd(gi-gj) =	2.85094	1.147543	0.358885	0.832449	12.10593	0.092042	0.530976	0.458871	0.135437
SEd(Sij-Sik) =	6.532324	2.629352	0.822308	1.907379	27.73817	0.210894	1.216618	1.051405	0.310325
SEd(Sij-Skl) =	6.374895	2.565985	0.80249	1.861411	27.06969	0.205811	1.187297	1.026066	0.302846

**p<0.01, *p<0.05

4.2.2.3 Number of primary branches per plants

The cross combination SAU sarisha 3 X BARI sarisha 15 (2.08) was found to be the best to improve this crop with more number of primary branches as it showed the highest significant and positive SCA effects for this trait (Table 6). Chowdhury *et al.* (2004a) found more primary branches in Sampad x Tori-7 in *Brassica rapa*. Singh *et al.* (2000) obtained maximum number of primary branches per plant in YSK-8501 x SS-1 in *Brassica rapa*. Sheikh and Sing (1998) observed best positive effect in Pusa x Barani in *Brassica juncea*.

4.2.2.4 Number of secondary branches per plant

The highest positive value for this character was revealed by Tori 7 X BARI sarisha 15 (2.15) followed by BARI sarisha 6 X BARI sarisha 15 (1.65). On the other hand, the highest negative SCA value was provided by Tori 7 X BARI sarisha 6 (-0.89). Thus, Tori 7 X BARI sarisha 15 (2.15) and BARI sarisha 6 X BARI sarisha 15 (1.65) were the best cross combinations to improve plants with more number of secondary branches and Tori 7 X BARI sarisha 6 was the best combination to obtain plants with minimum secondary branches (Table 6). Chowdhury *et al.* (2004a) found maximum secondary branches in Sampad x Din-2 in *Brassica rapa*. Singh *et al.* (2000) observed the highest secondary branches in YSC-68 x SS-2 in *Brassica rapa*. Acharya and Swain (2004) obtained more secondary branches in BM 20-12-3 x JC 26 in *Brassica juncea*.

4.2.2.5 Number of Siliquae per plant

Among the cross combinations, Tori 7 X BARI sarisha 15 showed the highest significant and positive SCA effects (60.71) followed by BARI sarisha 6 X BARI sarisha 15 (51.21). (Table 6). So, Tori 7 X BARI sarisha 15 was the best combination for this trait. Chowdhury *et al.* (2004a) found maximum siliquae in Sampad x Din-2 in *Brassica rapa*. Acharya and Swain (2004) obtained the highest siliquae per plant in Pusa Bahar x JC 26 in *Brassica juncea*.

4.2.2.6 Thousands seed weight:

The highest significant and positive SCA value was exhibited by SAU sarisha 3 X BARI sarisha 15 (1.12) followed by Tori 7 X SAU sarisha 2 (0.85), BARI sarisha 6 X BARI sarisha 15 (0.82), SAU sarisha 2 X BARI sarisha 6 (0.73). So, SAU sarisha 3 X BARI sarisha 15 might be the best cross combination for this trait.

4.2.2.7 Siliqua length:

Among the cross combinations, SAU sarisha 2 X BARI sarisha 15 (1.42) showed the highest significant and positive SCA effects followed by Tori 7 X BARI sarisha 6 (1.12) and Tori 7 X SAU sarisha 3 (1.09). On the other hand, the remaining combinations showed significant but positive SCA effects for the trait (Table 6). Hence, the cross combination SAU sarisha 2 X BARI sarisha 15 was the best for siliqua length. Sheikh and Singh (1998) and Acharya and Swain (2004) observed the maximum siliqua length in Pusa Barani x Glossy mutant and BM 20-12-3 x Pusa Bahar respectively in *Brassica juncea*.

4.2.2.8 Number of seeds per siliqua:

Among the cross combinations, SAU sarisha 2 X BARI sarisha 15 exhibited the highest significant and positive SCA value (5.26) followed by Tori 7 X BARI sarisha 6 (5.18), SAU sarisha 3 X BARI sarisha 6 (4.30), Tori 7 X SAU sarisha 3 (3.68), and SAU sarisha 2 X SAU sarisha 3 (2.80). Hence, SAU sarisha 2 X BARI sarisha 15 was the best cross combination to increase the number of seeds in the siliqua as well as seed yield (Table 6). Chowdhury *et al.* (2004a) found height seeds per siliqua in Dhali x Sampad in *Brassica rapa*. Singh *et al.* (2000) obtained more seeds per siliqua in YSP-842 x YSK-8501 in *Brassica campestris*. Acharya and Swain (2004) observed the maximum number of seeds per siliqua in BM 20-12-3 x Pusa Bahar in *Brassica juncea*.

4.2.2.9 Seed yield per plant:

The cross combination BARI sarisha 6 X BARI sarisha 15 (1.45) exhibited the highest positive SCA effects for seed yield per plant. On the other hand, the highest negative SCA value was provided by Tori 7 X SAU sarisha 3 (-0.11) but it was also insignificant. So, BARI sarisha 6 X BARI sarisha 15 might be the best cross combination for the improvement of seed yield per plant in *Brassica rapa* (Table 6). Chowdhury *et al.* (2004a) obtained the highest seed yield in M-27 x Din-2 in *Brassica rapa*. Singh *et al.* (2000) observed more seed yield per plant in YSP-842 x YSK-8501 in *Brassica campestris*. Acharya and Swain (2004) found the maximum seed yield in Pusa Bold x Pusa Bahar in *Brassica juncea*.

4.3 Heterosis

Nine yield contributing characters of *Brassica rapa* were studied in five parental genotypes and their 10 hybrids obtained from 5x5 half diallel crosses. Percent heterosis for 9 different characters of the F₁ hybrids over their respective mid and better parental values is shown in Table 7. These results on heterosis of 10 F₁'s are described by character wise below.

4.3.1 Plant height

Out of 10 crosses, two hybrids showed significant heterosis over mid parent and four over better parent for plant height (Table 7). The significant heterosis over mid-parent ranged from -3.93% to 10.83% which were represented by Tori 7 X BARI sarisha 15 (10.83%) and Tori 7 X SAU sarisha 2 (-3.93%). On the other hand, the significant value of better parent heterosis was ranged from -5.74% to 9.18%. The highest better parent heterosis was obtained from Tori 7 X BARI sarisha 15 (9.18%) and the lowest better parent heterosis was produced by SAU sarisha 2 X BARI sarisha 6 (-5.74%). Lefort *et al.* (1987) while studying *Brassica napus* of Asian and European parental lines and their hybrids reported that plant height and seed yield showed positive heterosis in the hybrids. So, the hybrid Tori 7 X SAU sarisha 2 showed the highest negative significant heterosis (-3.93% and -4.33% over mid and better parent, respectively) for plant height (Plate 5).

4.3.2 Days to 80% maturity

Out of 10 cross combinations, five cross combinations showed significant heterosis over mid parent including both positive and negative values. For this character, negative heterosis is usually useful to obtain early hybrid. In this regard, only SAU sarisha 3 X BARI sarisha 6 (-11.86%) showed significant and negative heterosis over mid parent (Plate 6). For better parent heterosis, five hybrids showed significant heterosis. The combination SAU sarisha 3 X BARI sarisha 6 (-16.13 %) possessed the highest negative heterosis. Thus, the combinations with significant and negative heterosis over both mid parent and better parent might be useful for development of early *Brassica* variety. Kumar *et al.* (2002), Mahak *et al.* (2003a) and Das *et al.* (2004) found significant heterosis for days to maturity over mid parent and better parent which was disagreement with the majority of the findings of the parent crossed.

Table 7. Percent heterosis over mid-parent and better parent for 9 different characters in intervarietal hybrids of oleiferous *Brassica rapa* L.

Cross combination	Plant height (cm)		Days to 80% maturity		No. of primary branches/plant	
	Heterosis over mid-parent (%)	Heterosis over better parent (%)	Heterosis over mid-parent (%)	Heterosis over better parent (%)	Heterosis over mid-parent (%)	Heterosis over better parent (%)
Tori 7 X SAU sarisha 2	-3.93*	-4.33*	0.65	-1.27	-10.53	-15.00
Tori 7 X SAU sarisha 3	-3.57	-4.20	-6.25*	-10.71**	-5.26	-10.00
Tori 7 X BARI sarisha 6	-0.76	-5.31*	5.33*	-4.30	-6.98	-13.04
Tori 7 X BARI sarisha 15	10.83*	9.18	2.50	-2.38	10.00*	10.00
SAU sarisha 2 X SAU sarisha 3	-2.98	-4.01	-6.75**	-9.52**	-5.56	-5.56
SAU sarisha 2 X BARI sarisha 6	-1.61	-5.74*	-2.33	-9.68**	-12.20*	-21.74*
SAU sarisha 2 X BARI sarisha 15	-3.91	-5.73*	11.66**	8.33**	10.53*	5.00
SAU sarisha 3 X BARI sarisha 6	3.18	-2.16	-11.86**	-16.13**	-7.32	-17.39
SAU sarisha 3X BARI sarisha 15	7.51	6.60	-4.76	-4.76	15.79**	10.00
BARI sarisha 6 X BARI sarisha 15	4.23	-1.95	2.82	-2.15	-2.33	-8.70
Maximum	10.83	9.18	11.66	8.33	15.79	10.00
Minimum	-3.93	-5.74	-11.86	-16.13	-12.20	-21.74

**p<0.01,*p<0.05

Table 7. Continued

Cross combination	No. of secondary branches/plant		No of Siliqua/plant		Siliqua length (cm)	
	Heterosis over mid-parent (%)	Heterosis over better parent (%)	Heterosis over mid-parent (%)	Heterosis over better parent (%)	Heterosis over mid-parent (%)	Heterosis over better parent (%)
Tori 7 X SAU sarisha 2	17.95*	9.52*	-7.10	-9.05	-11.39**	-13.04**
Tori 7 X SAU sarisha 3	-43.59	-47.62	-15.30	-17.26	-4.08	-4.97
Tori 7 X BARI sarisha 6	-41.46	-42.86	-4.14	-7.76	-2.24	-4.97
Tori 7 X BARI sarisha 15	42.86*	-28.57	40.17*	8.61*	-5.16	-8.70**
SAU sarisha 2 X SAU sarisha 3	-50.00*	-50.00	-17.27	-20.84	-2.24	-3.16
SAU sarisha 2 X BARI sarisha 6	-21.05	-25.00	1.52	-4.29	-0.98	-1.94
SAU sarisha 2 X BARI sarisha 15	-44.44*	-72.22*	11.86*	-11.98	5.92*	3.87*
SAU sarisha 3 X BARI sarisha 6	-36.84	-40.00	-12.95	-14.29	-3.23	-5.06
SAU sarisha 3X BARI sarisha 15	11.11*	-44.44	10.50	-15.79	-6.19	-8.86*
BARI sarisha 6 X BARI sarisha 15	20.00	-40.00	30.45*	-1.63	-1.00	-1.97
Maximum	42.86	9.52	40.17	8.61	5.92	3.87
Minimum	-50.00	-72.22	-17.27	-20.84	-11.39	-13.04

**p<0.01,*p<0.05

Table 7. Continued

Cross combination	Seeds/ Siliqua		Seed yield/plant (gm)		1000 seed weight (gm)	
	Heterosis over mid-parent (%)	Heterosis over better parent (%)	Heterosis over mid-parent (%)	Heterosis over better parent (%)	Heterosis over mid-parent (%)	Heterosis over better parent (%)
Tori 7 X SAU sarisha 2	-13.40*	-17.65*	-2.08	-12.96	12.42	1.18
Tori 7 X SAU sarisha 3	1.18	-6.52	-31.85	-33.95*	2.56	-9.09*
Tori 7 X BARI sarisha 6	11.63**	4.35	-5.30	-6.17	-5.00	-17.39
Tori 7 X BARI sarisha 15	-22.12**	-34.33**	10.00*	-4.94	13.89*	7.89*
SAU sarisha 2 X SAU sarisha 3	-6.67	-17.65*	-12.23	-19.74*	-9.83**	-11.36
SAU sarisha 2 X BARI sarisha 6	-18.68**	-27.45**	-2.46	-12.58	1.69	-2.17
SAU sarisha 2 X BARI sarisha 15	0.00	-11.94	17.21*	13.49*	-5.59**	-10.59**
SAU sarisha 3 X BARI sarisha 6	8.86*	7.50	-17.04	-18.87	-15.56	-17.39
SAU sarisha 3X BARI sarisha 15	-16.98**	-34.33**	7.41	-4.61	23.17*	14.77
BARI sarisha 6 X BARI sarisha 15	-14.02*	-31.34**	15.52*	0.63	10.71	1.09
Maximum	11.63	7.50	17.21	13.49	23.17	14.77*
Minimum	-22.12	-34.33	-31.85	-33.95	-15.56	-17.39

**p<0.01,*p<0.05



Tallest plant (single plant view)



Shortest plant (single plant view)



SAU sarisha 2 X BARI sarisha 15



Tori 7 X SAU sarisha 2

Plate 5: Photograph showing variation for plant height between SAU sarisha 2 X BARI sarisha 15 and Tori 7 X SAU sarisha 2 of *Brassica rapa* genotypes.



Early maturity line (SAU sarisha 3 X BARI sarisha 6)



Late maturity line (SAU sarisha 2 X BARI sarisha 15)

Plate 6: Photograph showing variation between early (SAU sarisha 3 X BARI sarisha 6) and late (SAU sarisha 2 X BARI sarisha 15) line of *Brassica rapa* genotype.

4.3.3 Number of primary branches per plant

Only three hybrids showed significant and positive mid-parent heterosis which ranged from 10.00 % to 15.79 % for number of primary branches per plant. The hybrid SAU sarisha 3 X BARI sarisha 15 showed the highest (15.79 %) significant and positive heterosis where as the lowest (10.00 %) value was found from Tori 7 X BARI sarisha 15. In case of better parent heterosis, the significant values ranged from -21.74 to 10.00 %. The combination SAU sarisha 3 X BARI sarisha 15 and Tori 7 X BARI sarisha 15 showed the highest positive value followed by SAU sarisha 2 X BARI sarisha 15 (5.00 %). Thus, the hybrid SAU sarisha 3 X BARI sarisha 15 showing higher branches than its parent (Plate 7). Thakur and Segwall (1997) found a heterosis value ranging from -26.0 to 193.6% over better parent for the character primary branches in rapeseed (*Brassica napus* L.). Yadav *et al.* (2004) observed that Trachystome × PR 905 showed 106.5% and 100.0% heterosis over BP and SV, respectively for the number of primary branches per plant.



SAU sarisha 2 X BARI sarisha 6



SAU sarisha 3 X BARI sarisha 15

Plate 7: Photograph showing variation for number of primary branches / plant between SAU sarisha 2 X BARI sarisha 6 (lowest) and SAU sarisha 3 X BARI sarisha 15 (highest) of *Brassica rapa* genotype.

4.3.4 Number of secondary branches plant

Out of 10 crosses five showed significant mid parent heterosis, in which three hybrids showed positive values ranging from 11.11 % to 42.86 %. The highest significant and positive mid-parent heterosis was produced by the combination Tori 7 X BARI sarisha 15 (42.86) followed by Tori 7 X SAU sarisha 2 (17.95%) and SAU sarisha 3 X BARI sarisha 15 (11.11%). The hybrid Tori 7 X BARI sarisha 15 showed higher branches than its parent (Plate 8). These hybrids could be considered for further development of heterotic *Brassica* hybrid. On the other hand, one cross combinations were found to be significant and positive for better parent heterosis Tori 7 X SAU sarisha 2 (9.52 %) exhibited the highest value . These hybrid might be selected for number of secondary branches per plant. Kumar *et al.* (1990) found positive heterosis for number of secondary branches per plant and they also recorded the highest heterobeltiosis for number of secondary branches per plant. Yadav *et al.* (2004) observed maximum heterosis over BP in *Trachystoma* × PHR-1 (125.1%) and over CV in *Moricandia* × NRCM-79 (9.6%) for the number of secondary branches per plant.



SAU sarisha 2 X SAU sarisha 3



Tori 7 X BARI sarisha 15

Plate 8: Photograph showing variation for number of secondary branches / plant between SAU sarisha 2 X SAU sarisha 3 (lowest) and Tori 7 X BARI sarisha 15 (highest) of *Brassica rapa* genotype.

4.3.5 Number of siliquae per plant

The highly significant and positive mid-parent heterosis for siliquae per plant was found in three hybrids ranged from 11.86% to 40.17%. The hybrid Tori 7 X BARI sarisha 15 represented the highest heterosis over mid-parent (40.17%), where as the lowest was found in SAU sarisha 3 X BARI sarisha 15 (11.86%). On the other hand, only one combination showed significant and positive value for better parent heterosis Tori 7 X BARI sarisha 15 produced the highest heterotic value (8.61%). However, the hybrid Tori 7 X BARI sarisha 15 having significant and positive heterosis produced more siliquae per plant than any parent and could be selected for further evaluation (Plate 9). Zheng and Fu (1991) found positive heterosis of 51.47% over mid parent in the F₁ hybrids in *Brassica nigra* for number of siliquae per plant. Thakur and Segwal (1997) estimated positive heterosis over better parent ranging from 21.9 to 162.6% in rape seed for siliquae per plant. Qi *et al.* (2003) observed the forty-seven crosses gave on average 28.02% (0.93-97.87%) more siliquae per plant.



Tori 7 X BARI sarisha 15



SAU sarisha 2 X SAU sarisha 3

Plate 9: Photograph showing variation for number of siliquae / plant between Tori 7 X BARI sarisha 15 (highest) and SAU sarisha 2 X SAU sarisha 3 (lowest) of *Brassica rapa* genotype.

4.3.6 Siliqua length

Out of 10 combinations, two combinations showed significant heterosis over mid parent but only one of them showed positive and significant heterosis SAU sarisha 2 X BARI sarisha 15 (5.92%), In case of better parent heterosis, the significant and positive heterosis was found in SAU sarisha 2 X BARI sarisha 15 (3.87%) Other hybrids produced significant but negative heterotic value. For siliqua length, the hybrid SAU sarisha 2 X BARI sarisha 15 showed higher siliqua length than its parent (Plate 10). Kumar *et al.* (1990) found positive heterosis for length of siliqua in *Brassica juncea*.



SAU sarisha 2 X BARI sarisha 15



Tori 7 X SAU sarisha 2

Plate 10: Photograph showing variation for siliqua length between SAU sarisha 2 X BARI sarisha 15 (highest) and Tori 7 X SAU sarisha 2 (lowest) of *Brassica rapa* genotype.

4.3.7 Number of seeds per siliqua

Out of 10 combinations, seven combinations showed significant heterosis over mid parent but only two of them showed positive and significant heterosis Tori 7 X BARI sarisha 6 (11.63%) and SAU sarisha 3 X BARI sarisha 6 (8.86%) . On the other hand, no combinations were found to be significant and positive for better parent heterosis except SAU sarisha 3 X BARI sarisha 6 (7.50%). However, the mentioned combinations showed significant and positive heterosis over mid parent as well as better parent and could be considered for further evaluation. Kumar *et al.* (1990) reported positive heterosis for number of seeds per siliqua in *Brassica juncea*. Yadav *et al.* (2004) observed that the Siifolia × SM-1 showed 54.1% heterosis over BP and 9.2% negative heterosis over SV for seeds per siliqua.

4.3.8 Seed yield per plant

In case of yield per plant, only SAU sarisha 2 X BARI sarisha 15 represented significant and positive value for both mid parent (17.21%) and better parent (13.49%) heterosis. This combination could be selected for improvement of yield performance. Tyagi *et al.* (2001) found the highest standard heterosis (206.14%) and heterobeltiosis (240.56%) for seed yield per plant in the cross BIO 772 × Rohini. Adefris *et al.* (2005) observed that seed yield showed the highest relative mid parent heterosis that varied from 25 to 145% with a mean of 67% and relative high parent heterosis varied from 16 to 124% with a mean of 53%. The presence of high levels of mid and high parent heterosis indicated a considerable potential to embark on breeding hybrid or synthetic cultivars in mustard. Shen *et al.* (2005) observed mid parent heterosis and high parent heterosis of seed yield per plant ranged from 5.50 to 64.11% and from -2.81 to 46.02% respectively. Wang *et al.*(1999) analysed heterosis and combining abilities of 20 reciprocal cross combinations of five rape (*Brassica napus*) cultivars (lines) showing high seed yield .

4.3.9 Thousand seed Weight

Out of 10 hybrids, only two showed significant and positive heterosis over mid parent. They were SAU sarisha 3 X BARI sarisha 15 (23.17%) and Tori 7 X BARI sarisha 15 (13.89%). Similarly for better parent heterosis, the combinations SAU sarisha 3 X BARI sarisha 15 (14.77%) and Tori 7 X BARI sarisha 15 (7.89%) showed the highest significant and positive heterosis. The hybrids of SAU sarisha 3 X BARI sarisha 15, Tori 7 X BARI sarisha 15, Tori 7 X SAU sarisha 2 , BARI sarisha 6 X BARI sarisha 15 were positively significant over both mid and better parent heterosis. Yadav *et al.* (2004) observed the highest heterosis for thousand seed weight in Moricandia × PHR-1 (48.80%) followed by Trachystoma NRCM 69 (20.6%) over BP and SV respectively. Qi *et al.* (2003) observed that eight crosses showed better parent heterosis (3.57 to 20.48%) in thousand seed weight.

4.4 Vr-Wr graph

Vr-Wr graphs, the two dimensional depiction made based on the parental variance (V_r) and parent offspring co-variance (W_r) are presented in the Fig. 2 to Fig.11. Hayman's graphic approach to diallel analysis is based on monogenic additive model. The regression coefficient differ significantly from zero and approaching to unity for all the traits studied suggesting that there was no epistasis for most of the traits indicated the validity of such type of analysis. Vr-Wr graphs for the ten characters are described below:

4.4.1 Plant height




Fig 2: Vr-Wr graph for plant height in *Brassica rapa*

The regression line intersected above the point of origin suggesting partial dominance gene action for controlling the trait (Fig. 2). The distribution of array points indicated two parents BARI sarisha 6 (P4) and SAU sarisha 2 (P2) contained most dominant alleles as they fell closer to the point of origin. The parents Tori 7 (P1) fallen at the middle portion means that contained equal frequencies of dominant and recessive alleles. Whereas, rest of the parents fallen far from the origin indicated that they possessed maximum frequency of recessive alleles. Chowdhury *et al.* (2004b) obtained nearly complete dominance in *Brassica rapa*.

4.4.2 Days to maturity




Fig 3: Vr-Wr graph for days to maturity in *Brassica rapa*

The regression line intersected below the point of origin suggesting over dominance gene action for controlling the trait (Fig. 3). The parents SAU sarisha 3 (P3) contained maximum dominant alleles as they fell very closer to the point of origin. The parents BARI sarisha 15 (P5) fallen at the middle portion means that contained equal frequencies of dominant and recessive alleles. Whereas, rest of the parents fallen far from the origin indicated that they possessed maximum frequency of recessive alleles. Chowdhury *et al.* (2004b) observed partial dominance in *Brassica rapa*, Trivedi and Mukharjee (1986) found over dominance in *Brassica juncea*.

4.4.3 Primary branches per plant




Fig 4: Vr-Wr graph for primary branches per plant in *Brassica rapa*

The regression line intersected the W_r axis below the point of origin indicating the existence of over dominance gene action for controlling the trait (Fig. 4). The parent BARI sarisha 15 (P5) fell closer to the origin means they contained maximum frequencies of dominant alleles. The parent SAU sarisha 2 (P2) fallen at the middle portion means that contained equal frequencies of dominant and recessive alleles. Whereas, rest of the parents fallen far from the origin indicated that they possessed maximum frequency of recessive alleles. Chowdhury *et al.* (2004b) observed nearly complete dominance for the character in *Brassica rapa*, and Yadav and Yadava (1996) found over dominance in *Brassica campestris*.

4.4.4 Secondary branches per plant




Fig 5: Vr-Wr graph for secondary branches per plant in *Brassica rapa*



The regression line intersected the W_r axis below the point of origin indicating the existence of over dominance gene action for controlling the trait (Fig. 5). The parents SAU sarisha 3 (P3) and BARI sarisha 6 (P4) fell closer to the point of origin suggesting they contained maximum number of dominant alleles. The parents Tori 7 (P1) and BARI sarisha 6 (P2) fallen at the middle portion means they contained equal frequencies of dominant and recessive alleles. The parent SAU sarisha 2 (P2) and BARI sarisha 15 (P5) fell far from the origin indicating the presence of maximum frequency of recessive alleles. Chowdhury *et al.* (2004b) obtained partial dominance in *Brassica rapa*, Yadav and Yadava (1996) observed over dominance in *Brassica campestris*.

4.4.5 Siliquae per plant




Fig 6: Vr-Wr graph for siliqua per plant in *Brassica rapa*

The regression line intersected the Wr axis below the point of origin indicating the existence of over dominance gene action for controlling the trait (Fig. 6). The parent BARI sarisha 6 (P4), SAU sarisha 2 (P2), SAU sarisha 3 (P3) and Tori 7 (P1) contained maximum number of dominant alleles as it fell closer to the point of origin. The parent BARI sarisha 15 (P5) fell far from the origin indicating the presence of maximum frequency of recessive alleles in that parent. Chowdhury *et al.* (2004b) and Trivedi and Mukharjee (1986) observed over dominance in *Brassica rapa* and *Brassica juncea*, respectively.

4.4.6 Length of Siliqua




Fig. 7: Vr-Wr graph for length siliquae in *Brassica rapa*

The regression line intersected the Wr axis above the point of origin suggesting partial dominance gene action for controlling the trait (Fig. 7). The parents BARI sarisha 6 (P4) contained maximum dominant alleles as it fell closer to the point of origin. The parents SAU sarisha 3 (P3) and BARI sarisha 15 (P5) fallen at the nearly middle portion means they contained nearly equal frequencies of dominant and recessive alleles. The parent SAU sarisha 2 (P2) and Tori 7 (P1) fell far from the origin and thus it contained maximum frequency of recessive alleles. Trivedi and Mukharjee (1986) found over dominance in *Brassica juncea*.

4.4.7 Seeds per siliquae




Fig. 8: Vr-Wr graph for seed/siliquae in *Brassica rapa*

The regression line intersected the W_r axis below the point of origin suggesting over dominance gene action for controlling the trait (Fig. 8). The parent Tori 7 (P1), SAU sarisha 3 (P3) and BARI sarisha 6 (P4) contained maximum dominant alleles as it fell closer to the point of origin. The parents SAU sarisha 2 (P2) and BARI sarisha 15 (P5) fallen far from the origin indicating the presence of maximum frequency of recessive alleles in this parent. Chowdhury *et al.* (2004b) observed over dominance in *Brassica rapa*, Trivedi and Mukharjee (1986) found over dominance in *Brassica juncea*.

4.4.8 Seed yield per plant




Fig. 9: Vr-Wr graph for seed yield per plant in *Brassica rapa*

The regression line intersected the W_r axis above the point of origin suggesting partial dominance gene action for controlling the trait (Fig. 9). The parents SAU sarisha 2 (P2) fell closer to the point of origin indicating that it contained maximum dominant alleles. The parents BARI sarisha 6 (P4), Tori 7 (P1) and SAU sarisha 3 (P3) fallen at the middle portion means they contained equal frequencies of dominant and recessive alleles. The parent BARI sarisha 15 (P5) fell far from the origin indicated that they possessed maximum frequency of recessive alleles. Chowdhury *et al.* (2004b) obtained over dominance in *Brassica rapa*, Trivedi and Mukharjee (1986) observed over dominance in *Brassica juncea*.

4.4.9 Thousand seed weight




Fig. 10: Vr-Wr graph for 1000 seed weight in *Brassica rapa*

The regression line intersected the W_r axis above the point of origin suggesting partial dominance gene action for controlling the trait (Fig. 10). The distribution of array points indicated that two parents SAU sarisha 2 (P2) and Tori 7 (P1) contained most dominant alleles as they fell closer to the point of origin. The parents BARI sarisha 6 (P4) fallen at the middle portion suggesting it contained equal frequencies of dominant and recessive alleles. The parent SAU sarisha 3 (P3) and BARI sarisha 15 (P5) fell far from the origin indicated that they possessed maximum frequency of recessive alleles. Chowdhury *et al.* (2004b) found partial dominance in *Brassica rapa*, Trivedi and Mukharjee (1986) observed over dominance in *Brassica juncea*.

Regression line intersected the W_r -axis below the origin for all the characters except plant height, length of Siliqua, Seed yield per plant, 1000-seed weight indicating the presence of over dominance. Such serious inflation of dominance has been postulated by Hayman (1954) and Jinks (1955). A further support to the existence of pseudo-over dominance was visualized in the estimates of D and H components and relative magnitude of GCA and SCA in variance component analysis for these traits. This was supported by the earlier findings reported by Chowdhury *et al.* (2004b) in turnip rape, Trivedi and Mukharjee (1986) in Indian mustard. The over dominance might not be an index of real over-dominance at gene level, since particular combination of positive and negative genes or a complementary type of gene interaction of simply correlated gene distribution could have caused serious inflation in particular combinations of unidirectional dominance which might have resulted in over-estimation of partial dominance (Comstock and Robinson, 1952; Hayman, 1954; Grafius, 1959). The presence of over dominance in the various components of oil yield and seed yield in the present study has also been substantiated by the findings of Singh *et al.* (1970, 1971) in *B. campestris*, Rawat (1975) in *B. juncea*, and Trivedi and Mukharjee (1985) in *B. juncea*, Chowdhury *et al.* (2004b) in *B. rapa*.

As non-fixable variation was high for all the attributes except days to maturity plant height and 1000-seed weight, considerable improvements of these traits might be possible by transferring complementary gene into non-epistatic high-dominance crosses or by eliminating duplicate genes from some of high-dominance crosses. A study of epistatic components would be helpful in formulating an efficient breeding programme.

The results obtained from both Griffing and Hayman's analysis indicated the importance of both additive and dominance genetic variances, the later being more important to utilise simultaneously both additive and dominant genetic variances. The graphical analysis also indicated the genetic diversity within the parents.



Chapter 5

Summary and Conclusion

SUMMARY AND CONCLUSION

A five parents (Tori 7, SAU sarisha 2, SAU sarisha 3, BARI sarisha 6 and BARI sarisha 15) half diallel cross hybrids were evaluated for estimating the magnitude of combining ability and heterosis over mid parent and better parent effects.

Analysis of combining ability following Griffing's approach showed significant GCA and SCA variance for all the characters studied, indicating the role of both additive and non-additive components in the genetic system controlling these characters. Estimates of GCA effects for different characters suggested that parent BARI sarisha 6 was the best general combiner for plant height, early maturity, number of primary branches, number of Siliqua per plant, seed yield per plant and thousand seed weight. The parent BARI sarisha 15 was the best general combiner for number of primary branches per plant and number of seeds per siliqua. The parent Tori 7 was good general combiner for number of secondary branches per plant. The parent SAU sarisha 2 was the best for seed per siliqua.

The higher magnitude of GCA variance was observed than that of SCA variance for plant height, days to maturity, no. of primary branches/ plant, no. of secondary branches/ plant, no. of siliquae/ plant, seed yield/ plant and 1000 seed weight which indicated the preponderance of additive component in their expression.

The SCA estimates of various characters revealed that Tori 7 X BARI sarisha 15 was the best cross combination for plant height, no. of secondary branches per plant, and no. of siliquae per plant. The combination SAU sarisha 2 X BARI sarisha 15 was the best for days to early maturity, Siliqua length and seed per siliqua. The hybrid SAU sarisha 3 X BARI sarisha 15 was the best for no. of primary branches per plant and thousand seed weight. For seed yield per plant the cross BARI sarisha 6 X BARI sarisha 15 was the best.

The Vr-Wr graph indicate over dominance for days to maturity, primary branches per plant, secondary branches per plant, siliquae per plant, and seeds per siliqua,. Partial dominance was observed for plant height, length of siliqua, seed yield per plant and thousand seed weight. The graphical analysis also indicates wide genetic diversity among the parents.

Among the genotypes, the parents had high GCA effects and wide genetic diversity and hybrids had high heterotic value and SCA effect should give maximum emphasis. These genotypes could be effectively used in future for developing varieties of rapeseed (*Brassica rapa* L).

It was observed that all the hybrids obtained did not perform well for many of the important characters and to find out the desirable hybrids, the crosses were scored on the basis of desirable heterotic values. Out of ten crosses, the hybrids SAU sarisha 2 X BARI sarisha 6 showed desirable negative heterosis for the character of short plant height. The hybrids SAU sarisha 3 X BARI sarisha 6 were best for early maturity. The hybrids Tori 7 X BARI sarisha 15, SAU sarisha 3 X BARI sarisha 15, Tori 7 X SAU sarisha 2 were found to be heterotic for no. of primary branches per plant, no. of secondary branches per plant and no. of siliquae per plant. While the cross Tori 7 X BARI sarisha 15 produced maximum no. of secondary branches and no. of siliquae per plant. Most of the crosses showed significant and negative performance for length of siliqua and no. of seeds per siliqua. The crosses SAU sarisha 2 X BARI sarisha 15 were best for siliqua length and the crosses SAU sarisha 3 X BARI sarisha 6 and Tori 7 X BARI sarisha 6 were best for seeds per siliqua. For thousand seed weight the hybrids SAU sarisha 3 X BARI sarisha 15, Tori 7 X BARI sarisha 15, BARI sarisha 6 X BARI sarisha 15 and Tori 7 X SAU sarisha 2 were best. For seed yield per plant the cross SAU sarisha 2 X BARI sarisha 15 was found to be the best. Thus selection out of these crosses in the subsequent generations might produce some suitable segregants.



References

REFERENCES

- Adefris T. and C.B. Heiko. (2005). Heterosis and combining ability in a diallel cross of Ethiopian mustard inbred lines. *Crop Sci.* 45: 2629-2635.
- Agrawal, P.K. and S.S. Badwal. (1998). Possible utilization of commercial heterosis in Indian mustard *Brassica juncea* L. Cross and Czern, *Indian J. Genet. Pl. Breed.* 58 (4): 513-516.
- Ahmad, M.R. (1993). Study of agronomic value of resynthesized rapeseed lines and early generations of crosses rsyn-lines \times improved varieties. *Iranian J. Agril. Sci.* 24 (3/4): 1-13.
- Ali, M., L.O. Copeland, S.G. Elias and J.D. Kelly. (1995). Relationship between genetic distance and heterosis for yield and morphological traits in winter canola (*Brassica napus* L.). *Theor. Appl. Genet.* 91 (1): 118-121.
- Anonymous, (2000). FAO, Production Year Book. Food and Agricultural Organization of United Nations, Rome, Italy. 48:115.
- Arya, A.S., D.Singh, S. Hari and H. Singh. (1989). Unified selection procedure for simultaneous improvement of characters in rapeseed (*Brassica napus* L.). *Crop Res. Haisar.* 2(2): 137-141.
- Badwal, S.S. and K.S. Labana. (1987). Diallel analysis for some metric traits in Indian mustard. *Crop Improv.* 14 (2): 191-194.
- Banga, S.S. and K. S. Labana. (1985). Male sterility in Indian mustard (*Brassica juncea* L.) Cross. IV Genetics of MS-4. *Canadian J. Genet. Cytol.* 27: 487-490.
- Banga, S.S. and K.S. Labana. (1984). Heterosis in Indian mustard. *Zeitschrift für pflanzenzüchtung.* 92(1):61-70.
- Barua, P.K. and M.H. Hazarika. (1993). Heterosis analysis in Indian mustard rapeseed. In: heterosis breeding in crop plant – theory and application: short communications: symposium Ludhiana, April 5-5, Ludhiana, India. pp. 4-5.
- BBS. (2006a). Statistical year book of Bangladesh. (2006). Bangladesh Bureau of Statistics, Dhaka. Bangladesh.p.142.
- BBS. (2006b). Statistical year book of Bangladesh. (2006). Bangladesh Bureau of Statistics, Dhaka, Bangladesh.p. 151.

- BBS. (2006c). Statistical year book of Bangladesh. (2006). Bangladesh Bureau of Statistics, Dhaka, Bangladesh.p. 535.
- Chaudhury, B.D., A. Kumar, D.P. Singh, P.Singh and A. Kumar. (1987). Combining abilities and components of variation in *Brassica campestris*. *Res. Report*. 4 (2): 125-129.
- Chauhan, Y.S. (1987). Genetic analysis for yield and its components in Indian mustard. *Indian J. Agril. Res.* 21 (2): 77-82.
- Chauhan, Y.S., K. Kumar, B. Ram and S.K.Singh. (1990). Breeding for increased yield. Research on rapeseed and mustard. In: Proceedings of an Indo Swedish Symposium, 4-6 September, UP Saha, Sweden [Ed. Ohlsson, I.; Kumar, P.R.]: 109-121.
- Cheema, K.L. and H.A. Sadaqat. (2004). Potential and genetic basis of drought tolerance in canola (*Brassica napus*): II. Heterosis manifestation in some morpho-physiological traits in canola. *Int. J. Agric. Biol.* 6 (1): 82:85.
- Chowdhury, M.A.Z., M.A.K Mian., M.A. Akbar and M.Z. Alam. (2004). Combining ability for seed yield and yield contributing characters in Turnip Rape (*Brassica rapa L.*). *Bangladesh. J. Pl. Breed. Genet.* 17 (1): 17-24.
- Das, G.G., M.A. Quddus and M.E. Kabir. (2004). Heterosis in interspecific *Brassica* hybrids grown under saline condition. *J. Biol. Sci.* 4 (5): 664-667.
- Engqvist, G.M. and H.C. Becker. (1991). Heterosis and epistasis in rape seed estimated from generation means. *Euphytica*. 58 (1): 1165-1170.
- FAO. (2001). Production Year Book of 1999. Food and Agricultural Organization of United Nations, Rome, Italy.pp.118.
- FAO. (2003). Production Year Book. Food and Agricultural Organization of United Nations, Rome, Italy. 57:115-133.
- Ghosh, S.K., S.C.Gulati, R. Rajani and R. Raman. (2002). Combining ability and heterosis for seed yield and its components in Indian mustard (*Brassica juncea L.*). *Indian J. Genet. Pl. Breed.* 62 (1): 29-33.
- Goffman, F.D.and H.C. Becker. (2001). Diallel analysis for tocopherol contents in seeds of rape seed. *Crop Sci.* 41(4): 1072-1079.
- Goswami, P.K., N.K. Thakral, R.K. Behl and R. Kumar. (2004). Heterosis breeding in Indian mustard (*Brassica juncea L.*). *Brassica*. 6 (1/2): 47-51.

- Grant, I. and W.D. Beversdorf. (1985). Heterosis and combining ability estimates in spring plated oil seed rape (*Brassica napus*). *Canadian J. Genet. Cytol.* 27:472-478.
- Grants, I. (1985). Heterosis and cytoplasmic-genetic male sterility in oil seed rape (*Brassica juncea* L.). *Dissertation Abstract International (Science and Engineering)*. 46 (2):419.
- Griffing, B. (1956a). A generalized treatment of the use in diallel cross in quantitative inheritance. *Heredity*. 10: 13-50.
- Griffing, B. (1956b). Concept of general and specific combining ability in relation to diallel crossing systems. *Australian J. Biol. Sci.* 9: 463-493.
- Griffing, J.E. (1958). Application of sampling variance in the identification of methods which yields unbiased estimates of genotypic variances. *Australian J. Biol. Sci.* 11: 219-245.
- Gupta, M.L., K.S. Labana and S.S. Badwal. (1987a). Combining ability analysis of seed yield and its components in Indian mustard. In 71 International rapeseed congress, Poznan, Poland, 11-14 May.
- Gupta, M.L., S.K. Banga and G.S. Sandha. (1993). Commercially exploitable heterosis in *Brassica campestris* L. Oleiferous var. toria. In heterosis breeding in crop plant-theory and application: symposium Ludhiana, India. Crop improvement Society of India. pp. 18-19.
- Gupta, M.L., V.P. Gupta and K.S. Labana. (1987b). Combining ability analysis of seed yield and its components in Indian mustard. *Crop Improv.* 14 (2):160-164.
- Gupta, S.K. and K.S. Labana (1995). Genetics of plant nitrogen distribution in rapeseed. *J. Oilseeds Res.* 12 (2): 251-253.
- Habetinek, J. (1993). Heterosis and combining ability in a diallel cross of five varieties of winter swede rape (*Brassica napus* L.). *Roslinna Vyraba.* 55: 151-160.
- Hari, S., V.S. Malik and D. Singh. (1995). Heterosis in some intervarietal crosses of rapeseed (*Brassica napus*). *J. Oilseeds Res.* 12 (12): 180-183.
- Hayman, B.I. (1954a). The analysis of variance of diallel table. *Biometrics.* 10: 235-244.
- Hayman, B.I. (1954b). The theory and analysis of diallel crosses. *Genetics.* 39:789-809.

- Hayman, B.I. (1958). The separation of epistatic from additive and dominance Variation in generation means. *Heredity*. 12: 371-391.
- Hayman, B.I., (1957). Interaction, heterosis and diallel crosses. *Genetics*. 42: 336-355.
- Hirve, C.D. and A.S. Tiwari. (1991). Heterosis and inbreeding depression in Indian mustard. *Indian J. Genet.* 51 (2):190-193.
- Hoque, R. (2005). Combining ability and gene action for yield and yield contributing characters in Indian musard, derivatives. MS. Thesis, SAU, Dhaka.
- Huang, Y.J., J. Chen and Y.C. Li. (2000). Genetic study of *Sclerotinia sclerotiorum* resistance in rapeseed (*Brassica napus* L.) I. Its inheritance and combining ability. *Chinese J. Oil Crop Sci.* 22 (4): 1-5.
- Huq, K. M. E. (2007). Heterosis and combining ability in rapeseed (*Brassica rapa* L.). M.S. Thesis, Dept of Genetics and Plant Breeding, SAU, Dhaka.
- Iftikhar, A., S.A. Shah, S. Hassan, S.S.J. Ahmad, A. Mumtaz and K.U. Rehman. (2000). Mutant heterosis in oilseed rape. *Cruciferae Newsl.* 22: 67-68.
- Kakroo, P. and Kumar, S. (1991). Genetic determination of seed yield through its components in Indian mustard. *Indian J. Genet. Pl. Breed.* 51(2): 82.
- Katiyar R.K., R. Chamola and V.L. Chopra, (2000b). Heterosis and combining ability in Indian mustard (*Brassica juncea* L.). *Indian J. Genet. Pl. Breed.* 60 (4): 557-559.
- Katiyar, R.K., R. chamola and V.L. chopra. (2000a). Study of heterosis for seed yield in Indian mustard (*Brassica juncea* L.). *Cruciferae Newsl.* 22: 41-42.
- Katiyar, R.K., R. Chamola, H.B. Singh and S. Tickoo. (2004). Heterosis and combining ability analysis for seed yield in yellow sarson (*Brassica campestris*). *Brassica.* 6 (1/2): 23-28.
- Khaleque, M. A. (1985). A Guide book on Production of Oil Crops in Bangladesh. DAE and FAO/UNDP Project BGD/79/034, " Strengthening the Agricultural Extension Service" Khamarbari, Farmgate, Dhaka.p.3.
- Krishnapal, K. and S.K. Ghose. (1992). Heterosis in relation to genetic divergence in rapeseed and mustard. *J. Oilseed Res.* 9 (1): 169-174.

- Krzymanski, J. T. Pietka and K. Krotka. (1993). Combining ability and heterosis in diallel crosses of double-zero winter swede rape. I. F₁ generation. *Postepy Nauk Rolniczych*. 40/45(5): 41-52.
- Krzymanski, J., T. Pietka and M. Ogradowczyk. (1999). Combining ability of oilseed rape cultivars and strains and expected yield of synthetics from them. *Rosliny Oleiste*. 20(2): 325-334.
- Krzymanski, J., T. Pietka and K. Krotka. (1994). Combining ability and heterosis in diallel crosses of double-zero winter swede rape. II. F₁ and F₂ generations. *Rosliny Oleiste*. 15(1): 21-32.
- Krzymanski, J., T. Pietka, K. Krotka and K. Michalski. (1995). Glucosinolate content in F₁ hybrids of Polish double-zero winter swede rape. *Rosliny Oleiste*. 16 (1): 13-24.
- Krzymanski, J., T. Pietka, M. Ogradowczyk and K. Krotka. (1999). Estimation of combining ability for hybrids between winter oilseed rape inbred lines and varieties in a factorial design. I. F₁ generation. *Rosliny Oleiste*. 20(2): 335-345.
- Kudla, M. (1993). Comparative analysis of winter swede rape genotypes. *Biuletyn Instytutu Hodowli Roslin*. 90:99-107.
- Kudla, M. (1996). General combining ability of inbred lines and heterosis effects of winter oilseed rape F₁ and F₂ hybrids. *Rosliny Oleiste*. 17(1): 61-71.
- Kudla, M. (1997). General and specific combining ability of inbred lines and cultivars of winter oilseed rape. *Biuletyn Instytutu Hodowli Aklimatyzacji Roslin*. 201: 361-371.
- Kumar, C., S. Muralia, G. Deepak and D. Gupta. (2002). Heterosis studies in Indian mustard (*Brassica juncea* L.). *Cruciferae Newsl*. 24:59-60.
- Kumar, P.R., R.K. Arora, N.P. Singh, R.C. Yadav and P. Kumar. (1990). A study of heterosis in Indian mustard. *Acta Agronomica Hungarica*. 39(1-2):137-143.
- Lefort-Buson, M. and Y. Dattee. (1982). Genetic study of some agronomic characters in winter oilseed rape (*Brassica napus*). I. Heterosis, II. Genetic parameters. *Agronomic*. 2: 315-332.
- Lefort-Busson, M., Y. Dattee and B. Guillot-lemaine. (1987). Heterosis and genetic distance in rape seed (*Brassica napus* L.). *Genomes*. 29:11-18.

- Li, J. N., Qiu, J. and Tang, Z. L. (1990). Analysis of variability of some genetic parameters in segregating hybrid generations of *B. napus*. *Hereditas Beijing*. 11(6): 4-7.
- Liersch, A., I. Bartkowiak-Broda and K. Krotka. (1999). Characteristics of winter oilseed rape CMS ogura lines and their recurrent lines. *Rosliny Oleiste*. 20(2): 311-324.
- Liu, X.X., Z.S. Dong, C.S. Liu, J.G. Dong and H.B. Li. (2001). Combining ability and heritability of the main agronomic characters of some fine hybrids of rapeseed (*Brassica napus* L.). *China J. Oil Crop Sci*. 23 (3): 1-4.
- Lu, C.M., F.S. Shen, K. Hu. (2001). Heterosis in interspecific hybrids between *brassica napus* and *B.rapa*. *SABRAO J. Breed. Genet*. 33 (2): 73-85.
- Mahak, S. and S. Lallu. (2004). Heterosis in relation to combining ability for seed yield and its contributing traits in Indian mustard, *Brassica juncea* (L.). *J. Oilseeds Res*. 21(1): 140-142.
- Mahak, S., R.L. Srivastava, R.K. Dixit and M. Singh. (2003b). Heterosis and inbreeding depression in relation to seed yield in Indian mustard (*Brassica juncea* L.). *Pl. Arch*. 3 (1): 71-72.
- Mahak, S., S. Lallu, G.B. Swarnkar, P. Lalta and R.K. Dixit. (2003a). Heterobeltiosis and inbreeding depression in India mustard (*Brassica juncea* L.). *Crop Res. Hisar*. 26(2): 318-325.
- Malik, V., Singh, H. and Singh, D. (1995). Gene action and seed yield and other desirable characters in rapeseed. *Annal. Biol. (Ludhiana)*. 11(1/2):94-97.
- Malik, V.S., H. Singh and D. Singh. (1995). Gene action of seed yield and other desirable characters in rape seed (*Brassica napus* L.). *Annal. Biol*. 11:97.
- Matho, J.L. and Z.A. Haider. (2001). Assessing suitable combiners in *Brassica juncea* L. for high altitude acidic soils. *Cruciferae Newsl*. 23. pp. 47-48.
- Nair, B., V. Kalamkar and S. Bansod. (2005). Heterosis studies in mustard. *J. Phyto. Res*. 18 (2): 231-233.
- Nasim, M. (1990). Combining Ability and Heterosis in Cultivars of *Barassica campestris* L. M.sc. (Ag.) in Genetics and Plant Breeding Dissertation, submitted to the Faculty of Agriculture. Bangladesh Agril. University, Mymensingh.

- Nasrin, S. (2008). Heterosis and combining ability analysis in Indian mustard (*Brassica napus*). MS. Thesis, SAU, Dhaka.
- Pankaj, S., D.P. Pant, S.P. Singh, S. Chanchal, S. Ritesh, P. Sharma, C. Sah and R. Sharma. (2002). Heterosis vis a vis GCA of parents in toria (*Brassica campestris* L.). *Cruciferae Newsl.* 24: 71-72.
- Patel, M.C., J.D. Malkhandale and J.S. Raut. (1996). Combining ability in interspecific crosses of mustard (*Brassica* spp.). *J. Soils Crops.* 6(1): 49-54.
- Patil, S., V. Khambalkar, N.S. Solao and A. Thul. (2005). Study of heterosis in Indian mustard (*Brassica juncea*). *J. Phyto. Res.* 18 (1): 107-109.
- Piazza, G.J. and T.A. Foglia. (2001). Rapeseed oil for oleo chemical usage. *European J. Lip. Sci. Tech.* 103: 450-454.
- Pietka, T., K. Krotka and J. Krzymanski. (1998). Oilseed rape (*Brassica napus* L.) winter hardiness analysis with the use of diallel cross from generations F₁ and F₂. *Rosliny Oleiste.* 19(2): 371-378.
- Pietka, T., K. Krotka and J. Krzymanski. (2001). Study on combining ability in respect to glucosinolate content in F₁, and F₂ and generations of winter rapeseed hybrids (*Brassica juncea* L.). *Rosliny Oleiste.* 22 (2): 303-314.
- Pietka, T., K. Krotka and J. Krzymanski. (2003). General combining ability and heritability of glucosinolate content of winter oilseed rape (*Brassica napus* L.) seeds. *Biuletyn Instytutu Hodowli Aklimatyzacji Roslin.* (226/227): 405-414.
- Pradhan, A.K., Y.S. Sodhi, A. Mukhopadhaya and D.Pental. (1993). Heterosis breeding in Indian mustard (*Brassica juncea* L.): analysis of component characters contributing to heterosis for yield. *Euphytica.* 69 (3): 219-229.
- Prakash, N., Chauhan, Y.S., Kumar, K. (1987). Combining ability and path coefficient analysis in F₂ generation in Indian mustard (*Brassica juncea* L.) In 7th International Rapeseed Congress, 11-54 May, Poznan. Poland.p.110.
- Prasad, L., M. Singh and R.K. Dixit. (2002). Combining ability analysis in Indian mustard (*Brassica juncea* L.). *Adva. Pl. Sci.* 15 (1): 307-314
- Pszczola, J. (1993). Estimation of combining ability for selected traits in winter swede rape in two sets of diallel crosses. *Postepy Nauk Rolniczych.* 40/45(5): 33-40.

- Pu, H.M. (1998). Breeding of double low sterile line Ning A6 in rapeseed (*Brassica napus*) and its utilization. *Jiangsu J. Agril. Sci.* 14 (1): 8.
- Qi, C.K., G.J. Gao and J.F. Zhang. (2000). Analysis of heterosis among varieties or inbred lines in *Brassica campestris*. *Jiangsu J. Agril. Sci.* 11 (3): 30-32.
- Qi, C.K., H.M. Pu, J.F. Zhang and S.Z. Fu. (2003). Analysis of heterosis of seed yield and its components of intervarietal cross in *Brassica napus* L. *Jiangsu J. Agril. Sci.* 19 (3): 145-150.
- Qian, W., X. Chen, D. Fu, J. Zou and J. Meng. (2005). Inter subgenomic heterosis in seed yield potential observed in a new type of *Brassica napus* introgressed with partial *Brassica rapa* genome. *TAG. Theo. Appl. Genet.* 110(7): 1187-1194.
- Rai, B. and P.K. Singh. (1994). Expression of yield heterosis in some inter varietal crosses in Indian rapeseed (*Brassica campestris* L.). *Indian J. Genet. Pl. Breed.* 54 (3): 310-314.
- Ramsay, L.D., J.E. Bradshaw and M.J. Kearsey. (1994). The inheritance of quantitative traits in swedes (*Brassica napus* L. spp. rapifera). Diallel analysis of dry matter yield. *J. Genet. Breed.* 48 (3): 253-257.
- Rawat, D.S. (1987). Line \times tester analysis and rank index in mustard. *Ann. Agril.* 8 (2): 173-182.
- Rawat, D.S. (1992). Analysis of reciprocal differences in Indian mustard. *Acta Agron. Hungarica.* 41 (3-4): 277-233.
- Ripley, V.L. and W.D. Beversdorf. (2003). Development of self-incompatible *Brassica napus*:(1) introgression of S-alleles from *Brassica oleracea* through interspecific hybridization. *Pl. Breed.* 122 (1): 1-5.
- Satwinder, K., S. Paramjit and V.P. Gupta. (1997). Combining ability analysis for oil yield and its components in *Brassica napus* L. *Crop Improv.* 24 (1): 140-142.
- Satyendra, T., H.L. Singh and S. Mahak. (2004). Evaluation of heterosis for seed yield and its components Indian mustard (*Brassica juncea* L.) *Pl. Arch.* 4 (2): 433-437.
- Schuster, W., A. Alawi and R.G. Seidy. (1978). Studies on Breeding and heterosis phenomenon in black mustard (*Brassica nigra* L.). *Angewandte Botani.* 52 (3/4): 215-232.
- Sernyk, J. L. and B. R. Stefanson. (1983). Heterosis in summer rape (*Brassica napus*). *Canadian J. Pl. Sci.* 63:407-413.

- Shen, J.X., T.D. Fu, G.S. Yang, C.Z. Ma and J.X. Tu. (2005). Genetic analysis of rapeseed self-incompatibility lines reveals significant heterosis of different patterns for yield and oil content traits. *Pl. Breed.* 124 (2):111-114.
- Shi, H.Q., R.F. Cyong, L.L. Zhuan and X.P. Jhong. (1991). Studies on the utilization of heterosis in mustard (*Brassica juncea* L.). *Acta Agron. Sci.* 17 (1): 32-41.
- Siddique, S.B., M.A. Quddus, M.H. Rahman, M. Wazuddin, L. Rahman. (1990). Combining ability and heterosis in *Brassica campestris*. *Bangladesh J. Agril. Sci.* 17 (1): 141-146.
- Sing, D.N. and D.K. Verma. (1997). Scope of heterosis breeding in rapeseed-mustard. *J. Oilseeds Res.* 14 (2): 157-164.
- Singh, A.K., Y.S. Chauhan and K. Kumar. (1999). Combining ability analysis in rapeseed (*Brassica campestris* L.) using recessive genic male sterility. *J. Oilseeds Res.* 16 (2): 314-316.
- Singh, D., H. Singh, AS. Arya and Naveenchandra. (1992). Combining ability analysis and genetic improvement in Ethiopian mustard (*Brassica carinata* Braun). *Crop Res. Hisar.* 5 (2): 279-285.
- Singh, J.N. and B.R. Murty. (1980). Combining ability and maternal effects in *Brassica campestris* L. var. Yellow sarson. *Theor. Appl. Genet.* 56: 265-272.
- Singh, J.N., Maheshc, Yadav and I.A. Sheikh. (1996). Genetical studies for yield and oil content in *Brassica juncea* L. *Indian J. Genet.* 56 (3): 299-304.
- Singh, M., S.P. Singh and S. Dhirendra. (2000). Genetic analysis for seed yield and its genotypes in yellow sarson (*Brassica campestris* L.). *Indian J. Agril. Sci.* 70 (9): 624-626.
- Singh, O.N. and Y.S. Chauhan. (1987). Triple test cross analysis for seed and oil yield and its components in Indian mustard (*Brassica Juncea*. L.). In 7th International Rapeseed Congress, Poznan, 11-14 May, Poland, pp.111.
- Singh, R.S., P. Singh and R.K. Dixit. (1987). Combining ability analysis of yield and developmental traits in Indian colza (*Brassica campestris*. L. var. Yellow Sarson Prain). *Farm Sci. J.* 2(2): 170-174.
- Singh. V.S., A.N. Srivasta and Z. Ahmad. (1989). Combining ability in F₁ and F₂ generations of diallel cross in Indian mustard. *Crop Improv.* 16 (2): 164-167.

- Srivastava, K. and B. Rai. (1993). Expression of heterosis for yield and its attributes in rapeseed (*Brassica napus*.) *Indian J. Agri. Sci.* 63 (4): 243-245.
- Swarnkar, G.B., M. Singhand, L. Prasad. (2001). Combining ability for seed yield and its contributing characters in Indian mustard (*Brassica juncea*. L.). *J. Pl. Archives.* 2 (1): 5-14.
- Tak, G. M. and M. N. Khan. Combining ability for seed yield and its attributes in Indian Mustard (*Brassica Juneca* L.). *Adv. Pl. Sci.* 13 (2): 423-426.
- Tamber, K.S., M. Singh and M.L. Gupta. (1991). Combining ability for earliness in Indian mustard (*Brassica juncea*. L.). *Indian J. Genet. Pl. Breed.* 49:233-226.
- Thakur, H.L. and J.C. Segwal. (1997). Heterosis and combining ability in rapeseed (*Brassica napus* L.). *Indian J. Genet. Pl. Breed.* 57 (2): 163-167.
- Thakur, H.L. and M.A. Zarger. (1989). Heterosis in relation to genetic divergence and specific combining ability in Indian mustard *Brassica juncea*. *Indian J. Genet. Pl. Breed.* 49 (2): 223-226.
- Tyagi, M.K. J.S. Chauhan, P.R. Kumar and K.H. Singh. (2001). Estimaiton of heterosis in Indian mustard [*Brassica juncea*. (L)]. *Annl. Agril. Bio. Res.* 6 (2): 193-200.
- Tyagi, M.K., J.S. Chauhan, S.K. Yadav, P.R. Kumar and P. Tyagi. (2000). Heterosis in intervarietal crosses in mustard (*Brassica juncea*. L.) *Annl. Biol.* 16 (2): 191-194.
- Varshney, S.K. and C.S. Rao. (1997). Combining ability, heterosis and inbreeding depression for yield and yield components of in yellow sarson. *Indian J. Genet. Pl. Breed.* 57 (1): 91-97.
- Verma, N.K., B. Singh and J.N. Sachn. (1989). Combining ability and Heterosis in yellow sarson. *J. Oilseeds Res.* 6: 32-40.
- Verma, R.P. (2000). Combining ability analysis of yield and its components through diallel crosses in indica coiza (*Brassica juncea* L.). *Indian J. Agril. Res.* 34 (2):91-96.
- Walker, K.C. and E.J. Booth. (2001). Agricultural aspects of rape and other *Brassica* products. *European J. Lip. Sci. Tech.* 103: 441-446.
- Wang, W.R., H.J. Liu, G.H. Fang, H. Zhao, Y.L. Li, X.F. Qian and S.C. Cai. (1999). Analyses of heteroses and combining abilities of five rapeseed cultivars (lines) in *Brassica napus* L. *Acta Agril. Shanghai.* 15 (2): 45-50.

- Wos, H., B.I. Bartkowiak, G. Budzianowski and J. Krzymanski. (2000). Breeding works on winter and spring oilseed rape hybrids at Malyszyn. *Rosliny Oleiste*. 21 (3): 777-784.
- Wos, H., J. Krzymanski and M. Ogradowczyk. (1999). Estimation of combining ability of winter oilseed rape CMS ogura lines. *Rosliny Oleiste*. 20 (2): 347-355.
- Wos, H., W. Mackowiak and S. Wegrzyn. (1997). General combining ability of selected spring oilseed rape inbred lines. *Rosliny Oleiste*. 18 (1): 171-178.
- Wos, H., S. Wegrzyn and J. Wos. (1998). General combining ability of selected inbred lines and heterosis effects of winter oilseed rape F₁ hybrids. *Rosliny Oleiste*. 19 (2): 379-387.
- Woyke, T., (1981). The variability and mutual relationship of characters of significance in the evaluation of varieties and lines of winter Swede rape free of erucic acid. *Biuletyn Instytutu Hodowli i Aklimatyzacji Roslin*, 146: 17-23 (Pl. Br. Abstr., 53: 6641;1983).
- Wu, X.M., H.L. Huang., Z.L. Ning. and D.J. Ding. (2001). Studies on breeding and application of thermo-sensitive genic male sterile line 402S in *Brassica napus*. II. Evaluation of heterosis and combining ability. *Hunan Agril. Sci. Tech. Newsl.* 2 (4): 16-21.
- Yadav, L.S., T.P. Yadava, P. Kumar and N.K. Thakral. (1998). Heterosis for seed yield and its components in rapeseed under rainfed conditions. *Cruciferae Newsl.* 20: 79.
- Yadav, O.P., T.P. Yadava and P. Kumar. (1992). Combining ability studies for seed yield, its component characters and oil content in Indian mustard (*Brassica juncea* L.). *J. Oilseeds Res.* 9 (1): 14-20.
- Yadav, O.P., T.P. Yadava and P. Kumar. (1996). Combining ability studies for seed yield, its component characters and oil content in Indian mustard (*Brassica juncea* L.). *J. oilseeds Res.* 9 (1): 14-20.
- Yadav, T.P., P. Kumar, L. Raj and N.K. Thakral. (1997). A study of heterosis in toria (*Brassica campestris* var. toria). *Cruciferae Newsl.* 19: 101-102.
- Yadav, C.B., Y.S. Chauhan and K. Kumar. (2004). Heterosis for seed yield and its components in Indian mustard (*Brassica juncea* L.). *J. Maharashtra Agril. Uni.* 29 (2): 217-218.
- Yadav, C.B., Y.S. Chauhan and R.K. Chaudhury. (2005). Combining ability analysis using CMS systems in rape seed (*Brassica napus* L.). *Farm Sci. J.* 14 (1): 72-74.
- Yarnell, S. H. (1956). Cytogenetics of vegetable crops 11. *Crucifers Bot. Rev.* 22 (2): 18-166.

- Yu, X.L. and Z.J. Tang. (1995). Prediction of heterosis at the seedling stage of rape (*Brassica napus* L.) with acid phosphatase isozyme analysis. *J. Southwest Agril. Uni.* 17 (5): 399-403.
- Zhang *et al.* (2000). Genome analysis in *Brassica napus*. *Jiangsu J. Agril. Sci.* 11 (3): 30-32.
- Zheng, Y.J. and T.D. Fu. (1991). Study on heterosis in CMS line-derived hybrids of rapeseed (*Brassica napus* L.). *Oil Crops China* , 1: 7-11.
- Zuberi. M.I. and S.V. Ahmed. (1973). Genetic study of yield and some of its components in *Brassica campestris* var. Toria. *Crop Sci.* 13(1): 13-15.



Appendices

APPENDIX

Appendix I. Morphological, physical and chemical characteristics of initial soil (0-15 cm depth) of the experimental site

A. Physical composition of the soil

Soil separates	%	Methods employed
Sand	36.90	Hydrometer method (Day, 1915)
Silt	26.40	Do
Clay	36.66	Do
Texture class	Clay loam	Do

B. Chemical composition of the soil

Sl. No.	Soil characteristics	Analytical data	Methods employed
1	Organic carbon (%)	0.82	Walkley and Black, 1947
2	Total N (kg/ha)	1790.00	Bremner and Mulvaney, 1965
3	Total S (ppm)	225.00	Bardsley and Lanester, 1965
4	Total P (ppm)	840.00	Olsen and Sommers, 1982
5	Available N (kg/ha)	54.00	Bremner, 1965
6	Available P (kg/ha)	69.00	Olsen and Dean, 1965
7	Exchangeable K (kg/ha)	89.50	Pratt, 1965
8	Available S (ppm)	16.00	Hunter, 1984
9	pH (1 : 2.5 soil to water)	5.55	Jackson, 1958
10	CEC	11.23	Chapman, 1965

Source: Soil Resource and Development Institute (SRDI), Dhaka

REFERENCE ONLY

Appendix II. Monthly average temperature, relative humidity and total rainfall of the experimental site during the period from October, 2011 to April, 2012

Month	Air temperature (°c)		Relative humidity (%)	Rainfall (mm) (total)	Sunshine (hr)
	Maximum	Minimum			
October, 2011	34.8	18.0	77	227	5.8
November, 2011	32.3	16.3	69	0	7.9
December, 2011	29.0	13.0	79	0	3.9
January, 2012	28.1	11.1	72	1	5.7
February, 2012	33.9	12.2	55	1	8.7
March, 2012	34.6	16.5	67	45	7.3
April, 2012	35.8	20.3	65	88	8.3

Source: Bangladesh Metrological Department (Climate division), Agargaon, Dhaka-1212.



Sher-e-Bangla Agricultural University
 Lai
 Accession No. 38915
 Sign. Date: 3-9-15